

CHILDRENS ACETAMINOPHEN- acetaminophen suspension
Grocery Outlet, Inc.

Grocery Outlet Acetaminophen 160 mg/5 mL Oral Suspension Children's - Grape Flavor

Drug Facts

-

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:

-the common cold

-flu

-headache

-sore throat

-toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product,

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

.

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use.**
 - **do not give more than directed (see overdose warning)**
 - **shake well before using**
 - mL = milliliter
 - find right dose on chart below. If possible, use weight to dose; otherwise, use age.
 - remove the child protective cap and squeeze your child's dose into the dosing cup
 - repeat dose every 4 hours while symptoms last
 - do not give more than 5 times in 24 hours
-

Weight (lb)	Age (yr)	Dose (mL)**
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

**or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: **sodium 2 mg**
- store between 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments?

Call **1-877-290-4008**

Varnish Omit Area

Back of Package

Drug Facts (continued)

- Directions**
- This product does not contain directions or complete warnings for adult use.
 - Do not give more than directed (See overdose warning)
 - Shake well before using
 - mL = milliliter
 - Find right dose on chart below. If possible, use weight to dose; otherwise, use age.
 - Remove the child protective cap and squeeze your child's dose into the dosing cup
 - Repeat dose every 4 hours while symptoms last
 - Do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)**
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

** or as directed by a doctor
 Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- Each 5 mL contains: 300mg 2 mg
- Store between 20-25°C (68-77°F)

Inactive ingredients: anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments? Call 1-877-290-4008

TAMPER EVIDENT. DO NOT USE IF SHRINKBAND SEAL AROUND THE BOTTLE IS BROKEN OR MISSING

DISTRIBUTED BY:
 Grocery Outlet, Inc. | Emeryville, CA 94608
 www.groceryoutlet.com
 Made in India.
 922L 0326



IMPORTANT:
 Please see dosing chart or ask a doctor for your child's correct dose.

DIRECTIONS:

STEP 1
 Shake well before using.



STEP 2
 Remove child protective cap.



STEP 3
 Squeeze your child's dose into the dosing cup.



*This product is not manufactured or distributed by Kenova Inc., owner of the registered trademark Children's Tylenol® Oral Suspension

Top of Package

FOR AGES 2 TO 11 YEARS

Children's Acetaminophen

ORAL SUSPENSION ACETAMINOPHEN,
 160 mg / 5 mL
 Pain Reliever / Fever Reducer

Front of Package

NDC 83828-012-05

Compare to active ingredient in
 Children's Tylenol® Oral Suspension*

FOR AGES 2 TO 11 YEARS

Children's Acetaminophen

ORAL SUSPENSION ACETAMINOPHEN,
 160 mg / 5 mL
 Pain Reliever / Fever Reducer
 Ibuprofen, Aspirin, and Alcohol Free

GRAPE FLAVOR



DOSING CUP INCLUDED

4 FL OZ (118 mL)

GO

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each 5 mL) Purpose
 Acetaminophen 160 mg Pain reliever/fever reducer

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes
- more than 5 doses in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



LOT:
EXP.:

Print/Varnish Omit Area

Bottom of Package

Varnish Omit Area

Print/Varnish Omit Area

Drug Facts (continued)

overdose warning) ■ shake well before using ■ mL = milliliter
 ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ remove the child protective cap and squeeze your child's dose into the dosing cup. ■ repeat dose every 4 hours while symptoms last
 ■ do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)**
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

** or as directed by a doctor
 Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
Other Information ■ each 5 mL contains: sodium 2 mg ■ store between 20-25°C (68-77°F)
Inactive ingredients anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, sucrose, xanthan gum
Questions or comments? Call 1-877-290-4008
IMPORTANT: Please see dosing chart or ask a doctor for your child's correct dose.

DIRECTIONS:

STEP 1
Shake well before using.

STEP 2
Remove child protective cap.

STEP 3
Squeeze your child's dose into the dosing cup.

STOP PEELING

Adhesive Area

FOR AGES 2 TO 11 YEARS NDC 85828-922-05

Children's Acetaminophen

ORAL SUSPENSION
 ACETAMINOPHEN, 160 mg / 5 mL

Pain Reliever /
 Fever Reducer

GRAPE
 FLAVOR

GO

4 FL OZ (118 mL)

TAMPER EVIDENT: DO NOT USE IF SHRINKBAND SEAL AROUND THE BOTTLE IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each 5 mL)
 Acetaminophen 160 mg

Purpose
 Pain reliever/fever reducer

Uses
 temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.
 Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
 Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 Acetaminophen is allergic to ■ other inactive ingredients in this product
Ask a doctor before use if your child has liver disease
 Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin
When using this drug do not exceed recommended dose (see overdose warning)
 Stop use and ask a doctor if ■ pain gets worse or lasts more than 5 days ■ redness or swelling is present. These could be signs of a serious condition, keep out of reach of children. Overdose warning: in case of overdose, get medical help or contact a Poison Control Center right away (1-800-422-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions ■ this product does not contain directions or complete warnings for adult use.

Drug Facts (continued under label)

DISTRIBUTED BY:
 Grocery Outlet, Inc. | Emeryville, CA 94608
 www.groceryoutlet.com
 Made in India. 20210226

GM0096
100% MONEY BACK GUARANTEE
NO SUNK COSTS

LOT:
EXP.:

Peel Here

Varmish Omit Area

1/16" Margin

CHILDRENS ACETAMINOPHEN

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85828-922
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
XANTHAN GUM (UNII: TTV12P4NEE)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
SUCROSE (UNII: C151H8M554)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	purple	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85828-922-05	1 in 1 CARTON	03/10/2026	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/10/2026	

Labeler - Grocery Outlet, Inc. (029161585)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(85828-922)

