

ADAPALENE- adapalene gel
YYBA CORP

Adapalene New 73581-215

Active ingredient

Adapalene USP 0.1% (retinoid)*

*read consumer information leaflet

Purpose

Acne treatment

Use

For the treatment of acne

Warnings

For external use only

Do not use

on damaged skin (cuts, abrasions eczema, sunburn)

if you are allergic to adapalene or any of the ingredients in this product

Pregnancy/Breastfeeding

ask a doctor before use.

When using this product

limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors.

do not wax to remove hair in areas where the product has been applied

during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that become severe

irritation (redness, itching, dryness, burning) is more likely to occur:

- in the first few weeks of use
- if using more than one topical acne medication at a time
- but irritation usually lessens with continued use of this product

it may take up to 3 months of once daily use to see result

avoid product contact eyes, lips, and mouth. If contact occurs immediately flush the area with water.

wash hands after use

Stop use and ask a doctor if

you become pregnant, or planning to become pregnant while using this product

you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)

irritation become severe

you see no improvement after 3 months of once daily use

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

clean the skin gently and pat dry before applying the product

cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.

do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation.

Children under 12 years of age: ask a doctor

Other information

store at room temperature 68° to 77°F

protect from freezing

Inactive ingredients

carbomer homopolymer type C, edetate disodium, methylparaben, poloxamer 182, propylene glycol, purified water. May contain hydrochloric acid and/or sodium hydroxide to adjust pH.

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

†Compare to the active ingredient in Differin® Gel

Adapalene Gel USP 0.1%

Acne Treatment

- Previously available only by prescription
- Once daily topical retinoid*
- Oil free
- Fragrance free
- Dermatologist developed and tested

NET WT OZ (g)

***READ CONSUMER INFORMATION LEAFLET BEFORE USE.**

*Read carton and enclosed consumer information leaflet before using this product. Keep this carton and consumer information leaflet. They contain important information.

†This product is not manufactured or distributed by Galderma Laboratories, L.P. distributor of Differin® Gel.

DISTRIBUTED BY

Wellspring Meds

Airmont, NY 10952

Product Label

OIL FREE


FRAGRANCE FREE

PARABEN FREE

DERMATOLOGIST DEVELOPED AND TESTED

First FDA-approved over-the-counter topical retinoid for acne treatment*

CODE: GO/DRUGS/361

 Compare to the active ingredient in Differin® Gel¹

ADAPALENE GEL USP 0.1%

ACNE TREATMENT

ONCE DAILY TOPICAL RETINOID*

PREVIOUSLY AVAILABLE ONLY BY PRESCRIPTION

DERMATOLOGIST DEVELOPED AND TESTED

FDA APPROVED

 OIL FREE
 FRAGRANCE FREE

VALUE SIZE


1.6 OZ (45 g)
TUBE

PLD-B7964 F008363



3 73581 00020 3

Drug Facts

Active ingredient	Purpose
Adapalene USP 0.1% (retinoid)*	Acne treatment <small>*read consumer information leaflet</small>
Use	
<ul style="list-style-type: none"> • For the treatment of acne 	
Warnings	
For external use only	
Do not use	
<ul style="list-style-type: none"> • on damaged skin (cuts, abrasions, eczema, sunburn) • if you are allergic to adapalene or any of the ingredients in this product. 	
If pregnant or breast-feeding, ask a doctor before use.	
When using this product	
<ul style="list-style-type: none"> • limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors • do not wax to remove hair in areas where the product has been applied • during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that becomes severe • irritation (redness, itching, dryness, burning) is more likely to occur: <ul style="list-style-type: none"> • in the first few weeks of use • if using more than one topical acne medication at a time • but irritation usually lessens with continued use of this product • it may take up to 3 months of once daily use to see results • avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water. • wash hands after use 	
Stop use and ask doctor if	
<ul style="list-style-type: none"> • you become pregnant, or are planning to become pregnant, while using the product • you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath) • irritation becomes severe • you see no improvement after 3 months of once daily use 	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
Adults and children 12 years of age and older:	
<ul style="list-style-type: none"> • use once daily • clean the skin gently and pat dry before applying the product • cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face. • do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation. 	
Children under 12 years of age: ask a doctor	
Other information	
<ul style="list-style-type: none"> • TAMPER EVIDENT: DO NOT USE IF THE SEAL ON THE TUBE IS PUNCTURED OR NOT VISIBLE. • store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. • protect from freezing 	
Inactive ingredients	
carbomer homopolymer type C, edetate disodium, phenoxethanol, poloxamer 182, propylene glycol, purified water, sodium hydroxide.	
Questions? 1-866-933-6337	
<small>¹All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Differin® Gel.</small>	
	
Distributed by: Wellspring Airmont, NY 10952, U.S.A. 866-933-6337	
Made in India	

ADAPALENE

adapalene gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73581-215
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ADAPALENE (UNII: 1L4806J2QF) (ADAPALENE - UNII:1L4806J2QF)	ADAPALENE	1 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLOXAMER 182 (UNII: JX0HIX6OAG)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
CARBOMER HOMOPOLYMER TYPE C (UNII: 4Q93RCW27E)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73581-215-45	1 in 1 CARTON	12/10/2025	
1		45 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090962	12/10/2025	

Labeler - YYBA CORP (006339772)