NWRB NICOTINE TRANSDERMAL PATCH- nicotine patch Dongkou County Hecheng Network Technology Co., Ltd.

84288-007 NWRB Nicotine Transdermal Patch

Nicotine 21mg

Stop smoking aid

Use Reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings if you are pregnant or breast feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking However, the risks to your child from this medicine are not fully known.

Heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.

High blood pressure not controlled with medication. Nicotine can increase your blood pressure.

An allergy to adhesive tape or have skin problems, because you are more likely to get rashes

.Stomach ulcer or diabetes.

History of seizures

If you have vivid dreams or other sleep disturbances, remove the patch at bedtime.

Skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash.

Irregular heartbeat or palpitations occur.

You get symptoms of nicotine overdose, such as nausea, vomiting, dizziness weakness, and rapid heartbeat.

You have symptoms of an allergic reaction(such as difficulty breathing or rash).

Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away, Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

If you are under 18 years of age, ask a doctor before use.

Before using this product, read the enclosed User's Guide for complete directions and other information.

Use 1 patch every 24 hours. Apply to a clean, dry, hairless area of skin on the upper arm, hip, or torso.

Begin using the patch on your quit day, use according to the following 8-week schedule.

Polyacrylate-13 Polyester Film Lauryl Alcohol





FUNCTIONS:

• Reduces withdrawal symptoms, including nicotine craving.

• Helps you quit smoking.

Drug Facts

Active ingredient (in each patch) Purpose
Nicotine 21 mg / 24 hours Stop smoking aid
(Contains approximately 17% nicotine)

Use Reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings If you are pregnant or breast feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- Ask a doctor before use if you have
 Heart disease, recent heart attack, or irregular
 heartheat. Nicotine can increase your heart rate,
 High blood pressure not controlled with medication.
 Nicotine can increase your blood pressure.
 An allery to adhesive tape or have skin problems,
 because you are more likely to get rashes.
 Stomach utcer or diabetes.
 History of seizures.

Ask a doctor or pharmacist before use if you are

Using a non-nicotine stop smoking drug.
 Taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

If you have vivid dreams or other sleep disturbances, remove the patch at bedtime.

Drug Facts

Stop use and ask a doctor if

- Stop use and ask a doctor if
 Skin redness caused by the patch does not
 go away after four days, or if your skin swells
 or you get a rash.
 Irregular heartbeat or palpitations occur.
 Viou get symptoms of nicotine overdose,
 such as nausea, vomining, dizzness,
 "Wou pet symptoms of an altergic reaction
 (such as difficulty breathing or rash).

Keep out of reach of children and pets.

Used patches have enough nicotine to poison child and pets. If swallowed, get medical help or contact Poison Control Center right away. Save pouch to us for patch disposal Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- Directions

 If you are under 18 years of age, ask a
 doctor before use.

 Before using this product, read the enclosed User's
 Guide for complete directions and other information.

 Use 1 patch eveny 24 hours. Apply to a dean, day,
 hairless area of skin on the upper arm, hip, or torso.
- Begin using the patch on your quit day, use according to the following 8, used school do:

Amount of cigarettes /day	STEP 1		
Morethan 10	Weeks 1-4 Use one 21 mg patch/day	Weeks 5-6 Use one 14 mg patch/day	Weeks 7-8 Use one 7 mg patch/day
10 or less		Weeks 1-6 Use one 14 mg patch/day	Weeks 7-8 Use one 7 mg patch/day

Other information store at 20–25°C(68–77°F)

Inactive ingredients
Polyacrylate-13, Polyester Film, Lauryl Alcohol

MANUFACTURER: ShantouSouth VoonBiotechnology Co., Ltd.
ADDRESS: Room 114, 1st Floor, Building8-2,
(Shenzhen Shantou Digital Science and Technology Innovation Industrial Park)
No.22 Discoyun Road, Longhu District, ShantouCity,515000
SHIELF LIFE: 3 Years
MADE IN CHINA














Skin-Friendly

Start Your Quit Journey Today!





FUNCTIONS:

• Reduces withdrawal symptoms, including nicotine craving.

• Helps you quit smoking.

Drug Facts

Active ingredient (in each patch) Purposit
Nicotine 21 mg / 24 hours Stop smoking a
(Contains approximately 17% nicotine) Stop smoking air

Use Reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings If you are pregnant or breast feeding, only use this medicine on the advice of your health care provider. Smoking can senously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smokin However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- Heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
 High blood pressure not controlled with medication
- Ingin bood pressure not controlled with medication. Micotine can increase your blood pressure.

 An allergy to adhesive tape or have skin problems, because you are more likely to get rashes.

 Stomach ulcer or diabetes.

 History of seizures.

Ask a doctor or pharmacist before use if you are

Using a non-nicotine stop smoking drug.
 Taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

If you have vivid dreams or other sleep disturbances remove the patch at bedtime.

Drug Facts

Stop use and ask a doctor if

- Stop use and ask a doctor if
 Skin redness caused by the patch does not
 go away after four days, or if your skin swells
 or you get a rash.
 Irregular heartbeat or palpitations occur.
 Viou get symptoms of nicotine overdose,
 such as nausea, vomiting, dizzness,
 weakness, and raph fearetbeat.
 Viou have symptoms of an allergic reaction
 such as difficulty breathing or rash).

Keep out of reach of children and pets.

Used patches have enough nicotine to poison children and pets. It swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal Dispose of the used patches by folding sti

Directions

- Directions

 If you are under 18 years of age, ask a
 doctor before use.
 Before using this product, read the enclosed User's
 Guide for complete directions and other information.
 Use 1 patch every 24 hours. Apply to a clean, dry,
 haitess area of skin or the upper arm, hip, or torso.

- Begin using the patch on your quit day, use according to the following 8—week schedule:

STEP 1		
Weeks 1-4 Use one 21 mg	Weeks 5-6 Use one 14 mg patch/day	Weeks 7-8 Use one 7 mg patch/day

Other information store at 20-25°C(68-77°F)

Inactive ingredients Polyacrylate-13, Polyester Film, Lauryl Alcohol

MANUFACTURER: ShantouSouth VoonBiotechnology Co., Ltd.
ADDRESS: Room 114, 1st Floor, Building8—2,
(Shenzhen Shantou Digital Science and Technology Innovation Industrial Park)
No.22 Olacyun Road, Longhu District, ShantouCity,515000
SHLET LIFE: 3 Years
MADE IN CHINA

36M





NARE

NARO **NICOTINE** TRANSDERMA PATCH Reduces Nicotine Cravings & Withdrawal Symptoms

Drug Facts

Active ingredient (in each patch) Nicotine 21 mg / 24 hours (Contains approximately 17% nicotine) Stop smoking aid

Use Reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings If you are pregnant or breast feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any incotine replacement medicine. This medicine is believed to be safer than smokin ver, the risks to your child from this medicine

are not fully known. Ask a doctor before use if you have

Drug Facts

Stop use and ask a doctor if

Stop use and ask a doctor if
Skin redness caused by the patch does not
go away after four days, or if your skin swells
or you get a rash.
Irregular heartbeat or palpitations occur.
Vion get symptoms of incotine overdose,
such as nausea, vomiting, dizzness,
wedkness, and rapid heartbeat
vedkness, and rapid fleartbeat
such as difficulty breathing or rash).

Keep out of reach of children and pets.

Used patches have enough nicotine to poison childre and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.



MANUFACTURER: StantouSouth VoonBiotechnology Co., Ltd.
ADDRESS: Room 114, 134 Floor, Building8-2,
(Shenzhen Sharbou Digital Science andTechnology Innovation Industrial Park)
No 22 Glacytun Road, Longhur District, Shantou





NWRB NICOTINE TRANSDERMAL PATCH

nicotine patch

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:84288-007

Route of Administration TRANS DERMAL

Active Ingredient/Active Moiety

Ingredient Name

Basis of Strength

NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)

NICOTINE

21 mg

Inactive Ingredients			
Ingredient Name	Strength		
LAURYL ALCOHOL (UNII: 178A96NLP2)			
POLYETHYLENE TEREPHTHALATE (INTRINSIC VISCOSITY 0.60-0.70) (UNII: 5YSH70HE6K)			
POLYACRYLATE-13 (UNII: FS2D4T67EA)			

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		NDC:84288-007- 01	30 in 1 BOTTLE; Type 0: Not a Combination Product	12/02/2025	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	505G(a)(3)	12/02/2025		

Labeler - Dongkou County Hecheng Network Technology Co., Ltd. (637818093)

Establishment					
Name	Address	ID/FEI	Business Operations		
Dongkou County Hecheng Network Technology Co., Ltd.		637818093	manufacture(84288-007)		

Revised: 12/2025 Dongkou County Hecheng Network Technology Co., Ltd.