

**RIGHT REMEDIES PAIN RELIEF CAPLET- acetaminophen tablet, coated**  
**Strive Pharmaceuticals Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**RIGHT REMEDIES Pain Relief Caplet**

**Drug Facts**

**Active Ingredient (in each caplet)**

Acetaminophen 500mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever.

**Warnings**

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug

warfarin.

**Stop use and ask a doctor if** • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children. OVERDOSE WARNING:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

### ***Directions***

Do not take more than directed. **SEE OVERDOSE WARNING.**

adults & children 12 years of age and over	<ul style="list-style-type: none"><li>• take 2 caplets every 6 hours while symptoms last</li><li>• do not take more than 6 caplets in 24 hours, unless directed by doctor</li><li>• do not take more than 10 days, unless directed by a doctor</li></ul>
children under 12 years of age	<ul style="list-style-type: none"><li>• ask a doctor</li></ul>

### ***Other information***

- store at room temperature between 20 - 25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.

**TAMPER EVIDENT: Do not use if imprinted safety seal is broken or missing.**

### ***Inactive ingredients***

hydroxypropyl methylcellulose, maltodextrin, microcrystalline cellulose, polyethylene glycol, povidone, sodium starch glycolate, starch, stearic acid

### ***Questions or comments?***

Call 1 (888) 577 - 8033

†compare to active ingredient in **TYLENOL®** Extra Strength Caplets

**EXTRA STRENGTH**

**Distributed by: STRIVE PHARMACEUTICALS INC.  
East Brunswick, NJ 08816**

# READ AND KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

†This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division., owner of the registered trademark Tylenol® Extra Strength Caplets.

## Packaging

**RIGHT REMEDIES**

**Pain Relief**

† pain reliever/fever reducer  
**EXTRA STRENGTH**

acetaminophen  
**500mg** each caplet

**24** caplets

**RIGHT REMEDIES**

**Pain Relief**

† pain reliever/fever reducer  
**EXTRA STRENGTH**

acetaminophen  
**500mg** each caplet

**24** caplets

**RIGHT REMEDIES**

**Pain Relief**

† pain reliever/fever reducer  
acetaminophen 500mg

**READ AND KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

†This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division., owner of the registered trademark Tylenol® Extra Strength Caplets.

†compare to active ingredient in  
**TYLENOL®** Extra Strength Caplets  
MDC 70692-141-42

†compare to active ingredient in  
**TYLENOL®** Extra Strength Caplets  
MDC 70692-141-42

Distributed by: STRIVE PHARMACEUTICALS INC.  
East Brunswick, NJ 08816  
CT7069214142 - REV00-042020

**3 70692 14042 2**

Lot. No.:  
Exp. Date:

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. **OVERDOSE WARNING:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**  
Do not take more than directed. **SEE OVERDOSE WARNING**

adults & children 12 years of age and over	<ul style="list-style-type: none"> <li>take 2 caplets every 6 hours while symptoms last</li> <li>do not take more than 6 caplets in 24 hours, unless directed by doctor</li> <li>do not take more than 10 days, unless directed by a doctor</li> </ul>
children under 12 years of age	ask a doctor

**Other information**

- store at room temperature between 20°-25°C (68°-77°F); avoid excessive heat, cold and humidity.
- close caplet(s) after use.
- TAMPER EVIDENT: Do not use if imprinted safety seal is broken or missing.**

**Inactive ingredients**  
hydroxypropyl methylcellulose, maltodextrin, microcrystalline cellulose, polyethylene glycol, povidone, sodium starch glycolate, starch, stearic acid

**Questions or comments?** Call 1 (888) 577-8833

**Drug Facts**

**Active Ingredient (in each caplet)**  
Acetaminophen 500mg Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - periodic and menstrual cramps
- temporarily reduces fever.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

**Stop use and ask a doctor if:**

- diarrhea occurs or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- itching or swelling is present. These could be signs of a serious condition.

## DRUG FACTS LABEL



## Drug Facts

Active Ingredient (in each caplet)	Purpose
Acetaminophen 500mg.....	Pain reliever/fever reducer

### Uses

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever.

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease  
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if 

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

 These could be signs of a serious condition.

## Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. **OVERDOSE WARNING:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

### Directions

Do not take more than directed. SEE OVERDOSE WARNING.

adults & children 12 years of age and over	<ul style="list-style-type: none"><li>take 2 caplets every 6 hours while symptoms last</li><li>do not take more than 6 caplets in 24 hours, unless directed by doctor</li><li>do not take more than 10 days, unless directed by a doctor</li></ul>
children under 12 years of age	<ul style="list-style-type: none"><li>ask a doctor</li></ul>

### Other information

- store at room temperature between 20 - 25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.

**TAMPER EVIDENT: Do not use if imprinted safety seal is broken or missing.**

### Inactive ingredients

hydroxypropyl methylcellulose, maltodextrin, microcrystalline cellulose, polyethylene glycol, povidone, sodium starch glycolate, starch, stearic acid

**Questions or comments?** Call 1 (888) 577-8033

## RIGHT REMEDIES PAIN RELIEF CAPLET

acetaminophen tablet, coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-141
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	capsule	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	S500
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-141-42	24 in 1 PACKAGE; Type 0: Not a Combination Product	04/01/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/01/2020	

**Labeler** - Strive Pharmaceuticals Inc. (080028013)

Revised: 7/2022

Strive Pharmaceuticals Inc.