

**CHLORAPREP- chloraprep solution**  
**HF Acquisition Co LLC, DBA HealthFirst**

-----  
Chlorhexidine gluconate (CHG), 2% w/v  
and Isopropyl alcohol (IPA), 70% v/v

Antiseptic

for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

For external use only. Flammable, keep away from fire or flame.

do not use with electrocautery procedures

**Allegery Alert**

This product may cause a severe allergic reaction.

Symptoms may include:

wheezing/difficulty breathing  
shock  
facial swelling  
hives  
rash

If an allergic reaction occurs, stop use and seek medical help right away.

on patients allergic to chlorhexidine gluconate or any other ingredient in this product  
for lumbar puncture or in contact with the meninges  
on open skin wounds or as a general skin cleanser

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

If swallowed, get medical help or contact a Poison Control Center right away.

use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

maximal treatment area for one applicator

one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm<sup>2</sup>)

tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip.

Place foam flat side down on the treatment area.

completely wet the treatment area with antiseptic

dry surgical sites (e.g., abdomen or arm): use repeated back-and-forth strokes for 30 seconds. (When using the triple swabstick applicators, use each swabstick sequentially

within the 30 seconds). Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away.

moist surgical sites (e.g., inguinal fold): use repeated back-and-forth strokes for 2 minutes. (When using the triple swabstick applicators, use each swabstick sequentially within the 2 minutes). Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.

discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

USP purified water

store between 15-30 °C (59-86 °F)

avoid freezing and excessive heat above 40 °C (104 °F)



## CHLORAPREP

chlora prep solution

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51662-1675(NDC:54365-401)
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL
<b>CHLORHEXIDINE GLUCONATE</b> (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	20 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51662-1675-1	1 in 1 POUCH	05/13/2024	
1		1.75 mL in 1 APPLICATOR; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021555	05/13/2024	

Labeler - HF Acquisition Co LLC, DBA HealthFirst (045657305)

Registrant - HF Acquisition Co LLC, DBA HealthFirst (045657305)

Establishment

Name	Address	ID/FEI	Business Operations
HF Acquisition Co LLC, DBA HealthFirst		045657305	repack(51662-1675)