THERMACARE MAX STRENGTH LIDOCAINE- thermacare max strength lidocaine spray Clarion Brands LLC

Benzyl alcohol 10%

Lidocaine HCl 4%

Topical anesthetic

Use

• for the termporary relief of pain

For external use only

Flammable

- contents under pressure
- keep away from fire, sparks, and heated surfaces
- do not puncture or incinerate
- exposure to temperatures above 130°F may cause bursting

Do not use

- in large quantities
- on wounds, damaged, broken, or irritated skin
- for more than one week without consulting a doctor

When using this product

- use only as directed. Read and follow all directions and warnings on this label
- avoid contact with the eyes and mucous membranes
- do not bandage tightly or apply local heat such as a heating pad or a medicated patch
- do not use with other topical analgesics

Stop use and ask a doctor if

- rash or irritation develops
- symptoms worsen or persist for more than 7 days or clear up and occur again within a few days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

- adults and children 2 years of age and older: apply to the affected area no more than 3 to 4 times daily
- children under 2 years of age: consult a doctor before use

1.2 hexanediol

alcohol

aloe barbadensis leaf juice

aminomethyl propanol

bisabolol

caprylyl glycol

disodium EDTA

isopropyl alcohol

propylene glycol

vanillyl butyl ether

water

zingiber officinale (ginger) root extract

1-800-323-3383

Benzyl alcohol 10%

Lidocaine HCI 4%

Directions

adults and children 2 years of age and older: apply to affected area no more than 3 to 4 times daily

children under 2 years of age: consult a doctor before use.

NEW! ThermaCare Heat Therapy

Soothing Heat 2x Dual Action Maximum Lidocaine

Heat + Max Strength Lidocaine Pain Relief

Fast Acting, Long Lasting

Penetrates to the Source

Number 1 Doctor Recommended

Spray NET WT 3oz (85g)

No Mess Application - 360 Degree Spray

Distributed by: Bridges Consumer Healthcare

811 Broad Street, Suite 600, Chattanooga, TN 37402



Drug Facts

Active ingredients

Purpose

Benzyl alcohol 10% Lidocaine HCl 4% Topical anesthetic Topical anesthetic

Use ■ for the temporary relief of pain

Warnings For external use only

Flammable Contents under pressure

- keep away from fire, sparks, and heated surfaces
- do not puncture or incinerate
- exposure to temperatures above 130°F may cause bursting

Do not use in large quantities

- on wounds, damaged, broken, or irritated skin
 for more than one week without consulting a doctor

- When using this product
 use only as directed, Read and follow all directions and warnings on this label avoid contact with the eyes and mucous membranes
- do not bandage tightly or apply local heat such as a heating pad or a medicated patch - do not use with other topical analgesics

Stop use and ask a doctor if a rash or imitation develops

symptoms worsen or persist for more than 7 days or clear up and occur again within a few days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children, if swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 2 years of age and older: apply to the affected area no more than 3 to 4 times daily
- m children under 2 years of age: consult a doctor before use

Inactive ingredients 1,2-hexanediol, alcohol, alce barbadensis leaf juice, aminomethyl propanol, bisabolol, caprylyl glycol, disodium EDTA, isopropyl alcohol, propylene glycol, vanillyl butyl ether, water, zingiber officinale (ginger) root extract

Questions? call 1-800-323-3383

AVOID INHALING OVERSPRAY WHILE USING

Distributed by: Bridges Consumer Healthcare 811 Broad Street, Suite 600, Chattanooga, TN 37402 ©2026

*ProClaim Solutions 2025 prescription

1000315



THERMACARE MAX STRENGTH LIDOCAINE

thermacare max strength lidocaine spray

Product Information

Product Type HUMAN OTC DRUG **Item Code (Source)** NDC:69693-004

TOPICAL Route of Administration

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------------------|-------------------|
| LIDOCAINE HCL (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987) | LIDOCAINE HYDROCHLORIDE ANHYDROUS | 4 g in 100 mL |
| BENZYL ALCOHOL (UNII: LKG8494WBH) (BENZYL ALCOHOL - UNII: LKG8494WBH) | BENZYL ALCOHOL | 10 g in 100 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| GINGER (UNII: C5529G5JPQ) | |
| AMINOMETHYL PROPANOL (UNII: LU49E6626Q) | |
| VANILLYL BUTYL ETHER (UNII: S2ULN37C9R) | |
| EDETATE DISODIUM (UNII: 7FLD91C86K) | |
| 1,2-HEXANEDIOL (UNII: TR046Y3K1G) | |
| ALCOHOL (UNII: 3K9958V90M) | |
| ISOPROPYL ALCOHOL (UNII: ND2M416302) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) | |
| WATER (UNII: 059QF0KO0R) | |
| ALOE BARBADENSIS LEAF JUICE (UNII: RUE8E6T4NB) | |
| BISABOLOL (UNII: 24WE03BX2T) | |

| Product Characteristics | | |
|-------------------------|-------|--------------|
| Color | | Score |
| Shape | ROUND | Size |
| Flavor | | Imprint Code |
| Contains | | |

| l | P | Packaging | | | | |
|---|---|-----------|---|-------------------------|-----------------------|--|
| | # | Item Code | Package Description | Marketing Start Date | Marketing End Date | |
| | | | 1 mL in 1 CANISTER; Type 0: Not a Combination Product | 12/02/2025 | | |

| Marketing In | Marketing Information | | | |
|-----------------------|---|-------------------------|-----------------------|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| OTC Monograph Drug | M017 | 12/02/2025 | | |
| | | | | |

Labeler - Clarion Brands LLC (079742703)

Revised: 12/2025 Clarion Brands LLC