

**VICKS NYQUIL HOT REMEDY COLD AND FLU- acetaminophen,
diphenhydramine hcl powder, for solution
Procter & Gamble Manufactura S de RL de CV**

**Vicks[®] NyQuil[™] HOT REMEDY
COLD & FLU**

Drug Facts

Active Ingredient (in each Packet)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Purpose

Pain reliever/fever reducer

Antihistamine, Cough suppressant

Uses

temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor sore throat & bronchial irritation
- cough to help you sleep
- runny nose
- sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (Certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

Age	Dose
Adults & children 12 yrs & over	one packet (dose) every 4 hours
Children under 12 yrs	Do not use

Other information

- **each packet contains:** potassium 10 mg
- phenylketonurics: contains phenylalanine 68 mg per dose
- store at no greater than 25°C (77°F)

Inactive ingredients

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF SACHET IS OPENED OR TORN

Keep outer package for complete product information.

MADE IN MEXICO

**DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202**

PRINCIPAL DISPLAY PANEL - 8 packet package

NEW FORMULA

Vicks® NyQuil™ HOT REMEDY

COLD & FLU

Acetaminophen,

Diphenhydramine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough, Sneezing, Runny Nose

HONEY LEMON FLAVORED

Nighttime Relief

8 PACKETS



Drug Facts (continued)	
• headache • cough due to minor throat & bronchial irritation • cough to help you sleep • runny nose • sneezing	
Warnings	
Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take	
• more than 6 doses in 24 hours, which is the maximum daily amount for this product	
• with other drugs containing acetaminophen	
• 3 or more alcoholic drinks every day while using this product	
Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash	
If a skin reaction occurs, stop use and seek medical help right away	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use	
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
• with any other product containing diphenhydramine, even one used on the skin.	
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have • liver disease • glaucoma • cough that occurs with too much phlegm (mucus)	
• a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema	
• trouble urinating due to enlarged prostate gland	
Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin	
When using this product • excitability may occur, especially in children • marked drowsiness may occur	
• alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks	
• be careful when driving a motor vehicle or operating machinery	
Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days	
• redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts	
These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions • take only as directed • do not exceed 6 doses per 24 hrs	
• Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes	
• If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat	
Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use
Other information • each packet contains: potassium 10 mg	
• phenylephrine: contains phenylephrine 68 mg per dose • store at no greater than 25°C (77°F)	
Inactive ingredients acacia, ascorbic acid, aspartame, carboxymethylcellulose, citric acid, D&G Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate	
Questions? 1-800-362-1683	



VICKS NYQUIL HOT REMEDY COLD AND FLU

acetaminophen, diphenhydramine hcl powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58933-555
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
TRICALCIUM PHOSPHATE (UNII: K4C08XP666)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ASPARTAME (UNII: Z0H242BBR1)	
SUCROSE (UNII: C151H8M554)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
CARBOXYMETHYLCELLULOSE (UNII: 05JZI7B19X)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor	HONEY (Honey Lemon)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-555-01	1 in 1 PACKET; Type 0: Not a Combination Product	06/11/2025	
2	NDC:58933-555-08	8 in 1 PACKAGE	06/11/2025	
2		1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/11/2025	

Labeler - Procter & Gamble Manufactura S de RL de CV (812807550)

Revised: 12/2025

Procter & Gamble Manufactura S de RL de CV