

## **PINWORM TREATMENT- pyrantel pamoate suspension**

### **STRATEGIC SOURCING SERVICES**

-----

#### **pinworm treatment-foster & thrive**

#### **Active ingredient (in each mL)**

Pyrantel Pamoate 144mg/mL

(the equivalent of 50mg pyrantel base)

#### **Purpose**

Anthelmintic

#### **Use**

for the treatment of pinworms

#### **Warnings**

**Do not exceed recommended dosage**

#### **Ask a doctor before use if**

■ you are pregnant ■ have liver disease ■ nursing a baby

#### **When using this product**

abdominal cramps, nausea, vomiting, diarrhea, headache, or dizziness sometimes occur after taking this drug. If any of these symptoms persist, consult a doctor.

#### **Keep out of reach of children.**

In case of overdose get medical help or contact a poison control center right away.

#### **Directions**

■ **shake well before use**

■ **read bottle label and package insert carefully before taking this medication**

■ treat the entire household unless otherwise advised. See Warnings

■ do not repeat treatment unless directed by a doctor

■ this product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during or after medication

■ if signs of pinworms persist after treatment, consult a doctor

---

**Weight**

Less than 25 lb. or under 2 yrs.old  
25-37 lb.  
38-62 lb.  
63-87 lb.  
88-112 lb.  
113-137 lb.  
138-162 lb.  
163-187 lb.  
188 lb. and over

**Dosage**

(taken as a single dose)

Do not use unless directed by a doctor  
1/2 teaspoonful  
1 teaspoonful  
1 1/2 teaspoonfuls  
2 teaspoonfuls  
2 1/2 teaspoonfuls  
3 teaspoonfuls  
3 1/2 teaspoonfuls  
4 teaspoonfuls

---

**Other information**

- each teaspoon contains **3mg of sodium**
- Store between 15°C-30°C (59°- 86°F)

**Inactive ingredients**

banana flavor, citric acid, glycerin, lecithin, magnesium aluminum silicate, methylcellulose, methylparaben, povidone, propylene glycol, propylparaben, purified water, simethicone, sodium benzoate, sodium citrate, sodium saccharin and sorbitol.

**Questions or comments?**

Call 8 **33-358-6431** Monday to Friday  
9:00am to 7:00pm EST

©2023 Northstar Healthcan UC

**Distributed by:**

McKesson Corp~  
viaStrategicSourcingServicesll.C.  
Memphis, TN 381--41  
MoneyBackGuarantee  
www.fosterandthrive.com

**Product of India**

**Code: TN/Drugs/TN00002222/2006  
949129350**

**L0001161**

**Rev. 09/25**

**READ THIS INSERT CAREFULLY BEFORE TAKING MEDICATION**

**PINWORM MEDICINE**

Pyrantel Pamoate 144 mg/mL

(the equivalent of 50 mg Pyrantel base per mL)

## **FOR THE TREATMENT OF PINWORMS**

### **DIRECTIONS**

Adults, children 12 years of age and over, and children 2 years to under 12 years of age: oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight not to exceed 1 gram.

Take only according to directions and do not exceed the recommended dosage unless directed by a doctor. Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See warnings. If any worms other than pinworms are present before or after treatment, consult a doctor. If any symptoms or pinworms are still present after treatment, consult a doctor.

**Shake well before using.**

<b>Weight</b>	<b>Dosage (taken as a single dose)</b>
Less than 25 pounds or under 2 years old	consult a doctor
25 to 37 pounds	½ teaspoonful
38 to 62 pounds	1 teaspoonful
63 to 87 pounds	1½ teaspoonfuls
88 to 112 pounds	2 teaspoonfuls
113 to 137 pounds	2½ teaspoonfuls
138 to 162 pounds	3 teaspoonfuls
163 to 187 pounds	3½ teaspoonfuls
188 pounds and over	4 teaspoonfuls

This product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during, or after medication.

### **WARNINGS**

Abdominal cramps, nausea, vomiting, diarrhea, headaches, or dizziness, sometimes occur after taking this drug, if any of these conditions persist consult a doctor.

If you are pregnant, nursing or have liver disease, do not take this product unless directed by your doctor.

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

### **OTHER INFORMATION**

This brochure will tell you how to identify and treat pinworm infections effectively with pinworm medicine. Many who have pinworms do not have symptoms, however, many will have itching in and around the rectal opening. The itching may be very annoying and continued scratching can cause an irritation in this area. The itching may be prominent

at night when sleeping. That is why restless sleep is sometimes a sign of pinworms, especially in children. Other symptoms include insomnia, gastrointestinal distress, irritability, enuresis (bed wetting) and secondary infection due to localized scratching. You should treat with pinworm medicine only after you have seen and identified the pinworms.

Pinworms look like tiny white threads. The female pinworm is about 1/2 inch long (the male pinworm is shorter); they live in the bowel. Usually, at night, the female pinworm travels to the rectal opening and lays eggs within the skin folds around the opening. This usually takes place within 1-2 hours after the child has been put to sleep for the night. The sticky gelatin like substance in which the eggs (they are too small to see) are deposited and the movement of the female pinworm may cause annoying itching and possibly restless sleep. By checking the rectal opening at this time, you can attempt to see the pinworm as described.

Scratching will cause pinworm eggs to stick to the fingers. Reinfection will result if the fingers are placed in the mouth. The eggs, which are too small to see, contaminate whatever they come in contact with: bed clothes, underwear, toys, hands, and food touched by contaminated hands. Even eggs floating in the air can be swallowed and cause infection. Eggs deposited can survive for as long as 3 weeks. Pinworms are highly contagious; therefore, after treatment with pinworm medication you should wash hands and fingernails with soap often during the day in order to prevent spreading to others and reinfection. Wear tight underpants both day and night. For several days after treatment clean the bedroom floor by vacuuming or damp mopping. After treatment, wash bed linens and night clothes (don't shake them). Keep toilet seats clean.

**Distributed by:**

McKesson Corp.

via Strategic Sourcing Services LLC.

Memphis, TN 38141

**Questions or comments?**

Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST.

**Product of India**

948026899

L0001175

REV 10/25

**carton label 1oz**

**NDC 70677-1312-1**

**Foster & Thrive**

**Pinworm**

**Treatment**

**Pyrantel Pamoate**

**Suspension\***

## ANTHELMINTIC

- Single dose effectiveness
- Measuring cup included
- For Ages 2 Years & Up

1 fl. oz. (30 ml)



## PINWORM TREATMENT

pyrantel pamoate suspension

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1312
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>PYRANTEL PAMOATE</b> (UNII: 81BK194Z5M) (PYRANTEL - UNII:4QIH0N49E7)	PYRANTEL	144 mg in 1 mL
---	----------	----------------

**Inactive Ingredients**

Ingredient Name	Strength
<b>CITRIC ACID</b> (UNII: 2968PHW8QP)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MAGNESIUM ALUMINUM SILICATE</b> (UNII: 6M3P64V0NC)	
<b>METHYLCELLULOSE</b> (UNII: Z944H5SN0H)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	

**Product Characteristics**

<b>Color</b>	yellow (pale yellow to yellow)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BANANA	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1312-1	1 in 1 CARTON	12/05/2025	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M024	12/05/2025	

**Labeler** - STRATEGIC SOURCING SERVICES (116956644)

**Registrant** - BIONPHARMA INC. (079637826)

**Establishment**

Name	Address	ID/FEI	Business Operations
OrBion Pharmaceuticals Private Limited		854403569	manufacture(70677-1312)

Revised: 12/2025

STRATEGIC SOURCING SERVICES