ARC FIRST AID CONTAINS 101 PIECES HARDCASE - benzalkonium chloride, lidocaine, water, isopropyl alcohol, aspirin American Red Cross

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ARC FIRST AID Contains 101 PIECES Hardcase

ACTIVE INGREDIENTS:

Benzalkonium Chloride 0.13% Lidocaine HCL 0.5%

Purpose: First aid antiseptic, external analgesic

Uses: First aid to help prevent infection and for the temporary relief of pain and itching associated with:

Minor Cuts

Scrapes

Burns

Warnings:

For external use only

Do not use: In eyes, in large quantities, over raw blistered areas, or on deep puncture wounds, animal bites or serious burns, for more than one week

Keep out of reach of Children.

If ingested, contact a Poison Control Center right away.

Directions: Clean affected area, Apply small amount not more than 3 times daily.

May be covered with a sterile bandage.

Other Information:

Store at room temperature

Genuine First Aid Burn Cream

Antiseptic Pain Relief With Aloe

Net Wt 0.9g (1/32 oz)

Manufactured in CHINA for

Genuine First Aid.

Reorder BC-001

Genuine **First Aid Burn Cream**

Antiseptic Pain Relief With Aloe

ACTIVE INGREDIENTS: Benzalkonium Chloride 0.13% Lidocaine HCL 0.5%

Net Wt 0.9g · (1/32 oz)



Drug Facts

Purpose: First aid antiseptic, external analgesic

Uses: First aid to help prevent infection and for the temporary relief of pain and itching associated with: • minor cuts

scrapes • burns

Warnings:

For external use only

Do not use • in eyes • in large quantities · over raw blistered areas, or on deep puncture wounds, animal bites or serious burns • for more than one week Keep out of reach of children. If ingested, contact a Poison Control Center right away. Directions: • clean affected area apply small amount not more than 3 times daily • may be covered with a sterile bandage Other information: Store at room temperature



LOT 600609



2015/01

Active Ingredient: **Purpose**

Benzalkonium Chloride 0.40%...... First Aid Antiseptic

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive Ingredient: Purified water

LOT/EXP: Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

GENUINE FIRST AID

Made in CHINA Inactive ingredient: Purified water. towelette appropriately after single use. and use to cleanse desired skin area. Discard Directions: Tear open packet, unfold towelette EAR HERE poqà. Do not use: In the eyes, or over large areas of the occnr, consult a physician immediately. It unusual redness, swelling or other symptoms or contact a Poison Control Center right away. reach of children. It swallowed, get medical help Warnings: For external use only. Keep out of hands and body without soap and water. prevent infection. Antiseptic cleansing of face, Use: For Professional and Hospital use. Helps Benzalkonium Chloride 0.40%...First Aid Antiseptic Purpose: Active Ingredient:

DRUG FACTS - Antiseptic Towelette



Active Ingredient:

Purified Water USP...q.s.

Inactive Ingredients:

Sodium Chloride USP	44mg
Monobasic Sodium Phosphate USP Sodium Phosphate Dibasic USP	18 mg 111 mg
Edetate Disodium USP	10 mg
Benzalkonium Chloride	0.5mg

NF (as preservative)
Store in a cool place. For irrigation only.
Discard unused portion of the solution.
Not for injection. Warning:
If you experience eye pain, changes in vision, continued redness or irritation of the eye,
or if the condition worsens or persists for more than 72 hours, discontinue use and consult a doctor.
Do not use if solution changes color or becomes cloudy. Directions
Remove contacts before using.
Twist top to remove.
Flush the affected area as needed. Control
Rate of flow by pressure on the bottle. Do not touch
tip of the container to any surface. Do not reuse.
If necessary continue flushing with emergency eyewash or shower.
Discard bottle after use. Uses:
For flushing or irrigating the eyes to
remove loose foreign material, air pollutants,
or chlorinated wate Code No.: GUJ/DRUG/G/1080
Batch No.:
Mfg Date:
Exp: Date:
10ml
Sterile Isotonic Buffered Genuine
Eyewash
For single use only



Directions: Remove contact lens before using. Twist top to remove. Flush the affected area as needed. Control rate of flow by pressure on the bottle. Do not touch tip of the container to any surface. Do not reuse. If necessary, continue flushing with emergency eyewash or shower. Discard bottle after use.

Uses: For flushing or irrigating the eyes to remove loose foreign material, air pollutants, or chlorinated water.

Each 10 ml contains:

Active Ingredient: Purified Water ...q.s.

Inactive Ingredients:

Sodium Chloride Monobasic Sodium Phosphate Dibasic Sodium Phosphate Edetate Disodium Benzalkonium Chloride NF (as preservativa)

Store in a cool place. For irrigation only Discard unused portion of the solution.

Mfg.Lic.No.: KTK/28/332/200 Batch No.: Mfg. date:

Purpose



Warning:

If you experience eye pain, changes in vision, continued redness or irritation of the eye, or if the condition worsens or persists for more than 72 hours, discontinue use and consult a doctor. Do not use if solution changes color or becomes cloudy





Active Ingredient:

Isopropyl Alcohol, 70% v/v..... Antiseptic

Use: For preparation of skin before injection.

Warnings: For external use only.

Flammable - keep away from fire or flame

Store at room temperature 15-30 degree Celsius (59-86 degree Fahrenheit)

Do not use: with electrocautery, in the eyes.

Enter section text here

Keep out of reach of children. If swallowed,

get medical help or contact a Poison Control

Center right away.

Wipe Injection site vigorously and discard.

Inactive Ingredient: Purified water.

LOT/EXP: Made in CHINA

20140301

Alcohol Cleansing Pad Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID



DRUG FACTS - Alcohol Cleansing Pad

Active Ingredient: Isopropyl Alcohol, 70% v/v . . . Use: For preparation of the skin before injection. Warnings: For external use only. Flammable - keep away from fire or flame. Do not use: with electrocautery, in the eyes Stop use if irritation and redness develop. If condition persists for more than 72 hours, consult your doctor. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away Directions: Wipe injection site vigorously and Other information: Store at room temperature 15°-30° C (59°-86° F) Inactive ingredient: Purified water. LOT/EXP: Made in CHINA

Enter section text here

20140301

Enter section text here

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of reye's syndrome, a rare but serious illness.

Stomach bleeding warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if the user; has had stomach ulcers or bleeding problems takes a blood thinning (anticoagulant) or steroid drug takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) takes more or for a longer time than directed is age 60 or older has 3 or more alcoholic drinks every while using this product.

Do not use

if you are allergic to aspirin or any other pain relievers/fever reducers

for more than 10 days for pain unless directed by a doctor

for more than 3 days for fever unless directed by a doctor

for at least 7 days after a tonsillectomy or oral surgery

Ask a doctor before use if you have;

asthma

stomach problems (such as heartburn, upset stomach, or stomach pain) that persist or recur, ulcers or bleeding problems

a child experiencing arthritis pain

if you are;

allergic to aspirin

taking prescription drug for anticoagulation (thinning of blood), diabetes, gout or arthritis Stop use and ask a doctor if;

an allergic reaction occurs, Seek medical help right away

pain or fever persists

new or unexpected symptoms occur

redness or swelling is present

ringing in ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in unborn child or complications during delivery.

Enter section text here

Enter section text here

Enter section text here

Enter section text here

MADE IN USA

Distributed by GENUINE FIRST AID

600 Cleveland Str Suite 400, Clearwater, FL 33755

2 Tablets

GENUINE FIRST AID.

Chewable Aspirin 81mg

2 Tablets



Chewable Aspirin 81mg

Uses

■ temporary relief of minor aches, pains and headaches

to reduce fever associated with colds, sore throats, teething

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Stomach bleeding warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if the user ■ has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed ■ is age 60 or older ■ has 3 or more alcoholic drinks every day while using this product

Do not use ■ if you are allergic to aspirin or any other pain relievers/fever reducers ■ for more than 10 days for pain unless directed by a doctor ■ for more than 3 days for fever unless directed by a doctor ■ for at least 7 days after a tonsillectomy or oral surgery

American Red Cross

Carrying Case

1 Soft-side Bag

1 Easy Access Pocket System

Small Cuts and Burns

25 Adhesive Plastic Bandages 3/4"x3"

25 Junior Plastic Bandages 3/8"x1-1/2"

1 Adhesive Plastic Bandage 1"x3"

1 Adhesive Spot Bandage 7/8"x7/8"

1 Elbow/Knee Adhesive Bandage

1 Burn Cream

20 Cotton Tipped Applicators

Medium Cuts and Scratches

- 1 Antiseptic Towelettes
- 1 Sterile Gauge Pad 2"x2"
- 1 Roller Gauge Bandage 2"x4.1yds.
- 1 Sterile Eye Pad
- 1 Sterile Eye Wash 10ml
- 2 Butterfly Wound Closure

Severe Bleeding and Burns

- 1 First Aid Tape Roll 1/2"x2.5 yds.
- 1 Combine Pad 5"x9"
- 1 Roller Gauge Bandage 3"x4.1yds

Warnings (continued)

Ask a doctor before use if you have asthma stomach problems (such as heartburn, upset stomach, or stomach pain) that persist or recur, uclers or bleeding problems a child experiencing arthritis pain

if you are ■allergic to aspirin ■ taking a prescription drug for anticoagulation (thinning of the blood), diabetes, gout or arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away ■ pain or fever persists ■ new or unexpected symptoms occur ■ redness or swelling is present ■ ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions Do not use more than directed Adults and Children (12 years and older): Chew 4 to 8 tablets with water every 4 hours. Do not exceed 48 tablets in 24 hours unless directed by a doctor.

Children Under 12 Years: Consult a doctor

Other information

■ store at room temperature
 ■ tamper evident sealed packets
 ■ do not use any opened or torn packets

Inactive ingredients flavor, FD&C yellow #6, saccharin sodium, silicon dioxide, starch, stearic acid, sucrose

MADE IN USA Distributed by GENUINE FIRST AID

600 Cleveland Str Suite 400, Clearwater, FL 33755

CPR

1 CPR Breathing Barrier

Protection

- 2 Medical Grade Vinyl Gloves
- 1 Alcohol Cleansing Pad
- 2 Chewable Aspirin Tablets
- 1 Triangular Bandage 42"x42"x59"

Instruments

- 1 Emergency First Aid Guide
- 6 Assorted Safety Pins
- 1 Plastic Tweezers

Visit RedCross.org for information on training in First Aid, CPR, AED and more

Manufactured in China for:

Genuine First Aid LLC. 600 Cleveland Street Suite 400 Clearwater FL 33755

www.GenuineFirstAid.com

GENUINE FIRST AID

Copyright c 2009 Genuine First Aid LLC. All rights reserved.



ARC FIRST AID CONTAINS 101 PIECES HARDCASE

benzalkonium chloride, lidocaine, water, isopropyl alcohol, aspirin kit

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:52164-1103

Packaging

ш	0 0			
Ш	# Item Code	Package Description	Marketing Start Date	Marketing End Date
П	1 NDC:52164-1103-1	1 in 1 KIT		

Quantity of Parts

Quan	Quantity of Furth		
Part #	Package Quantity	Total Product Quantity	
Part 1	1 PACKAGE	0.9 g	
Part 2	1 PACKAGE	0.8 mL	
Part 3	1 BOTTLE	10 mL	
Part 4	1 PACKAGE	0.5 mL	
Part 5	1 PACKAGE	2	

Part 1 of 5

GENUINE FIRST AID BURN ANTISEPTIC PAIN RELIEF WITH ALOE

benzalkonium chloride, lidocaine cream

Product Information

Item Code (Source)	NDC:52124-0004
Route of Administration	TOPICAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BENZALKO NIUM CHLO RIDE (UNII: F5UM2KM3W7) (BENZALKO NIUM - UNII:7N6 JUD5X6 Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
LIDO CAINE (UNII: 98 PI200987) (LIDO CAINE - UNII:98 PI200987)	LIDOCAINE	0.5 g in 100 g

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0004-1	0.9 g in 1 PACKAGE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part345	04/28/2010	

Part 2 of 5

ANTISEPTIC TOWELETTE

benzalkonium chloride liquid

Product Information

Item Code (Source) NDC:52124-0001

Route of Administration TOPICAL

Active Ingredient/Active Moiety

y		
Ingredient Name	Basis of Strength	Strength
(BENZALKONIUM CHLORIDE	0.40 mL in 100 mL

Inactive Ingredients

Ingredient Name Strength

WATER (UNII: 059QF0KO0R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0001-1	0.8 mL in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	nart333	0.4/28/2010	

Part 3 of 5

STERILE ISOTONIC BUFFERED GENUINE EYEWASH

water liquid

Product Information

Item Code (Source)	NDC:52124-0005
Route of Administration	OPHTHALMIC

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	

Inactive Ingredients		
Ingredient Name	Strength	
SODIUM CHLORIDE (UNII: 451W47IQ8X)		
SODIUM PHO SPHATE, MO NO BASIC (UNII: 3980 JIH2SW)		
SO DIUM PHO SPHATE, DIBASIC (UNII: GR686LBA74)		
EDETATE DISO DIUM (UNII: 7FLD9 1C86K)		
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)		

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:52124-0005-1	10 mL in 1 BOTTLE		

Marketing Information				
Marketing Category Application Number or Monograph Citation Marketing Start Date Marketing En				
OTC monograph final	part349	04/28/2010		

Part 4 of 5

ALCOHOL CLEANSING PAD

isopropyl alcohol liquid

Product Information	
Item Code (Source)	NDC:52124-0002
Route of Administration	TOPICAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	70 mL in 100 mL	

Inactive Ingredients	
Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	04/28/2010	

Part 5 of 5

CHEWABLE ASPIRIN

aspirin tablet, chewable

Product Information

Item Code (Source) NDC:52124-0012

ORAL Route of Administration

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients			
Ingredient Name	Strength		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
SACCHARIN (UNII: FST467XS7D)			
SODIUM CATION (UNII: LYR4M0NH37)			
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)			
STARCH, CORN (UNII: O8232NY3SJ)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
SUCROSE (UNII: C151H8 M554)			

Product Characteristics				
Color	orange (ORANGE)	Score	no score	
Shape	ROUND	Size	11mm	
Flavor		Imprint Code	ASPIRIN	
Contains				

Packaging					
# Item Code	Package Description	Marketing Start Date	Marketing End Date		
1 NDC:52124-0012-1	2 in 1 PACKAGE				

Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
OTC monograph final	part345	04/28/2010				
Marketing Info	rmation					
Marketing Info	rmation Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
		Marketing Start Date 04/28/2010	Marketing End Date			

Labeler - American Red Cross (003255213)

Establishment						
Name	Address	ID/FEI	Business Operations			
GFA Production (Xiamen) Co., Ltd		421256261	manufacture			

Revised: 6/2010 American Red Cross