

**IBUPROFEN- ibuprofen tablet, film coated**  
**Better Living Brands, LLC**

-----  
**Signature Select 44-393**

***Active ingredient (in each orange caplet)***

Ibuprofen USP, 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - headache
  - the common cold
  - menstrual cramps
  - backache
  - toothache
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- facial swelling
- blisters
- hives
- shock
- skin reddening
- asthma (wheezing)
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - slurred speech
  - leg swelling
  - trouble breathing
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 caplet, 2 caplets may be used
  - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

***Other information***

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

***Inactive ingredients***

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

***Questions or comments?***

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

***Principal Display Panel***

**COMPARE TO**

Active Ingredient of Motrin® IB Caplets

NDC 21130-393-14

***Signature***

**SELECT®**

**IBUPROFEN**

**TABLETS USP, 200 mg**

PAIN RELIEVER /  
FEVER REDUCER  
**(NSAID)**

Actual Size

500 COATED  
CAPLETS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

†This product is not  
manufactured or  
distributed by  
Kenvue Inc.,  
distributors of  
Motrin® IB Caplets.

50844  
REV1221A39314

DISTRIBUTED BY: BETTER LIVING BRANDS LLC  
P.O. BOX 99, PLEASANTON, CA 94566-0009  
1-888-723-3929

COMPARE TO  
Active Ingredient of Motrin® IB Caplets†

NDC 21130-393-14

Signature  
SELECT™

IBUPROFEN

TABLETS USP, 200 mg

PAIN RELIEVER /  
FEVER REDUCER  
(NSAID)

Actual Size

500 COATED  
CAPLETS

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredient  
(in each orange caplet)

Ibuprofen USP,  
200 mg (NSAID)\* ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

Purpose

Uses

■ temporarily relieves minor aches and pains  
due to: ■ headache ■ the common cold  
■ menstrual cramps ■ backache ■ toothache  
■ muscular aches ■ minor pain of arthritis  
■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic  
reaction, especially in people allergic to aspirin.  
Symptoms may include:  
■ facial swelling ■ hives ■ shock  
■ skin reddening ■ asthma (wheezing) ■ rash  
If an allergic reaction occurs, stop use and seek medical  
help right away.

**Stomach bleeding warning:** This product contains an  
NSAID, which may cause severe stomach bleeding. The  
chance is higher if you  
■ take more or for a longer time than directed  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or  
nonprescription NSAIDs (aspirin, ibuprofen, naproxen,  
or others) ■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ have 3 or more alcoholic drinks every day while using  
this product

**Heart attack and stroke warning:** NSAIDs, except  
aspirin, increase the risk of heart attack, heart failure,

Drug Facts (continued)

and stroke. These can be fatal. The risk is higher if you  
use more than directed or for longer than directed.

Do not use

■ if you have ever had an allergic reaction  
to any other pain reliever/fever reducer  
■ right before or after heart surgery

Ask a doctor before use if

■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as  
heartburn ■ you are taking a diuretic  
■ you have high blood pressure, heart disease, liver  
cirrhosis, kidney disease, asthma, or had a stroke  
■ you have problems or serious side effects from taking  
pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition  
■ taking aspirin for heart attack or stroke, because  
ibuprofen may decrease this benefit of aspirin  
■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of  
the following signs of stomach bleeding: ■ feel faint  
■ have bloody or black stools ■ vomit blood  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain ■ slurred speech  
■ leg swelling ■ trouble breathing  
■ weakness in one part or side of body  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health  
professional before use. It is especially important not  
to use ibuprofen at 20 weeks or later in pregnancy  
unless definitely directed to do so by a doctor because  
it may cause problems in the unborn child or  
complications during delivery.

**Keep out of reach of children.** In case of overdose,  
get medical help or contact a Poison Control Center  
right away.

Directions

■ do not take more than directed  
■ the smallest effective dose should be used  
■ adults and children 12 years and over: take 1 caplet  
every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 caplet,  
2 caplets may be used  
■ do not exceed 6 caplets in 24 hours, unless  
directed by a doctor  
■ children under 12 years: ask a doctor

Other information

■ store between 20°-25°C (68°-77°F)  
■ avoid excessive heat 40°C (104°F)  
■ use by expiration date on package

Inactive ingredients

carnauba wax, colloidal  
silicon dioxide, corn starch, FD&C yellow #6 aluminum  
lake, hypromellose, lactose anhydrous, magnesium  
stearate, microcrystalline cellulose, polydextrose,  
polyethylene glycol, sodium starch glycolate, stearic  
acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391  
8:30 AM-4:00 PM ET, Monday-Friday

51714

RD 25125

3

21130 7875

50844

REV1221A39314

smartlabel®

Scan here  
for more  
information

DISTRIBUTED BY: BETTER LIVING BRANDS LLC  
P.O. BOX 99, PLEASANTON, CA 94566-0009  
1-888-723-3929

†This product is not  
manufactured or  
distributed by  
Kenvue Inc.,  
distributor of  
Motrin® IB Caplets.

Signature Select 44-393

IBUPROFEN

ibuprofen tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-393
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>FD&amp;C YELLOW NO. 6 ALUMINUM LAKE</b> (UNII: GYP6Z2JR6Q)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	44;393
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-393-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/1999	
2	NDC:21130-393-15	1 in 1 CARTON	03/01/1999	04/30/2022
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:21130-393-12	1 in 1 CARTON	03/01/1999	05/21/2017
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/01/1999	

**Labeler** - Better Living Brands, LLC (009137209)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(21130-393)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(21130-393)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(21130-393)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(21130-393)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(21130-393)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(21130-393)

Revised: 7/2025

Better Living Brands, LLC