

**MARCS MUCUS RELIEF- guaifenesin tablet**  
**Marc Glassman Inc**

-----  
**Drug Facts**

**Active ingredient** (per tablet)

Guaifenesin 400mg

**Purpose**

Expectorant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus
- helps make coughs more productive

**Warnings**

**Ask doctor before use if you have**

- persistent or chronic cough, such as occurs with smoking, asthma, bronchitis or emphysema
- cough is accompanied by excessive phlegm (mucus)

**Stop use and ask doctor if**

- Symptoms are accompanied by fever, rash or persistent headache
- cough persists for more than 1 week or tends to recur

**A persistent cough may be a sign of a serious condition.**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control

Center immediately.

**Directions**

- **Adults and children 12 years of age and over:** take 1 tablet every 4 hours as needed
- **Children 6 to 10 under 12 years of age:** take 1/2 tablet every 4 hours as needed
- **Children under 6 years of age:** consult a doctor

**Do not exceed 6 doses in a 24 hour period or as directed by a doctor**

**Other Information**

store at 15°-30°C (59°-86°F)

**Inactive ingredients**

magnesium stearate, microcrystalline cellulose. May also contain (colloidal) silicon dioxide, (co) povidone, dicalcium phosphate, maltodextrin, sodium starch glycolate, stearic acid.



Caution: do not use if imprinted seal under cap is broken or missing

**Drug Facts**

| Active ingredient (in each caplet) | Purpose     |
|------------------------------------|-------------|
| Guaifenesin 400mg                  | Expectorant |

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus
- helps make coughs more productive

**Warnings**

**Ask a doctor before use if you have**

- persistent or chronic cough, such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough is accompanied by excessive phlegm (mucus)

**Stop use and ask a doctor if**

- cough persists for more than 1 week or tends to recur
- A persistent cough may be a sign of a serious condition.
- symptoms are accompanied by fever, rash or persistent headache

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.**

**Directions**

|  |  |
|--|--|
| For adults and children 12 years and older | take 1 caplet every 4 hours as needed. Do not exceed 6 caplets in 24 hours   |
| For children 6 to under 12 years           | take 1/2 caplet every 4 hours as needed. Do not exceed 3 caplets in 24 hours |
| For children under 6 years                 | consult a doctor   |

**Other information** store at room temperature 15°-30°C (59°-86°F)

**Inactive ingredients** magnesium stearate, methylcellulose, microcrystalline cellulose, polydextrose K30, polydextrose 90F, silicon dioxide, stearic acid

Distributed by: Marc Glassman, Inc. REV 2/20 GUAFMAR  
West 130th Street, Cleveland, OH 44130

No Varnish in this area  
DO NOT PRINT BOX



# MARCS MUCUS RELIEF

guaifenesin tablet

## Product Information

|                     |                |                           |               |
|---------------------|----------------|---------------------------|---------------|
| <b>Product Type</b> | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:68998-845 |
|---------------------|----------------|---------------------------|---------------|

|                                |      |
|--------------------------------|------|
| <b>Route of Administration</b> | ORAL |
|--------------------------------|------|

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength |
|---|-------------------|----------|
| <b>Guaifenesin</b> (UNII: 495W7451VQ) (Guaifenesin - UNII:495W7451VQ) | Guaifenesin       | 400 mg   |

### Inactive Ingredients

| Ingredient Name                              | Strength |
|--|----------|
| <b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)       |          |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)       |          |
| <b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)    |          |
| <b>COPOVIDONE</b> (UNII: D9C330MD8B)         |          |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30) |          |

### Product Characteristics

|                 |       |                     |          |
|-----------------|-------|---------------------|----------|
| <b>Color</b>    | white | <b>Score</b>        | 2 pieces |
| <b>Shape</b>    | OVAL  | <b>Size</b>         | 17mm     |
| <b>Flavor</b>   |       | <b>Imprint Code</b> | PH063    |
| <b>Contains</b> |       |                     |          |

### Packaging

| # | Item Code        | Package Description  | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:68998-845-50 | 1 in 1 CARTON  | 02/26/2020           |                    |
| 1 |                  | 50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012                                     | 02/26/2020           |                    |

**Labeler** - Marc Glassman Inc (094487477)

**Registrant** - Reese Pharmaceutical Co (004172052)

### Establishment

| Name     | Address | ID/FEI    | Business Operations    |
|----------|---------|-----------|------------------------|
| Pharbest |         | 557054835 | manufacture(68998-845) |

