

I2BLUE- i2pure iodine antibacterial soak liquid

I2Pure Corp.

Active Ingredient



I2BLUE IODINE ANTIBACTERIAL SOAK

IODINE TOPICAL SOLUTION

Drug Facts

Active ingredient: Iodine Topical Solution (Titratable Iodine 0.15%)

Purpose: Antiseptic

Use:

First aid antiseptic to help prevent the risk of bacterial contamination and infection in minor cuts, scrapes and burns.

WARNINGS:

For external use only. Do not use on infants or if pregnant or lactating. Do not use in the eyes or apply over large areas of the body. In case of deep or puncture wounds, animal bites or severe burns, consult a doctor.

Stop use and ask a doctor if; irritation and redness develop; condition persists for more than 72 hours.

Do not use longer than one week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Clean the affected area.
- Apply product to the affected area or immerse the affected area in product and soak for 1-3 minutes.
- May be covered with a sterile bandage.
- May be used several times per day.

Other information:

Store at controlled room temperature 15°-30°C (59°-86°F). Keep out of direct sunlight.

Inactive Ingredients:

Citric Acid, Glycerin, Sodium Iodide, Starch, Water

**Sterile When Unopened
For Professional and Hospital Use**

16 fl oz / 473 mL

NDC: 83465-401-16
Manufactured by I2Pure Corp.
850 Warren Wilson Rd., Swannanoa, NC 28778
Patent Pending. MADE IN THE USA.



ML01-005A



I2BLUE IODINE ANTIBACTERIAL SOAK

IODINE TOPICAL SOLUTION

Drug Facts

Active ingredient: Iodine Topical Solution (Titratable Iodine 0.15%)

Purpose: Antiseptic

Use:

First aid antiseptic to help prevent the risk of bacterial contamination and infection in minor cuts, scrapes and burns.

WARNINGS:

For external use only. Do not use on infants or if pregnant or lactating. Do not use in the eyes or apply over large areas of the body. In case of deep or puncture wounds, animal bites or severe burns, consult a doctor.

Stop use and ask a doctor if; irritation and redness develop; condition persists for more than 72 hours.

Do not use longer than one week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Clean the affected area.
- Apply product to the affected area or immerse the affected area in product and soak for 1-3 minutes.
- May be covered with a sterile bandage.
- May be used several times per day.

Other information:

Store at controlled room temperature 15°-30°C (59°-86°F). Keep out of direct sunlight.

Inactive Ingredients:

Citric Acid, Glycerin, Sodium Iodide, Starch, Water

**Sterile When Unopened
For Professional and Hospital Use**

16 fl oz / 473 mL

NDC: 83465-401-16
Manufactured by I2Pure Corp.
850 Warren Wilson Rd., Swannanoa, NC 28778
Patent Pending. MADE IN THE USA.



ML01-005A



I2BLUE IODINE ANTIBACTERIAL SOAK

IODINE TOPICAL SOLUTION

Drug Facts

Active ingredient: Iodine Topical Solution (Titratable Iodine 0.15%)

Purpose: Antiseptic

Use:

First aid antiseptic to help prevent the risk of bacterial contamination and infection in minor cuts, scrapes and burns.

WARNINGS:

For external use only. Do not use on infants or if pregnant or lactating. Do not use in the eyes or apply over large areas of the body. In case of deep or puncture wounds, animal bites or severe burns, consult a doctor.

Stop use and ask a doctor if; irritation and redness develop; condition persists for more than 72 hours.

Do not use longer than one week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Clean the affected area.
- Apply product to the affected area or immerse the affected area in product and soak for 1-3 minutes.
- May be covered with a sterile bandage.
- May be used several times per day.

Other information:

Store at controlled room temperature 15°-30°C (59°-86°F). Keep out of direct sunlight.

Inactive Ingredients:

Citric Acid, Glycerin, Sodium Iodide, Starch, Water

**Sterile When Unopened
For Professional and Hospital Use**

16 fl oz / 473 mL

NDC: 83465-401-16
Manufactured by I2Pure Corp.
850 Warren Wilson Rd., Swannanoa, NC 28778
Patent Pending. MADE IN THE USA.



ML01-005A



I2BLUE IODINE ANTIBACTERIAL SOAK

IODINE TOPICAL SOLUTION

Drug Facts

Active ingredient: Iodine Topical Solution (Titratable Iodine 0.15%)

Purpose: Antiseptic

Use:

First aid antiseptic to help prevent the risk of bacterial contamination and infection in minor cuts, scrapes and burns.

WARNINGS:

For external use only. Do not use on infants or if pregnant or lactating. Do not use in the eyes or apply over large areas of the body. In case of deep or puncture wounds, animal bites or severe burns, consult a doctor.

Stop use and ask a doctor if; irritation and redness develop; condition persists for more than 72 hours.

Do not use longer than one week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Clean the affected area.
- Apply product to the affected area or immerse the affected area in product and soak for 1-3 minutes.
- May be covered with a sterile bandage.
- May be used several times per day.

Other information:

Store at controlled room temperature 15°-30°C (59°-86°F). Keep out of direct sunlight.

Inactive Ingredients:

Citric Acid, Glycerin, Sodium Iodide, Starch, Water

**Sterile When Unopened
For Professional and Hospital Use**

16 fl oz / 473 mL

NDC: 83465-401-16
Manufactured by I2Pure Corp.
850 Warren Wilson Rd., Swannanoa, NC 28778
Patent Pending. MADE IN THE USA.



ML01-005A



I2BLUE IODINE ANTIBACTERIAL SOAK

IODINE TOPICAL SOLUTION

Drug Facts

Active ingredient: Iodine Topical Solution (Titratable Iodine 0.15%)

Purpose: Antiseptic

Use:

First aid antiseptic to help prevent the risk of bacterial contamination and infection in minor cuts, scrapes and burns.

WARNINGS:

For external use only. Do not use on infants or if pregnant or lactating. Do not use in the eyes or apply over large areas of the body. In case of deep or puncture wounds, animal bites or severe burns, consult a doctor.

Stop use and ask a doctor if; irritation and redness develop; condition persists for more than 72 hours.

Do not use longer than one week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Clean the affected area.
- Apply product to the affected area or immerse the affected area in product and soak for 1-3 minutes.
- May be covered with a sterile bandage.
- May be used several times per day.

Other information:

Store at controlled room temperature 15°-30°C (59°-86°F). Keep out of direct sunlight.

Inactive Ingredients:

Citric Acid, Glycerin, Sodium Iodide, Starch, Water

**Sterile When Unopened
For Professional and Hospital Use**

16 fl oz / 473 mL

NDC: 83465-401-16
Manufactured by I2Pure Corp.
850 Warren Wilson Rd., Swannanoa, NC 28778
Patent Pending. MADE IN THE USA.



ML01-005A



I2BLUE IODINE ANTIBACTERIAL SOAK

IODINE TOPICAL SOLUTION

Drug Facts

Active ingredient: Iodine Topical Solution (Titratable Iodine 0.15%)

Purpose: Antiseptic

Use:

First aid antiseptic to help prevent the risk of bacterial contamination and infection in minor cuts, scrapes and burns.

WARNINGS:

For external use only. Do not use on infants or if pregnant or lactating. Do not use in the eyes or apply over large areas of the body. In case of deep or puncture wounds, animal bites or severe burns, consult a doctor.

Stop use and ask a doctor if; irritation and redness develop; condition persists for more than 72 hours.

Do not use longer than one week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Clean the affected area.
- Apply product to the affected area or immerse the affected area in product and soak for 1-3 minutes.
- May be covered with a sterile bandage.
- May be used several times per day.

Other information:

Store at controlled room temperature 15°-30°C (59°-86°F). Keep out of direct sunlight.

Inactive Ingredients:

Citric Acid, Glycerin, Sodium Iodide, Starch, Water

**Sterile When Unopened
For Professional and Hospital Use**

16 fl oz / 473 mL

NDC: 83465-401-16
Manufactured by I2Pure Corp.
850 Warren Wilson Rd., Swannanoa, NC 28778
Patent Pending. MADE IN THE USA.



ML01-005A



I2BLUE IODINE ANTIBACTERIAL SOAK

IODINE TOPICAL SOLUTION

Drug Facts

Active ingredient: Iodine Topical Solution (Titratable Iodine 0.15%)

Purpose: Antiseptic

Use:

First aid antiseptic to help prevent the risk of bacterial contamination and infection in minor cuts, scrapes and burns.

WARNINGS:

For external use only. Do not use on infants or if pregnant or lactating. Do not use in the eyes or apply over large areas of the body. In case of deep or puncture wounds, animal bites or severe burns, consult a doctor.

Stop use and ask a doctor if; irritation and redness develop; condition persists for more than 72 hours.

Do not use longer than one week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Clean the affected area.
- Apply product to the affected area or immerse the affected area in product and soak for 1-3 minutes.
- May be covered with a sterile bandage.
- May be used several times per day.

Other information:

Store at controlled room temperature 15°-30°C (59°-86°F). Keep out of direct sunlight.

Inactive Ingredients:

Citric Acid, Glycerin, Sodium Iodide, Starch, Water

**Sterile When Unopened
For Professional and Hospital Use**

16 fl oz / 473 mL

NDC: 83465-401-16
Manufactured by I2Pure Corp.
850 Warren Wilson Rd., Swannanoa, NC 28778
Patent Pending. MADE IN THE USA.



ML01-005A



I2BLUE IODINE ANTIBACTERIAL SOAK

IODINE TOPICAL SOLUTION

Drug Facts

Active ingredient: Iodine Topical Solution (Titratable Iodine 0.15%)

Purpose: Antiseptic

Use:

First aid antiseptic to help prevent the risk of bacterial contamination and infection in minor cuts, scrapes and burns.

WARNINGS:

For external use only. Do not use on infants or if pregnant or lactating. Do not use in the eyes or apply over large areas of the body. In case of deep or puncture wounds, animal bites or severe burns, consult a doctor.

Stop use and ask a doctor if; irritation and redness develop; condition persists for more than 72 hours.

Do not use longer than one week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Clean the affected area.
- Apply product to the affected area or immerse the affected area in product and soak for 1-3 minutes.
- May be covered with a sterile bandage.
- May be used several times per day.

Other information:

Store at controlled room temperature 15°-30°C (59°-86°F). Keep out of direct sunlight.

Inactive Ingredients:

Citric Acid, Glycerin, Sodium Iodide, Starch, Water

**Sterile When Unopened
For Professional and Hospital Use**

16 fl oz / 473 mL

NDC: 83465-401-16
Manufactured by I2Pure Corp.
850 Warren Wilson Rd., Swannanoa, NC 28778
Patent Pending. MADE IN THE USA.



ML01-005A



I2BLUE IODINE ANTIBACTERIAL SOAK

IODINE TOPICAL SOLUTION

Drug Facts

Active ingredient: Iodine Topical Solution (Titratable Iodine 0.15%)

Purpose: Antiseptic

Use:

First aid antiseptic to help prevent the risk of bacterial contamination and infection in minor cuts, scrapes and burns.

WARNINGS:

For external use only. Do not use on infants or if pregnant or lactating. Do not use in the eyes or apply over large areas of the body. In case of deep or puncture wounds, animal bites or severe burns, consult a doctor.

Stop use and ask a doctor if; irritation and redness develop; condition persists for more than 72 hours.

Do not use longer than one week unless directed by a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Clean the affected area.
- Apply product to the affected area or immerse the affected area in product and soak for 1-3 minutes.
- May be covered with a sterile bandage.
- May be used several times per day.

Other information:

Store at controlled room temperature 15°-30°C (59°-86°F). Keep out of direct sunlight.

Inactive Ingredients:

Citric Acid, Glycerin, Sodium Iodide, Starch, Water

**Sterile When Unopened
For Professional and Hospital Use**

16 fl oz / 473 mL

NDC: 83465-401-16
Manufactured by I2Pure Corp.
850 Warren Wilson Rd., Swannanoa, NC 28778
Patent Pending. MADE IN THE USA.



ML01-005A

I2BLUE

i2pure iodine antibacterial soak liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83465-401
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	750 mg in 500 mL

Inactive Ingredients	
Ingredient Name	Strength
STARCH, POTATO (UNII: 8I089SAH3T)	4000 mg in 500 mL
WATER (UNII: 059QF0KO0R)	445 mL in 500 mL
CITRIC ACID (UNII: 2968PHW8QP)	2 mL in 500 mL
SODIUM IODIDE (UNII: F5WR8N145C)	4500 mg in 500 mL
GLYCERIN (UNII: PDC6A3C0OX)	48.75 mL in 500 mL

Product Characteristics			
Color	blue	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83465-401-16	475 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/12/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	11/12/2025	

Labeler - I2Pure Corp. (117480499)

Establishment			
Name	Address	ID/FEI	Business Operations
I2Pure Corp.		117480499	manufacture(83465-401)