

**VICKS DAYQUIL NYQUIL HOT REMEDY COLD AND FLU- acetaminophen,
dextromethorphan hbr, diphenhydramine hcl
Procter & Gamble Manufactura S de RL de CV**

**Vicks ®DayQuil™ HOT REMEDY Vicks ®NyQuil™ HOT REMEDY
COLD & FLU**

Vicks® DayQuil™ HOT REMEDY COLD & FLU

Drug Facts

DayQuil™

Active Ingredient (in each Packet)

Acetaminophen 500 mg

Dextromethorphan HBr 20 mg

Purpose

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening

- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

Other information

- **each packet contains:** potassium 10 mg
- phenylketonurics: contains phenylalanine 68 mg per dose
- store at no greater than 25°C (77°F)

Inactive ingredients:

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate

Questions?

1-800-362-1683

MADE IN MEXICO

DIST. BY PROCTER & GAMBLE,

CINCINNATI OH 45202

Vicks® NyQuil™ HOT REMEDY COLD & FLU

Drug Facts

NyQuil™

Active Ingredient (in each Packet)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Purpose

Pain reliever/fever reducer

Antihistamine/Cough suppressant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor sore throat & bronchial irritation
- cough to help you sleep
- runny nose
- sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- with any other product containing diphenhydramine, even one used on skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (Certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

Other information

- **each packet contains:**potassium 10 mg
- phenylketonurics: contains phenylalanine 68 mg per dose
- store at no greater than 25°C (77°F)

Inactive ingredients

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF SACHET IS OPENED OR TORN

Keep outer package for complete product information.

MADE IN MEXICO

**DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202**

PRINCIPAL DISPLAY PANEL - 6 packet package

NEW FORMULAS

VALUE PACK

Vicks[®] DayQuil[™] HOT REMEDY

COLD & FLU

Acetaminophen, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough

Non-Drowsy

Vicks[®] NyQuil[™] HOT REMEDY

COLD & FLU

Acetaminophen, Diphenhydramine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough, Sneezing, Runny Nose

Nighttime Relief

HONEY LEMON FLAVORED

3 DAYQUIL PACKETS; 3 NYQUIL PACKETS; 6 TOTAL PACKETS



Drug Facts	NyQuil™	Drug Facts	DayQuil™
Active Ingredients (in each Packet)	Purpose	Active Ingredients (in each Packet)	Purpose
Acetaminophen 500 mg	Pain reliever/Fever reducer	Acetaminophen 500 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 25 mg	Cough suppressant	Dextromethorphan HBr 20 mg	Cough suppressant
Uses Temporarily relieves common cold/flu symptoms: • minor aches & pains • sore throat • fever • headache • cough due to minor throat & bronchial irritation • cough to help you sleep • runny nose • sneezing		Uses Temporarily relieves common cold/flu symptoms: • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 6 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 6 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • with any other product containing diphenhydramine, even one used on skin • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist.		Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist.	

Drug Facts (continued)	NyQuil™	Drug Facts (continued)	DayQuil™
Ask a doctor before taking this product. Ask a doctor before use if you have • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin When using this product • excitability may occur, especially in children • marked drowsiness may occur • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.		Ask a doctor before taking this product. Ask a doctor before use if you have • liver disease • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, or emphysema Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions • take only as directed • do not exceed 6 doses per 24 hrs • Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes. • If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat. Age Dose adults & children 12 yrs & over one packet (dose) every 4 hours children under 12 yrs do not use Other information • each packet contains: potassium 10 mg • phenylephrine: contains phenylephrine 58 mg per dose • store at no greater than 25°C (77°F) Inactive ingredients acacia, potassium, aspartame, carboxymethylcellulose, citric acid, D&G Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate Questions? 1-800-362-1683		Directions • take only as directed • do not exceed 6 doses per 24 hrs • Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes. • If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat. Age Dose adults & children 12 yrs & over one packet (dose) every 4 hours children under 12 yrs do not use Other information • each packet contains: potassium 10 mg • phenylephrine: contains phenylephrine 58 mg per dose • store at no greater than 25°C (77°F) Inactive ingredients acacia, potassium, aspartame, carboxymethylcellulose, citric acid, D&G Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate Questions? 1-800-362-1683	

VICKS DAYQUIL NYQUIL HOT REMEDY COLD AND FLU

acetaminophen, dextromethorphan hbr, diphenhydramine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58933-556
--------------	----------------	--------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-556-16	1 in 1 PACKAGE; Type 0: Not a Combination Product	06/11/2025	
2	NDC:58933-556-06	1 in 1 PACKAGE; Type 0: Not a Combination Product	06/11/2025	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 PACKET	1
Part 2	1 PACKET	5 g

Part 1 of 2

VICKS NYQUIL HOT REMEDY COLD AND FLU

acetaminophen, diphenhydramine hcl powder, for solution

Product Information

Item Code (Source)	NDC:58933-555
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
TRICALCIUM PHOSPHATE (UNII: K4C08XP666)	
SUCROSE (UNII: C151H8M554)	
ASPARTAME (UNII: Z0H242BBR1)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
CARBOXYMETHYLCELLULOSE (UNII: 05JZI7B19X)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor	HONEY (Honey Lemon)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-555-01	1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/11/2025	

Part 2 of 2

VICKS DAYQUIL HOT REMEDY COLD AND FLU

acetaminophen, dextromethorphan hbr powder, for solution

Product Information

Item Code (Source)	NDC:58933-552
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 5 g
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg in 5 g

Inactive Ingredients

Ingredient Name	Strength
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SUCROSE (UNII: C151H8M554)	
TRICALCIUM PHOSPHATE (UNII: K4C08XP666)	
CARBOXYMETHYLCELLULOSE (UNII: 05JZI7B19X)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ASPARTAME (UNII: Z0H242BBR1)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Product Characteristics

Color	white	Score	
Shape		Size	

Flavor		HONEY (Honey Lemon)	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-552-01	5 g in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	06/11/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	06/11/2025	

Labeler - Procter & Gamble Manufactura S de RL de CV (812807550)