

**AROAMAS COT NICOTINE TRANSDERMAL PATCHES STEP 2- nicotine
transdermal patches patch
PCHEERY LLC**

**Initial Drug Listing - AROAMAS COT NICOTINE TRANSDERMAL PATCHES STEP
2**

Nicotine 12%

Menthol 0.3%

Stop smoking aid

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

If you have

- Heart disease, recent heart attack, or irregular heartbeat, Nicotine can increase your heart rate.
- High blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- An allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- Stomach ulcer or diabetes.
- History of seizures.

If you are

- Using a non-nicotine stop smoking drug
- Taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

If you have vivid dreams or other sleep disturbances, remove this patch at bedtime.

- Skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash.
- Irregular heartbeat or palpitations occur
- You get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat.
- You have symptoms of an allergic reaction (such as difficulty breathing or rash)

Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

- if you are under 18 years of age, ask a doctor before use.
- before using this product, read the enclosed self help guide for complete directions

and other information

- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day,
- use the following schedule below:

STEP1 Use one 21 mg patch/day	STEP2 Use one 14 mg patch/day	STEP3 Use one 7 mg patch/day
Weeks 1-6	Weeks 7-8	Weeks 9-10

- If you smoke 10 or less cigarettes per day, do not use Step 1(21 mg). Start with Step 2(14 mg) for 6 weeks, then Step3(7 mg) for 2 weeks and then stop.
- apply one new patch every 24 hours on skin that is dry, clean and hairless.
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch Save pouch to use for patch disposal Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day.
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.
- you may wear the patch for 18 to 24 hours
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing and MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

CODONOPSIS PILOSULA ROOT, CHRYSANTHEMUM INDICUM FLOWER, CLOVE LEAF OIL, TOBACCO LEAF, CAMPHOR OIL

AROAMAS



AROAMAS
Nicotine
Transdermal Patches

STOP SMOKING AID

AROAMAS

- ✓ 100% pure natural plant
- ✓ Help you stop smoking
- ✓ Stop smoking aid

21
PATCHES

Store at 20-25°C
(68-77°F)

STEP
2

Delivered Over 24 Hours

14
mg

If you smoke more than 10 cigarettes per day: Start with Step 1
If you smoke 10 or less cigarettes per day: Start with Step 2

Comments or Questions?
Call toll free: 833-976-2377
Or visit us at: aroamas.com



Drug Facts

Active ingredient (in each patch)	Purpose
Nicotine 12%	Stop smoking aid
menthol 0.3%	Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product • If you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- If you are under 18 years of age, ask a doctor before use.
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

STEP1	STEP2	STEP3
Use one 21 mg patch/day	Use one 14 mg patch/day	Use one 7 mg patch/day
Weeks 1-6	Weeks 7-8	Weeks 9-10

- If you smoke 10 or less cigarettes per day, do not use Step 1 (21 mg). Start with Step 2 (14 mg) for 6 weeks, then Step 3 (7 mg) for 2 weeks and then stop.
- apply one new patch every 24 hours on skin that is dry, clean and hairless.
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day.
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.
- you may wear the patch for 18 to 24 hours
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing and MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

MADE IN CHINA



NDC 82739-025-01
Manufactured for P-Cherry LLC
Republic, MO 65728
Email us: service@aroamas.com

AROAMAS COT NICOTINE TRANSDERMAL PATCHES STEP 2

nicotine transdermal patches patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85831-034
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	12 g in 100
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	0.3 g in 100

Inactive Ingredients

Ingredient Name	Strength
CLOVE LEAF OIL (UNII: VCA5491KVF)	
CODONOPSIS PILOSULA ROOT (UNII: 02VJ3CEY9G)	
CAMPBOR OIL (UNII: 75IZZ8Y727)	
TOBACCO LEAF (UNII: 6YR2608RSU)	
CHRYSANTHEMUM INDICUM FLOWER (UNII: I6OER6U04L)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85831-034-01	21 in 1 CARTON	12/30/2025	
1		1 in 1 PATCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	12/30/2025	

Labeler - PCHEERY LLC (061286986)

Revised: 12/2025

PCHEERY LLC