

WOFANLI QUIT SMOKING NICOTINE PATCHES- nicotine patches patch
Shenzhen Paxi Trading Co., Ltd.

85835-016 V2 add Pack 180Patches

Active Ingredient

Nicotine 3mg

Purpose

Stop smoking aid

Use

Apply to dry, clean, hair-free skin.

Remove backing from patch and immediately press onto skin. Hold for 10 seconds.

Wash hands after applying or removing patch. Save pouch to use for patch disposal.

Dispose of the used patches by folding sticky ends together and putting in pouch.

If vivid dreams occur, remove the patch before bedtime.

Do not cut patch in half or into smaller pieces

It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Warnings

If you are pregnant or breast feeding, only use this medicine on the advice of your health care provider.

Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Do not use

Ask a doctor or pharmacist before use:

- Contraindicated in heart disease, recent MI, or arrhythmia. Nicotine may increase heart rate.
- History of seizures.
- Uncontrolled hypertension. Nicotine may elevate blood pressure.
- Stomach ulcer or diabetes .
- If allergic to adhesive or with pre-existing skin conditions, higher risk of dermal reactions may occur.

Ask Doctor

Ask a doctor or pharmacist before use if you:

- are using non-nicotine smoking cessation drug .
- are taking prescription medications for depression or asthma, your dosage may require adjustment.

When Using

When using this product

- if you experience vivid dreams or other sleep disturbances, remove the patch at bedtime.

Stop Use

- Patch-induced erythema persists beyond 4 days, localized edema develops, or a cutaneous rash appears.
- irregular heartbeat or palpitations occur.
- Symptoms of nicotine overdose may include persistent nausea, vomiting, dizziness, weakness, and rapid heartbeat.
- You have symptoms of an allergic reaction (such as difficulty breathing or rash).

Keep Out Of Reach Of Children

KEEP OUT OF REACH OF CHILDREN AND PETS.

Used patches contain enough nicotine to poison children and pets. If swallowed, seek medical help immediately or contact a Poison Control Center 1-800-222-1222 right away. Save the pouch for patch disposal. Dispose of used patches by folding the sticky ends together and placing them in the pouch.

Directions

If you are under 18 years of age, ask a doctor before use.

before using this product, read the enclosed self-help guide for complete directions and other information

begin using the patch on your quit day

If you smoke more than 10 cigarettes per day, follow the schedule below:

Weeks 1 and 4

STEP 1

1 patch every 1 to 2 hours

Weeks 5 and 7

STEP 2

1 patch every 2 to 4 hours

Weeks 8 and 10

STEP 3

use 1/2

patch/day

Weeks 9 and 10

STEP 3

Inactive ingredients: acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer.

Manufactured for: Shenzhen Faxi Trading Co., Ltd.

Address: 102, 48 Zhenye Road, Hehua Community, Pinghu Street, Longgang District, Shenzhen, Guangdong 518000, China

After-sales contact information: Faxitrading@outlook.com

MADE IN CHINA

ING

LOW-NICOTINE CONTENT
The content of small patches is **3mg** PATCHES

HELP TO QUIT SMOKING
Nicotine Transdermal System Patch

• Not for sale to persons under 18 years of age.
• Proof of age required.

Trade Mark: WOFANLI MF Date: 11/18/2025
Lot number: NP720251118 EXP Date: 11/17/2028

FaNIli

Drug Facts

Active ingredient (in each patch) Nicotine, 3mg delivered over 2 hours. **Purpose** Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings If you are pregnant or breast feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor or pharmacist before use:

- Contraindicated in heart disease, recent MI, or arrhythmia. Nicotine may increase heart rate.
- History of seizures.
- Uncontrolled hypertension. Nicotine may elevate blood pressure.
- Stomach ulcer or diabetes.
- If allergic to adhesive or with pre-existing skin conditions, higher risk of dermal reactions may occur.

Ask a doctor or pharmacist before use if you:

- are using non-nicotine smoking cessation drug.
- are taking prescription medications for depression or asthma, your dosage may require adjustment.

When using this product • If you experience vivid dreams or other sleep disturbances, remove the patch at bedtime.

Stop Use:

- Patch-induced erythema persists beyond 4 days, localized edema develops, or a cutaneous rash appears.
- Irregular heartbeat or palpitations occur.
- Symptoms of nicotine overdose may include persistent nausea, vomiting, dizziness, weakness, and rapid heartbeat.
- You have symptoms of an allergic reaction (such as difficulty breathing or rash).

KEEP OUT OF REACH OF CHILDREN AND PETS. Used patches contain enough nicotine to poison children and pets. If swallowed, seek medical help immediately or contact a Poison Control Center 1-800-222-1222 right away. Save the pouch for patch disposal. Dispose of used patches by folding the sticky ends together and placing them in the pouch.

Directions • If you are under 18 years of age, ask a doctor before use. • Before using this product, read the enclosed self-help guide for complete directions and other information. • Begin using the patch on your quit day. If you smoke more than 10 cigarettes per day, follow the schedule below:

Weeks 1 and 4	Weeks 5 and 7	Weeks 8 and 10
STEP 1 1 patch every 1 to 2 hours	STEP 2 1 patch every 2 to 4 hours	STEP 3 1 patch every 4 to 8 hours

- To increase your chances of quitting, use at least 9 nicotine patches daily during the first week. If you experience strong or frequent cravings, you may apply a second patch within two hours. However, do not use one patch after another consecutively.
- If you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 3 weeks.
- Apply to dry, clean, hair-free skin.
- Remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- Wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- If vivid dreams occur, remove the patch before bedtime.
- Do not cut patch in half or into smaller pieces.
- It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information: Store at room temperature

Inactive ingredients: acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer.

Manufactured for: Shenzhen Faxi Trading Co., Ltd.

Address: 102, 48 Zhenye Road, Hehua Community, Pinghu Street, Longgang District, Shenzhen, Guangdong 518000, China

After-sales contact information: Faxitrading@outlook.com

MADE IN CHINA

For family's protection patches are supplied in child-resistant pouches.
DO NOT USE IF INDIVIDUAL PATCHES OPEN OR TORN.

QUIT SMOKING
Nicotine Patches

FaNIli

Nicotine Patches

QUIT SMOKING
STOP SMOKING AID

CIRCULAR DESIGN
LOW-NICOTINE CONTENT

The content of small patches is **3mg**

120
PATCHES

QUIT SMOKING AID
HELP TO QUIT SMOKING
Nicotine Transdermal System Patch

• Not for sale to persons under 18 years of age.
• Proof of age required.
• Not for sale for vending machines or from any source where a price of age cannot be verified.

Trade Mark: WOFANLI MF Date: 11/18/2025
Lot number: NP820251118 EXP Date: 11/17/2028

**Drug Facts**

Active ingredient (in each patch) **Purpose**
 Nicotine, 3 mg delivered over 2 hours Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings If you are pregnant or breast feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor or pharmacist before use:

- Contraindicated in heart disease, recent MI, or arrhythmia, Nicotine may increase heart rate.
- History of seizures.
- Uncontrolled hypertension, Nicotine may elevate blood pressure.
- Stomach ulcer or diabetes.
- If allergic to adhesive or with pre-existing skin conditions, higher risk of dermal reactions may occur.

Ask a doctor or pharmacist before use if you:

- are using non-nicotine smoking cessation drug.
- are taking prescription medications for depression or asthma, your dosage may require adjustment.

When using this product • If you experience vivid dreams or other sleep disturbances, remove the patch at bedtime.

Stop Use:

- Patch-induced erythema persists beyond 4 days, localized edema develops, or a cutaneous rash appears.
- Irregular heartbeat or palpitations occur.
- Symptoms of nicotine overdose may include persistent nausea, vomiting, dizziness, weakness, and rapid heartbeat.
- You have symptoms of an allergic reaction (such as difficulty breathing or rash).

KEEP OUT OF REACH OF CHILDREN AND PETS. Used patches contain enough nicotine to poison children and pets. If swallowed, seek medical help immediately or contact a Poison Control Center 1-800-222-1222 right away. Save the pouch for patch disposal. Dispose of used patches by folding the sticky ends together and placing them in the pouch.

Directions • If you are under 18 years of age, ask a doctor before use. • Before using this product, read the enclosed self-help guide for complete directions and other information. • Begin using the patch on your quit day. If you smoke more than 10 cigarettes per day, follow the schedule below.

Weeks 1 to 4	Weeks 5 to 7	Weeks 8 to 10
STEP 1	STEP 2	STEP 3
1 patch every 1 to 2 hours	1 patch every 2 to 4 hours	1 patch every 4 to 8 hours

- To increase your chances of quitting, use at least 8 nicotine patches daily during the first week. If you experience strong or frequent cravings, you may apply a second patch within two hours. However, do not use one patch after another consecutively.
- If you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 3 weeks.
- Apply to dry, clean, hair-free skin.
- Remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- Wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- If vivid dreams occur, remove the patch before bedtime.
- Do not cut patch in half or into smaller pieces.
- It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information: Store at room temperature

Inactive ingredients: acrylate adhesive, aluminum polyester, cellulose paper, methacrylic acid copolymer.

Manufactured for: Shenzhen Paxi Trading Co., Ltd.

Address: 102, 48 Zhenye Road, Heliu Community, Pinghu Street, Longgang District, Shenzhen, Guangdong 518000, China

After-sales contact information: Paxitrading@outlook.com

MADE IN CHINA

For family's protection patches are supplied in child-resistant pouches. DO NOT USE IF INDIVIDUAL PATCHES OPEN OR TORN.

QUIT SMOKING Nicotine Patches



Nicotine Patches

QUIT SMOKING
STOP SMOKING AID

CIRCULAR DESIGN
LOW-NICOTINE CONTENT

The content of small patches is **3mg**

180
PATCHES

QUIT SMOKING AID
HELP TO QUIT SMOKING

Nicotine Transdermal
System Patch

• Not for sale to persons under 18 years of age.
• Proof of age required.
• Not for sale to vending machines or from any source where proof of age cannot be verified.

Trade Mark: WOFANLI MF Date: 11/18/2025
Lot number: NP820251118 EXP Date: 11/17/2028

WOFANLI QUIT SMOKING NICOTINE PATCHES

nicotine patches patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85835-016
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	3 mg in 100

Inactive Ingredients

Ingredient Name	Strength
ACRYLIC ACID (UNII: J94PBK7X8S)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
POLYESTER-7 (UNII: 0841698D2F)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85835-016-01	60 in 1 BOX; Type 0: Not a Combination Product	12/09/2025	
2	NDC:85835-016-02	120 in 1 BOX; Type 0: Not a Combination Product	12/09/2025	
3	NDC:85835-016-03	180 in 1 BOX; Type 0: Not a Combination Product	12/09/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	12/09/2025	

Labeler - Shenzhen Paxi Trading Co., Ltd. (633886766)

Establishment

Name	Address	ID/FEI	Business Operations
Shenzhen Paxi Trading Co., Ltd.		633886766	manufacture(85835-016)

Revised: 1/2026

Shenzhen Paxi Trading Co., Ltd.