

**READYPREP CHG- chlorhexidine gluconate 2% cloth**  
**Medline Industries**

-----  
**ReadyPrep CHG**

Chlorhexidine gluconate 2% solution

Antiseptic

**Uses**

- helps reduce bacteria that can potentially cause skin infection
- for preparation of skin prior to surgery

For external use only

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar punctures or in contact with the meninges
- on open skin wounds
- as a general skin cleanser

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

- keep out of eyes, ears and mouth. May cause serious or permanent injury if chlorhexidine is permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- do not microwave
- product and packaging are not sterile. Follow your hospital policy for skin preparation with non-sterile products.
- to open package: identify the tear notch labeled on the front of the package. Grasp with both hands on both sides of the tear notch and tear to expose cloth.
- transfer contents onto prep table, avoiding contact between cloth and outside of package to reduce risk of cloth contamination
- use first cloth to prepare the skin area indicated for a moist or dry site, making certain to keep the second cloth where it will not be contaminated. Use second cloth to prepare larger areas.
- dry surgical sites: (such as abdomen or arm): use one cloth to cleanse each 161 cm<sup>2</sup> area

(approximately) 5 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to dry for one (1) minute. Do not rinse.

- moist surgical sites (such as inguinal fold): use one cloth to cleanse each 65 cm<sup>2</sup> area (approximately 2 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to dry for one (1) minute. Do not rinse.
- discard each cloth after a single use
- after package has been opened, discard any unused cloths

#### **Other Information**

- store product flat
- store between 20-25 degrees Celsius (68-77 degrees Fahrenheit)
- avoid excessive heat above 40 degrees Celsius (104 degrees Fahrenheit)
  
- benzalkonium chloride
- dimethicone emulsion
- glycerin
- isopropyl alcohol
- propylene glycol
- purified water USP

DO NOT FLUSH

Non-Sterile.

Product is not made with Natural Rubber Latex.

Cloth: 100% Polyester

REF MSC095CHG

[www.medline.com](http://www.medline.com)

©2017 Medline Industries, Inc.

Medline is a registered trademark of Medline Industries, Inc.

Manufactured by

Medline Industries, Inc.,

Northfield, IL 60093 USA.

Made in USA with domestic and foreign materials.

Questions, comments, or ordering call: 1-800-MEDLINE

Customer Service Hours:

(Monday to Friday, 7 AM to 7 PM CST)

RI17MSC

#### **Pouch label**

READYPREPTM CHG

CHLORHEXIDINE GLUCONATE 2%

PATIENT PREOPERATIVE SKIN PREPARATION CLOTH

**Drug Facts (continued)**

- discard each cloth after a single use.
- after package has been opened, discard any unused cloths.

**Other information**

- store product flat.
- store between 20-25°C (68-77°F).
- avoid excessive heat above 40°C (104°F).

**Inactive ingredients**

benzalkonium chloride, dimethylsiloxane emulsion, glycerin, isopropyl alcohol, propylene glycol, purified water USP



**DO NOT FLUSH**  
**Non-Sterile.**  
**Product is not made with Natural Rubber Latex.**  
**Cloth: 100% Polyester**

**REF** MSC095CHG



[www.medline.com](http://www.medline.com)  
 ©2017 Medline Industries, Inc. Medline is a registered trademark of Medline Industries, Inc. Manufactured by Medline Industries, Inc., Northfield, IL 60093 USA. Made in USA with domestic and foreign materials.  
**Questions, comments, or ordering call: 1-800-MEDLINE**  
 Customer Service Hours (Monday to Friday, 7 AM to 7 PM CST) R18M5C



**MEDLINE** NDC 5329-244-01

**READYPREP<sup>®</sup> CHG**

**2% CHLORHEXIDINE GLUCONATE\* CLOTH**

**PATIENT PREOPERATIVE SKIN PREPARATION**

**NON-STERILE**

\*EQUIVALENT TO 500 MG CHLORHEXIDINE GLUCONATE PER CLOTH

- Provides rapid bactericidal action against a broad spectrum of microorganisms
- Significantly reduces the number of microorganisms on intact skin
- Demonstrates continued antimicrobial activity for up to 6 hours after application

**FRAGRANCE FREE**

**RINSE FREE**

SINGLE USE ONLY.  
 FOR EXTERNAL USE ONLY.  
 DO NOT MICROWAVE.

**2** 9x10.5 in (22.9x26.7 cm)  
**DISPOSABLE CLOTHS** **REF** MSC095CHG

**Drug Facts**

Active ingredient	Purpose
Chlorhexidine gluconate 2% solution	Antiseptic

**Uses**

- helps reduce bacteria that can potentially cause skin infection.
- for preparation of skin prior to surgery.

**Warnings**

**For external use only**

**Allergy alert:**  
 This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product.
- for lacerated punctures or in contact with the meninges.
- on open skin wounds.
- as a general skin cleanser.

**When using this product**

- keep out of eyes, ears and mouth. May cause serious or permanent injury if chlorhexidine is permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.**  
 If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- do not microwave.
- product and packaging are not sterile. Follow your hospital policy for skin preparation with non-sterile products.
- **to open package:** identify the tear notch labeled on the front of the package. Grasp with hands on both sides of the tear notch and tear to expose cloth.
- transfer contents onto prep table, avoiding contact between cloth and outside of package to reduce risk of cloth contamination.
- use first cloth to prepare the skin area indicated for a moist or dry site, making certain to keep the second cloth where it will not be contaminated. Use second cloth to prepare larger areas.
- **dry surgical sites** (such as abdomen or arm): use one cloth to cleanse each 101 cm<sup>2</sup> area (approximately 8 X 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to dry for one (1) minute. Do not rinse.
- **moist surgical sites** (such as inguinal RAO): use one cloth to cleanse each 60cm<sup>2</sup> area (approximately 2x3 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to dry for one (1) minute. Do not rinse.

**NON-STERILE**

**EQUIVALENT TO 500MG CHLORHEXIDINE GLUCONATE PER CLOTH**

Provides rapid bactericidal action against a broad spectrum of microorganisms

Significantly reduces the number of microorganisms on intact skin

Demonstrates continued antimicrobial activity for up to 6 hours after application

**FRAGRANCE FREE**

**RINSE FREE**

SINGLE USE ONLY.

FOR EXTERNAL USE ONLY.

DO NOT MICROWAVE

2 9 X 10.5 IN (22.9 X 26.7 CM) DISPOSABLE CLOTHS

REF MSC095CHG

## READYPREP CHG

chlorhexidine gluconate 2% cloth

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53329-244
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	500 mg

### Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
DIMETHICONE 350 (UNII: 2Y53S6ATLU)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53329-244-02	24 in 1 CARTON	01/14/2019	
1	NDC:53329-244-01	2 in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA207964	01/14/2019	

**Labeler** - Medline Industries (025460908)