

LICE TREATMENT DEDICATED- ivermectin lotion
Shenzhen Zhenshangzekai Technology Co., Ltd.

lice treatment

ivermectin

Ask a doctor before use if you

Have any skin conditions or sensitivities.

- Are pregnant or plan to become pregnant.

It is not known if product can harm your unborn baby.

Are currently or planning to breast feed!

Avoid getting product on your breast to help prevent contact by your baby.

-Breathing difficulty occurs

-Eye irritation occurs

-Skin or scalp irritation continues or infection occurs

-Rash develops

Keep out of reach of children.If swallowed, get medical help or contact Poison Control Center(1-800-222-1222)right away.

If pregnant or breastfeeding, ask a health professional before use.

lice treatment

925-899-0135

TAMPER EVIDENT: DO NOT USE IF THEPUMP IS DAMAGED OR SEAL IS BROKEN. Store at room temperature 68°F-77°

Do not freeze.

keep carton and see Consumer Information Leaflet and warnings before use

lotion, topical

Butylated hydroxyanisole, castor oil, cetylalcohol, citric acid anhydrous, crodalan AWS,cyclomethicone, glycerin, imidurea, lanolinalcohols, methylparaben, oleyl alcohol, oliveoil, propylene glycol, propylparaben, purifiedwater, shea butter, sodium citrate anhydroussorbitan tristearate and stearyl alcohol.

Directions

Important: Read warnings before use

.Children 6 months of age to under 12 years of age :

an adult should supervise use

Adults and children 6 months of age and over:

Inspect

All houschold members should be checked by anothe!person for lice and/or nits (eggs)

Use a magnifying glassin bright light to help you see

thelice and/or nits(eggs)

Use a tool, such as a comb or two unsharpened pencilsto lift and part the hair

Look for tiny nits near the scalp, beginning at the back of the neck and behind the ears
Examine small sections of the hair (1-2 inches wide) at a time
Unlike dandruff, nits stick to the hair, Dandruff should move when lightly touched
If either lice or nits (eggs) are found, treat with product

Treat

Your hair and scalp must be DRY before applying product

Apply the product directly onto dry hair and scalp

Completely cover your scalp and hair closest to the scalp first, and then apply outwards towards the ends of your hair

Use only as much as needed to fully cover hair and scalp up to one full bottle, and discard the remaining
Rub product throughout your hair

It is important to completely cover your entire head so that all lice and eggs are exposed to the lotion, Be sure that each hair is coated from the scalp to the tip

Wait and Rinse

Allow product to stay on your hair and scalp for 10 minutes after it has been applied, Use a timer or clock
Start timing after you have completely covered your hair and scalp with product.

After 10 minutes, rinse product completely from your hair and scalp using only water.

After rinsing, dry and style hair as usual. Wait 24 hours before applying shampoo.

After treatment

Wash your hands after applying product.

-Nit combing is not necessary when treating with product for it to work, but if desired, a fine-tooth comb or special nit comb may be used to remove dead lice and nits. This is a single use product, Discard

bottle after use. Do not use again on the same person and same lice infestation without talking to a healthcare provider first. If infestation continues, see a doctor for other treatments.

Machine wash any bedding and clothing used by anyone having lice, Machine wash at high temperatures (150°F) and tumble in a hot dryer for 20 minutes. After finishing treatment with lice medicine, check everyone in your family for lice after one week.

Consider treatment for those who have lice.

Drug Facts (continued)

Directions

Important: Read warnings before use.

- Children 6 months of age to under 12 years of age: an adult should supervise use.
- Adults and children 6 months of age and over:

Inspect

- All household members should be checked by another person for lice and/or nits (eggs).
- Use a magnifying glass in bright light to help you see the lice and/or nits (eggs).
- Use a tool, such as a comb or two unsharpened pencils to lift and part the hair.
- Look for tiny nits near the scalp, beginning at the back of the neck and behind the ears.
- Examine small sections of the hair (1/2 inches wide) at a time.
- Unlike dandruff, nits stick to the hair. Dandruff should move when lightly touched.
- If either lice or nits (eggs) are found, treat with product.

Treat

- Your hair and scalp must be DRY before applying product.
- Apply the product directly onto dry hair and scalp.
- Completely cover your scalp and hair closest to the scalp first, and then apply outwards towards the ends of your hair.
- Use only as much as needed to fully cover hair and scalp - up to one full bottle, and discard the remaining.
- Rub product throughout your hair.
- It is important to completely cover your entire head so that all lice and eggs are exposed to the lotion. Be sure that each hair is coated from the scalp to the tip.

Wait and Rinse

- Allow product to stay on your hair and scalp for 10 minutes after it has been applied. Use a timer or clock. Start timing after you have completely covered your hair and scalp with product.
- After 10 minutes, rinse product completely from your hair and scalp using only water.
- After rinsing, dry and style hair as usual. Wait 24 hours before applying shampoo.



Drug Facts (continued)

After treatment.
Wash your hands after applying product.

- Nit combing is not necessary when treating with product for it to work, but if desired, a fine-tooth comb or special nit comb may be used to remove dead lice and nits.
- This is a single use product. Discard bottle after use.
- Do not use again on the same person and same lice infestation without talking to a healthcare provider first.
- If infestation continues, see a doctor for other treatments.
- Machine wash any bedding and clothing used by anyone having lice. Machine wash at high temperatures (150°F) and tumble in a hot dryer for 20 minutes.
- After finishing treatment with lice medicine, check everyone in your family for lice after one week. Consider treatment for those who have lice.


DANGER - FLAMMABLE: DO NOT USE IF THE PUMP IS DAMAGED OR SEAL IS BROKEN.

- Store at room temperature 68°F-77°F (20°C-25°C).
- Do not freeze.
- Keep carton and see Consumer Information Leaflet and warnings before use.

Inactive ingredients: Butylated hydroxyanisole, castor oil, cetyl alcohol, citric acid anhydrous, crodolan AWS, cyclomethicone, glycerin, inidurea, lanolin alcohols, methylparaben, oleyl alcohol, olive oil, propylene glycol, propylparaben, purified water, shea butter, sodium citrate anhydrous, sorbitan tristearate and stearyl alcohol.

Questions? 925-899-0135

LOT#1465652
MFG 03/09/2005
EXP DATE 03/09/2008
DISTRIBUTED BY: 112 Parkway Street
Needles, CA 92363, USA



Ivormetrn

0.5% Lotion Ivermectin

Lice Treatment
Dedicated



10 Minutes kills
lice eggs at the root!

Easy to rinse; Itch Control; Preventive effect

NET WT. 4OZ (125ml)

Drug Facts

Active ingredient: Ivermectin 0.5%
Purpose: Lice Treatment

Use: treats head lice

Warnings

For external use only. Use only on the scalp.

Do not use:

- On children under 6 months of age.
- Near the eyes.
- Inside the mouth, nose, ears, or vagina.
- On lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you:

- Have any skin conditions or sensitivities.
- Are pregnant or plan to become pregnant. It is not known if product can harm your unborn baby.
- Are currently or planning to breastfeed. Avoid getting product on your breast to help prevent contact by your baby.

When using this product:

- Keep eyes tightly closed and protect eyes with a washcloth or towel.
- If product gets into eyes, gently flush with water.
- Eye redness, soreness, or irritation can occur.
- Dandruff, dry skin, or burning sensation of the skin can occur.

Stop use and ask a doctor if:

- Breathing difficulty occurs.
- Eye irritation occurs.
- Skin or scalp irritation continues or infection occurs.
- Rash develops.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children.
If swallowed, get medical help or contact Poison Control Center (1-800-222-1222) right away.

LICE TREATMENT DEDICATED

ivermectin lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:87276-000
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IVERMECTIN (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)	IVERMECTIN	0.5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
LANOLIN ALCOHOLS (UNII: 884C3FA9HE)	
SHEA BUTTER (UNII: K49155WL9Y)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
OLIVE OIL (UNII: 6UYK2W1W1E)	
ANHYDROUS TRISODIUM CITRATE (UNII: RS7A450LGA)	
CASTOR OIL (UNII: D5340Y2I9G)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GLYCERIN (UNII: PDC6A3C0OX)	
IMIDUREA (UNII: M629807ATL)	
OLEYL ALCOHOL (UNII: 172F2WN8DV)	
SORBITAN TRISTEARATE (UNII: 6LUM696811)	
CYCLOMETHICONE (UNII: NMQ347994Z)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87276-000-01	1 g in 1 BOTTLE; Type 0: Not a Combination Product	12/10/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M031	12/10/2025	

Labeler - Shenzhen Zhenshangzekai Technology Co., Ltd. (518601007)

Revised: 12/2025

Shenzhen Zhenshangzekai Technology Co., Ltd.