LICE TREATMENT DEDICATED- ivermet lotion Shenzhen Zhenshangzekai Technology Co., Ltd.

Ivermectin 0.5%

Have any skin conditions or sensitivities.

Are pregnant or plan to become pregnant.

It is not known if product can harm your unborn baby.

Are currently or planning to breastfeed.

Avoid getting product on your breast to help prevent contact by your baby.

Breathing difficulty occurs.

Eye irritation occurs.

Skin or scalp irritation continues or infection occurs.

Rash develops.

Keep out of reach of children.

If swallowed, get medical help or contact Poison Control Center (1-800-222-1222) right away.

If pregnant or breastfeeding, ask a health professional before use.

Lice Treatment

925-899-0135

lotion, topical

TAMPER EVIDENT: DO NOT USE IF THEPUMP IS DAMAGED OR SEAL IS BROKEN. Store at room temperature 68°CF-77°

Do not freeze.

keep carton and see Consumer Information Leaflet and warnings before use

Butylated hydroxyanisole, castor oil, cetyl alcohol, citric acid anhydrous, crodalan AWS,cyclomethicone, glycerin, imidurea, lanolinalcohols, methylparaben, oleyl alcohol, olive oil, propylene glycol, propylparaben, purified water, shea butter, sodium ditrate anhydrous, sorbitan tristearate and stearyl alcohol.

Directions

Important: Read warnings before use

.Children 6 months of age to under 12 years of age :

an adult should supervise use

Adults and children 6 months of age and over:

Inspect

All household members should be checked by anothe!person for lice and/or nits (eggs) Use a magnifying glassin bright light to help you see thelice and/or nits(eggs)

Use a tool, such as a comb or two unsharpened pencilsto lift and part the hair Look for tiny nits near the scalp, beginning at the backof theneck and behind the ears Examine small sections of the hair (1-2 inches wide) at a time

.Unlike dandruff, nits stick to the hair, Dandruff shouldmove when lightly touched

If either lice or nits (eggs) are found, treat with product

Treat

Your hair and scalp must be DRY before applying product

Apply the product directly onto dry hair and scalp

Completely cover your scalp and hair closest to the scalp first.and then apply outwards towards the endsofyour hair

Use only as much asneeded to fully cover hair and scalp up to one full bottle, and discard the remaining Rub product throughout your hair

Itis important to completely cover your entire head sothat all lice and eggs are exposed to the lotion, Be surethat each hair is coated from the scalp to the tip

Wait and Rinse

Allow product to stay on your hair and scalp for 10minutes after it has been applied, Use a timer or clockStart timing after you have completely covered yourhair and scalp with product.

After 10 minutes, rinse product completely from youhair and scalp using only water. After rinsing, dry and style hair as usual. Wait 24 hoursbefore applying shampoo.

After treatment

Wash your hands after applying product.

-Nit combing is not necessary whentreating with product for it to work, but ifdesired, a fne-tooth comb or special nitcomb may be used to remove dead liceand nits. This is a single use product, Discard

bottle after use. Do not use again on the same personand same lice infestation without talkingto a healthcare provider frst. If infestation continues, see a doctor forother treatments.

Machine wash any bedding and clothingused by anyone having lice, Machinewash at high temperatures (150°F) and tumble in a hot dryer for 20 minutes. After fnishing treatment with licemedicine, check everyone in your family for lice after one week. Considertreatment for those who have lice.

Kills Lice: Just once is all it takes

IVERMET
0.5% Lice Treatment Lotion

ONCE 10 MINUTES! NO MESS

NET WT: 4 OZ (125 mL)



LICE TREATMENT DEDICATED

ivermet lotion

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:87276-001
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IVERMECTIN (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)	IVERMECTIN	0.5 g in 100 g	

Inactive Ingredients		
Ingredient Name	Strength	
CETYL ALCOHOL (UNII: 936JST6JCN)		
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)		
CASTOR OIL (UNII: D5340Y2I9G)		
CYCLOMETHICONE (UNII: NMQ347994Z)		
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)		
SORBITAN TRISTEARATE (UNII: 6LUM696811)		
SODIUM CITRATE (UNII: 1Q73Q2JULR)		
OLEYL ALCOHOL (UNII: 172F2WN8DV)		
PROPYLPARABEN (UNII: Z8IX2SC10H)		
WATER (UNII: 059QF0KO0R)		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
LANOLIN ALCOHOLS (UNII: 884C3FA9HE)		
OLIVE OIL (UNII: 6UYK2W1W1E)		
GLYCERIN (UNII: PDC6A3C0OX)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
METHYLPARABEN (UNII: A2I8C7HI9T)		
IMIDUREA (UNII: M629807ATL)		
SHEA BUTTER (UNII: K49155WL9Y)		

Product Characteristics			
Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87276-001- 01	1 g in 1 BOTTLE; Type 0: Not a Combination Product	12/10/2025	

Marketing In	Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M031	12/10/2025		

Shenzhen Zhenshangzekai Technology Co., Ltd.

Revised: 12/2025