

IVOMETN IVERMECTIN LICE TREATMENT- ivermectin lotion

Shenzhen Zhenshangzekai Technology Co., Ltd.

Ivermectin

Have any skin conditions or sensitivities.

Are pregnant or plan to become pregnant.

It is not known if product can harm your unborn baby.

Are currently or planning to breastfeed.

Avoid getting product on your breast to help prevent contact by your baby.

Breathing difficulty occurs

Eye irritation occurs

Skin or scalp irritation continues or infection occurs

Rash develops

Keep out of reach of children.

If swallowed, get medical help or contact Poison Control Center (1-800-222-1222) right away.

If pregnant or breastfeeding, ask a health professional before use.

Lice treatment

lotion, topical

Butylated hydroxyanisole, castor oil, cetylalcohol, citric acid anhydrous, crodalan AWS, cyclomethicone, glycerin, imidurea, lanolinalcohols, methylparaben, oleyl alcohol, oliveoil, propylene glycol, propylparaben, purifiedwater, shea butter, sodium citrate anhydroussorbitan tristearate and stearyl alcohol

Directions

Important: Read warnings before use

.Children 6 months of age to under 12 years of age :

an adult should supervise use

Adults and children 6 months of age and over:

Inspect

All household members should be checked by another person for lice and/or nits (eggs)

Use a magnifying glass in bright light to help you see

the lice and/or nits (eggs)

Use a tool, such as a comb or two unsharpened pencils to lift and part the hair

Look for tiny nits near the scalp, beginning at the back of the neck and behind the ears

Examine small sections of the hair (1-2 inches wide) at a time

.Unlike dandruff, nits stick to the hair, Dandruff should move when lightly touched

If either lice or nits (eggs) are found, treat with product

Treat

Your hair and scalp must be DRY before applying product

Apply the product directly onto dry hair and scalp

Completely cover your scalp and hair closest to the scalp first. and then apply outwards towards the ends of your hair

Use only as much as needed to fully cover hair and scalp up to one full bottle, and discard the remaining. Rub product throughout your hair.

It is important to completely cover your entire head so that all lice and eggs are exposed to the lotion. Be sure that each hair is coated from the scalp to the tip.

Wait and Rinse

Allow product to stay on your hair and scalp for 10 minutes after it has been applied. Use a timer or clock. Start timing after you have completely covered your hair and scalp with product.

After 10 minutes, rinse product completely from your hair and scalp using only water.

After rinsing, dry and style hair as usual. Wait 24 hours before applying shampoo.

After treatment

Wash your hands after applying product.

-Nit combing is not necessary when treating with product for it to work, but if desired, a fine-tooth comb or special nit comb may be used to remove dead lice and nits. This is a single use product. Discard

bottle after use. Do not use again on the same person and same lice infestation without talking to a healthcare provider first. If infestation continues, see a doctor for other treatments.

Machine wash any bedding and clothing used by anyone having lice. Machine wash at high temperatures (150°F) and tumble in a hot dryer for 20 minutes. After finishing treatment with lice medicine, check everyone in your family for lice after one week.

Consider treatment for those who have lice.

TAMPER EVIDENT: DO NOT USE IF THE PUMP IS DAMAGED OR SEAL IS BROKEN. Store at room temperature 68°F-77°F

Do not freeze.

Keep carton and see Consumer Information Leaflet and warnings before use.

IVORMETN

Ivermectin

Lotion, 0.5%

Lice Treatment

No second application required

Not nit combing required


Ivermectin lotion kills lice

One Bottle

One Time

10 Minutes

NET WT 4.4 OZ (125 g)



IVOMETN IVERMECTIN LICE TREATMENT

ivermectin lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:87276-003
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
IVERMECTIN (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)			IVERMECTIN	0.5 g in 100 g
Inactive Ingredients				
Ingredient Name				Strength
GLYCERIN (UNII: PDC6A3C0OX)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
SHEA BUTTER (UNII: K49155WL9Y)				
SORBITAN TRISTEARATE (UNII: 6LUM696811)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0KO0R)				
ANHYDROUS TRISODIUM CITRATE (UNII: RS7A450LGA)				
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)				
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)				
OLEYL ALCOHOL (UNII: 172F2WN8DV)				
CETYL ALCOHOL (UNII: 936JST6JCN)				
IMIDUREA (UNII: M629807ATL)				
CYCLOMETHICONE (UNII: NMQ347994Z)				
LANOLIN ALCOHOLS (UNII: 884C3FA9HE)				
OLIVE OIL (UNII: 6UYK2W1W1E)				
CASTOR OIL (UNII: D5340Y2I9G)				
Product Characteristics				
Color	white	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87276-003-01	1 g in 1 BOTTLE; Type 0: Not a Combination Product	12/10/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M031	12/10/2025	

Labeler - Shenzhen Zhenshangzekai Technology Co., Ltd. (518601007)

