

BACTOSHIELD CHG CHLORHEXIDINE GLUCONATE- chlorhexidine gluconate solution
SC Johnson Professional USA, Inc.

Bactoshield CHG
Chlorhexidine Gluconate Solution

Drug Facts

Active ingredient

Chlorhexidine gluconate 2% solution

Purpose

Antiseptic

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease
- **patient preoperative skin preparation:** preparation of the patient's skin prior to surgery
- **skin wound and general skin cleansing**

Warnings

For external use only

Allergy alert

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area
- as a preoperative skin preparation of the head or face

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub

- wet hands and forearms under running water for 30 seconds. Clean fingernails using a nailstick or similar cleaner.
- scrub for 1.5 minutes with about 8 mL of product with or without a wet brush paying close attention to the nails, cuticles, and skin between the fingers
- rinse thoroughly under running water for 30 seconds
- wash for an additional 1.5 minutes with 8 mL of product and rinse under running water for 30 seconds
- dry thoroughly

Healthcare personnel handwash

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Patient preoperative skin preparation

- apply product liberally to surgical site and swab for at least 2 minutes and dry with a sterile towel
- repeat procedure for an additional 2 minutes and dry with a sterile towel

Skin wound and general skin cleaning

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

Questions or comments?

call 1-800-587-3721

Monday through Friday 8:00 AM to 4:30 PM CST

PRINCIPAL DISPLAY PANEL - 118 mL Bottle Label

Bactoshield®

CHG 2%

Surgical Scrub

Chlorhexidine Gluconate 2% Solution Antiseptic

Non-Sterile Solution

SCJ PROFESSIONAL

HEALTHCARE

NDC 11084-038-03

Surgical Scrub

2% CHG

Antimicrobial

REORDER #

1322-39

SAP # 4000011062

1322-52J(622)

2SCJ04BTLLBL

deb

SKIN CARE

Net Contents:

4 fl oz (118 mL)

For Single Use

Bactoshield®
CHG 2%

Surgical Scrub

Chlorhexidine Gluconate 2% Solution Antiseptic
Non-Sterile Solution



Surgical Scrub



2% CHG



Antimicrobial

REORDER #
1322-39

SAP # 4000011062
1322-52J(622)
2SCJ04BTLLBL



Net Contents:
4 fl oz (118 mL)

For Single Use



4

Laundering/Cleaning Instructions:
Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

Manufactured for: SCJ Professional USA, Inc.
Charlotte, NC 28217
1-866-783-0422 www.scp.com

WARNING: Cancer -- www.P65Warnings.ca.gov.
Keep from freezing. Made in U.S.A.

Drug Facts

Active ingredient Chlorhexidine gluconate 2% solution Antiseptic

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease

Drug Facts (continued)

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Patient preoperative skin preparation:

- apply product liberally to surgical site and swab for at least 2 minutes and dry with a sterile towel
- repeat procedure for an additional 2 minutes and dry with a sterile towel

Skin wound and general skin cleaning:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients citric acid, coccamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

Questions or comments?

call 1-800-587-3721
Monday through Friday 8:00 AM to 4:30 PM CST

Drug Facts

- patient prepare surgery
- skin wk

Warning
For exte

Allergy a
This prod
reaction.

- wheezi
 - shock
 - facial s
 - hives
 - rash
- If an allen
seek mec

Do not u

- if you c
- chlorhe
- ingredi
- in cont
- in the g
- as a pr
- head o

▲ Lift Here

Warnings (continued)

Preoperative skin preparation: Application of the patient's skin prior to surgery and general skin cleansing

Topical use only

Alert: Product may cause a severe allergic reaction/difficulty breathing

Warning: Irritation occurs, stop use and seek medical help right away.

See: For the patient is allergic to chlorhexidine gluconate or any other ingredient in this product. Do not use with meningitis or other genital area. Do not use with operative skin preparation of the face.

When using this product

- Keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums.
- If solution should contact these areas, rinse out promptly and thoroughly with water.
- Wounds which involve more than the superficial layers of the skin should not be routinely treated.
- Repeated general skin cleansing of large body areas should not be done except when advised by a health care provider.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- Wet hands and forearms under running water for 30 seconds. Clean fingernails using a nailstick or similar cleaner.
- Scrub for 1.5 minutes with about 8 mL of product with or without a wet brush paying close attention to the nails, cuticles, and skin between the fingers.
- Rinse thoroughly under running water for 30 seconds.
- Wash for an additional 1.5 minutes with 8 mL of product and rinse under running water for 30 seconds.
- Dry thoroughly.

Drug Facts

Active ingredient Chlorhexidine gluconate 2% solution

Purpose Antiseptic

Uses

- surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease



Bactoshield® CHG 2%
Surgical Scrub
Chlorhexidine Gluconate 2% Solution Antiseptic Non-Sterile Solution

Icons: Surgical Scrub, 2% CHG, Antimicrobial

Net Contents: 4 fl oz (118 mL)

For Single Use

REORDER # 1322-39

SAP # 4000011062 1322-52J(622) 2SCJ048TLLBL

6 91240 06200 4

BACTOSHIELD CHG CHLORHEXIDINE GLUCONATE

chlorhexidine gluconate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11084-038
---------------------	----------------	---------------------------	---------------

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	20 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
COCO DIETHANOLAMIDE (UNII: 92005F972D)	
HYDROXYETHYL CELLULOSE (100 MPA.S AT 2%) (UNII: R33S7TK2EP)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
LAURAMINE OXIDE (UNII: 4F6FC4MI8W)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	YELLOW (Clear)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11084-038-03	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2022	12/31/2027

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019422	12/01/2022	12/31/2027

Labeler - SC Johnson Professional USA, Inc. (607378015)

Establishment

Name	Address	ID/FEI	Business Operations
Xttrium Laboratories, Inc.		007470579	MANUFACTURE(11084-038)