

**ACETAMINOPHEN 250 MG AND IBUPROFEN 125 MG - acetaminophen 250 mg and ibuprofen 125 mg tablets tablet, film coated**  
**MARKSANS PHARMA LIMITED**

-----  
**Acetaminophen 250 mg and Ibuprofen 125 mg Tablets**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg

Ibuprofen 125 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever

Pain reliever

**Uses**

- temporarily relieves minor aches and pains due to:
  - o headache
  - o toothache
  - o backache
  - o menstrual cramps
  - o muscular aches
  - o minor pain of arthritis

**Warnings**

**Acetaminophen liver damage warning:**

This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product

**Acetaminophen allergy alert:** may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away.

**NSAID allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**NSAID stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to acetaminophen or any other pain reliever
- right before or after heart surgery

**Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

## When using this product

- take with food or milk if stomach upset occurs

## Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - o feel faint
  - o vomit blood
  - o have bloody or black stools
  - o have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - o chest pain
  - o trouble breathing
  - o weakness in one part or side of body
  - o slurred speech
  - o leg swelling
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- any new symptoms appear

## If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- **do not take more than directed**

adults and children 12 years and over	• <b>take 2 caplets every 8 hours</b> while symptoms persist
children under 12 years	• ask a doctor

- do not take more than 6 caplets in 24 hours, unless directed by a doctor

## **Other information**

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

## **Inactive ingredients**

colloidal anhydrous silica, corn starch, croscarmellose sodium, glyceryl dibehenate, hypromellose, microcrystalline cellulose, polydextrose, povidone, red iron oxide, titanium dioxide, triacetin, yellow iron oxide

## **Questions or comments?**

Call **1-877-376-4271** (weekdays 9 AM to 5 PM)

Distributed by:

### **Time-Cap Labs, Inc.**

7 Michael Avenue,  
Farmingdale,  
NY 11735, USA

## **Made in India**

## **PRINCIPAL DISPLAY PANEL**

NDC 25000-080-72

Acetaminophen 250 mg and Ibuprofen 125 mg Tablets

18 count carton label

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

Drug Facts

Active ingredients (in each caplet)
Acetaminophen 250 mg
Ibuprofen 125 mg (NSAID\*)

Purposes
Pain reliever
Pain reliever

Uses
temporarily relieves minor aches and pains due to:
headache, toothache, backache, menstrual cramps, muscular aches, minor pain of arthritis

Warnings
Acetaminophen liver damage warning:
This product contains acetaminophen. Severe liver damage may occur if you take:

Drug Facts (continued)

with other drugs containing acetaminophen
more than 6 caplets in 24 hours, which is the maximum daily amount for this product
3 or more alcoholic drinks every day while using this product
Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:
skin reddening, blisters, rash

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
facial swelling, asthma (wheezing), hives, shock, skin reddening, rash, blisters

Drug Facts (continued)

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
if you have ever had an allergic reaction to acetaminophen or any other pain reliever
right before or after heart surgery

Ask a doctor before use if
you have liver disease
stomach bleeding warning applies to you
you have problems or serious side effects from taking pain relievers
you have a history of stomach problems, such as heartburn
you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
you are taking a diuretic

Ask a doctor or pharmacist before use if you are

Drug Facts (continued)

under a doctor's care for any serious condition
taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
taking any other drug

When using this product
take with food or milk if stomach upset occurs

Stop use and ask a doctor if
you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools
have stomach pain that does not get better
you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling
pain gets worse or lasts more than 10 days
redness or swelling is present in the painful area

Drug Facts (continued)

any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
do not take more than directed
adults and children 12 years and over: take 2 caplets every 8 hours while symptoms persist
children under 12 years: ask a doctor
do not take more than 6 caplets in 24 hours, unless directed by a doctor

Drug Facts (continued)

Other information
read all warnings and directions before use. (Keep carton. Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
Inactive ingredients: colloidal anhydrous silica, corn starch, croscarmellose sodium, glyceryl dibehenate, hypromellose, microcrystalline cellulose, polydextrose, povidone, red iron oxide, titanium dioxide, triacetin, yellow iron oxide

Questions or comments? Call 1-877-376-4271 (weekdays 9 AM to 5 PM)
Lot No.:
Exp Date:
Varnish Omit Area

Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.
NDC 2500008072
3125000108072
3
This product is not manufactured or distributed by Helsco US Holdings, LLC, owner of the registered trademark Advil® Dual Action with Acetaminophen.
Distributed by: Time-Sun Labs, Inc.
7 Mitchell Avenue, Farmingdale, NY 11735, USA
Mfg. Lic. No.: 600PHUS515 R31275

NDC 25000-080-72
Acetaminophen and Ibuprofen (NSAID) Tablets, 250 mg/125 mg

Pain Reliever
CONTAINS 2 MEDICINES
DUAL ACTION
18 Caplets\*
\*Capsule-Shaped Tablets
Marksans

Compare to the active ingredients in Advil® Dual Action with Acetaminophen†



ACTUAL SIZE

NDC 25000-080-72
Acetaminophen 250 mg and Ibuprofen 125 mg Tablets
18 count bottle label

### Drug Facts (continued)

not notice any signs or symptoms.

#### Directions

- do not take more than directed
- adults and children 12 years and over: take 2 caplets every 8 hours while symptoms persist
- children under 12 years: ask a doctor
- do not take more than 6 caplets in 24 hours, unless directed by a doctor

#### Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

#### Inactive ingredients

anhydrous silica, corn starch, croscarmellose sodium, glyceryl dibehenate, hypromellose, microcrystalline cellulose, polydextrose, povidone, red iron oxide, titanium dioxide, tracetin, yellow iron oxide

#### Questions or comments?

Call 1-877-376-4271 (weekdays 9 AM to 5 PM)

Distributed by:

Time-Cap Labs, Inc.

7 Michael Avenue,

Farmingdale, NY 11735, USA

Made in India

Mfg. Lic. No.: G0DRUSS515

R312/25

Adhesive Area

### Drug Facts (continued)

medical help right away.

**NSAID allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters.

If an allergic reaction occurs, stop use and seek medical help right away.

#### NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use

ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do

### Varnish Omit Area

Lot No.:  
Exp. Date:

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

acetaminophen

■ with other drugs containing

■ more than 9 caplets in 24 hours, which is the maximum daily amount for this product

■ 3 or more alcoholic drinks every day while using this product

**Acetaminophen allergy alert:** may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash.

if skin reaction occurs, stop use and seek

severe skin reactions. Symptoms may include:

# Acetaminophen 250 mg and Ibuprofen 125 mg Tablets

90 count carton label

**Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.**

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.**

### Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Ibuprofen 125 mg (NSAID)*	Pain reliever *nonsteroidal anti-inflammatory drug

### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - muscular aches
  - minor pain of arthritis

### Warnings

- Acetaminophen liver damage warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
  - with other drugs containing acetaminophen

### Drug Facts (continued)

- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product
- Acetaminophen allergy alert:** may cause severe skin reactions. Symptoms may include:
  - skin reddening
  - blisters
  - rash
- If skin reaction occurs, stop use and seek medical help right away.
- NSAID allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
  - hives
  - facial swelling
  - asthma (wheezing)
  - shock
  - skin reddening
  - rash
  - blisters
- If an allergic reaction occurs, stop use and seek medical help right away.

### NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

### Drug Facts (continued)

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to acetaminophen or any other pain reliever
- right before or after heart surgery

### Drug Facts (continued)

- Ask a doctor before use if**
  - you have liver disease
  - stomach bleeding warning applies to you
  - you have problems or serious side effects from taking pain relievers
  - you have a history of stomach problems, such as heartburn
  - you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
  - you are taking a diuretic

### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### When using this product

- take with food or milk if stomach upset occurs

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

### Drug Facts (continued)

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** do not take more than directed

### Drug Facts (continued)

adults and children 12 years and over	take 2 caplets every 8 hours while symptoms persist
children under 12 years	ask a doctor

- do not take more than 6 caplets in 24 hours, unless directed by a doctor

### Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients** colloidal anhydrous silica, corn starch, croscarmellose sodium, glyceryl dibehenate, hypromellose, microcrystalline cellulose, polydextrose, povidone, red iron oxide, titanium dioxide, triacetin, yellow iron oxide

**Questions or comments?** Call 1-877-376-4271 (weekdays 9 AM to 5 PM)

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.



†This product is not manufactured or distributed by Helicon US Holdings LLC, owner of the registered trademark Advil® Dual Action with Acetaminophen  
 Distributed by: Time-Cap Labs, Inc.  
 7 Michael Avenue, Farmingdale, NY 11735, USA  
 Made in India  
 Mfg. Lic. No.: GODRUGS515 R312/25

Lot No.:  
Exp Date:  
Varnish Omit Area

NDC 25000-080-07

## Acetaminophen and Ibuprofen (NSAID) Tablets, 250 mg/125 mg

Pain Reliever

CONTAINS 2 MEDICINES  
DUAL ACTION

90 Caplets\*

\*Capsule-Shaped Tablets



Compare to the active ingredients in Advil® Dual Action with Acetaminophen†



ACTUAL SIZE

NDC 25000-080-07

Acetaminophen 250 mg and Ibuprofen 125 mg Tablets

90 count bottle label

**Adhesive Area**

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>have had stomach ulcers or bleeding problems</li> <li>take a blood thinning (anticoagulant) or steroid drug</li> <li>take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> <li>have 3 or more alcoholic drinks every day while using this product</li> <li>take more or for a longer time than directed</li> </ul> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than directed</li> <li>adults and children 12 years and over: <b>take 2 caplets every 8 hours</b> while symptoms persist</li> <li>children under 12 years: ask a doctor</li> <li>do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>read all warnings and directions before use. Keep carton.</li> <li>store at 20-25°C (68-77°F)</li> <li>avoid excessive heat above 40°C (104°F)</li> </ul> <p><b>Inactive ingredients</b></p> <p>colloidal anhydrous silica, corn starch, croscarmellose sodium, glyceryl dibehenate, hypromellose, microcrystalline cellulose, polydextrose, povidone, red iron oxide, titanium dioxide, triacetin, yellow iron oxide</p> <p><b>Questions or comments?</b> Call 1-877-376-4271 (weekdays 9 AM to 5 PM)</p>
--

NDC 25000-080-07

**Acetaminophen and Ibuprofen (NSAID) Tablets, 250 mg/125 mg**

*Pain Reliever*

**CONTAINS 2 MEDICINES  
DUAL ACTION**

**90 Caplets\***  
\*Capsule-Shaped Tablets

  
**8 Hour Relief**

  
ACTUAL SIZE

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

**Do not use with other medicines containing ACETAMINOPHEN, can cause liver damage.**

<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each caplet)</b></p> <p>Acetaminophen 250 mg ..... Pain reliever</p> <p>Ibuprofen 125 mg (NSAID)* ..... Pain reliever</p> <p>*nonsteroidal anti-inflammatory drug</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves minor aches and pains due to:             <ul style="list-style-type: none"> <li>headache</li> <li>toothache</li> <li>backache</li> <li>menstrual cramps</li> <li>muscular aches</li> <li>minor pain of arthritis</li> </ul> </li> </ul> <p><b>Warnings</b></p> <p><b>Acetaminophen liver damage warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take:         <ul style="list-style-type: none"> <li>with other drugs containing acetaminophen</li> <li>more than 6 caplets in 24 hours, which is the maximum daily amount for this product</li> </ul> </p> <p><b>Acetaminophen allergy alert:</b> may cause severe skin reactions. Symptoms may include:         <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul> </p> <p><b>NSAID allergy alert:</b> ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:         <ul style="list-style-type: none"> <li>facial swelling</li> <li>asthma (wheezing)</li> <li>hives</li> <li>skin reddening</li> <li>rash</li> <li>blisters</li> <li>stomach</li> <li>skin reddening</li> </ul> </p> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>NSAID stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:         <ul style="list-style-type: none"> <li>are age 60 or older</li> </ul> </p>	<p>Lot No.:</p> <p>Exp. Date:</p> <p style="text-align: center;"><b>FREE HERE</b></p> <p style="text-align: center;"><b>Varnish Omit Area</b></p> <p style="text-align: right;">R31225 Mfg. Lic. No.: 6006UC5514</p>
--	--

Distributed by: Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale, NY 11735, USA  
Made in India

NDC 25000-080-74

Acetaminophen 250 mg and Ibuprofen 125 mg Tablets

288 count bottle label



**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>Acetaminophen</b> (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	250 mg
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	125 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>GLYCERYL DIBEHENATE</b> (UNII: R8WTH25YS2)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

**Product Characteristics**

<b>Color</b>	YELLOW	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	80
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:25000-080-72	1 in 1 CARTON	07/11/2023	
1		18 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:25000-080-07	1 in 1 CARTON	07/11/2023	
2		90 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:25000-080-74	288 in 1 BOTTLE; Type 0: Not a Combination Product	07/11/2023	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA216994	07/11/2023	

**Labeler** - MARKSANS PHARMA LIMITED (925822975)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
MARKSANS PHARMA LIMITED		925822975	MANUFACTURE(25000-080)

Revised: 12/2025

MARKSANS PHARMA LIMITED