

ASPIRIN AND DIPYRIDAMOLE- aspirin and dipyridamole capsule, extended release

NorthStar Rx LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ASPIRIN AND EXTENDED-RELEASE DIPYRIDAMOLE CAPSULES safely and effectively. See full prescribing information for ASPIRIN AND EXTENDED-RELEASE DIPYRIDAMOLE CAPSULES.

ASPIRIN and extended-release DIPYRIDAMOLE capsules, for oral use
Initial U.S. Approval: 1999

----- **INDICATIONS AND USAGE** -----

Aspirin and extended-release dipyridamole capsules are a combination of aspirin and dipyridamole, antiplatelet agents, indicated to reduce the risk of stroke in patients who have had transient ischemia of the brain or completed ischemic stroke due to thrombosis (1)

----- **DOSAGE AND ADMINISTRATION** -----

- One capsule twice daily (morning and evening) with or without food (2)
- In case of intolerable headaches during initial treatment, switch to one capsule at bedtime and low-dose aspirin in the morning; resume BID dosing within one week (2.1)
- Do not chew capsule (2)
- Not interchangeable with the individual components of aspirin and dipyridamole tablets(2)
- Dispense in this unit-of-use container (16)

----- **DOSAGE FORMS AND STRENGTHS** -----

- Capsule: 25 mg aspirin and 200 mg extended-release dipyridamole (3)

----- **CONTRAINDICATIONS** -----

- Hypersensitivity to any product ingredients (4.1)
- Patients with known allergy to NSAIDs (4.2)
- Patients with the syndrome of asthma, rhinitis, and nasal polyps (4.2)

----- **WARNINGS AND PRECAUTIONS** -----

- Aspirin and extended-release dipyridamole capsules increases the risk of bleeding (5.1)
- Avoid use in patients with severe hepatic or renal insufficiency (5.2, 5.3)
- Interrupt aspirin and extended-release dipyridamole capsules 48 hours before using intravenous dipyridamole or other adenosinergic agents for stress testing (5.6, 7.1)

----- **ADVERSE REACTIONS** -----

- The most frequently reported adverse reactions (>10% and greater than placebo) were headache, dyspepsia, abdominal pain, nausea, and diarrhea (6)

To report SUSPECTED ADVERSE REACTIONS, contact NorthStar RxLLC at 1-800-206-7821 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **DRUG INTERACTIONS** -----

- Co-administration with anticoagulants, antiplatelets, or NSAIDs can increase risk of bleeding (7.1)
- Decreased renal function can occur with co-administration with NSAIDs (7.1)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 5/2021

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Alternative Regimen in Case of Intolerable Headaches

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

4.1 Hypersensitivity

4.2 Allergy

4.3 Reye Syndrome

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Bleeding

5.2 Renal Failure

5.3 Hepatic Insufficiency

5.4 Coronary Artery Disease

5.5 Hypotension

5.6 Stress Testing with Intravenous Dipyridamole and Other Adenosinergic Agents

5.7 General

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Post-Marketing Experience

7 DRUG INTERACTIONS

7.1 Drug Interaction Study Information Obtained From Literature

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Patients with Severe Hepatic or Severe Renal Dysfunction

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment Of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Aspirin and extended-release dipyridamole capsules are indicated to reduce the risk of stroke in patients who have had transient ischemia of the brain or completed ischemic stroke due to thrombosis.

2 DOSAGE AND ADMINISTRATION

Aspirin and extended-release dipyridamole capsules are not interchangeable

with the individual components of aspirin and dipyridamole tablets.

The recommended dose of aspirin and extended-release dipyridamole capsules is one capsule given orally twice daily, one in the morning and one in the evening. Swallow capsules whole without chewing. Aspirin and extended-release dipyridamole capsules can be administered with or without food.

2.1 Alternative Regimen in Case of Intolerable Headaches

In the event of intolerable headaches during initial treatment, switch to one capsule at bedtime and low-dose aspirin in the morning. Because there are no outcome data with this regimen and headaches become less of a problem as treatment continues, patients should return to the usual regimen as soon as possible, usually within one week.

3 DOSAGE FORMS AND STRENGTHS

25 mg aspirin and 200 mg extended-release dipyridamole capsule contains yellow to dark yellow colored extended-release pellets incorporating dipyridamole and white to off white round biconvex film coated tablet incorporating immediate-release aspirin, in size '0xEL' hard gelatin capsule with orange colored cap and white to off white colored body, imprinted '339' on the body and 'RDY' on cap with black ink.

4 CONTRAINDICATIONS

4.1 Hypersensitivity

Aspirin and extended-release dipyridamole capsules are contraindicated in patients with known hypersensitivity to any of the product components.

4.2 Allergy

Aspirin is contraindicated in patients with known allergy to nonsteroidal anti-inflammatory drug (NSAID) products and in patients with the syndrome of asthma, rhinitis, and nasal polyps. Aspirin may cause severe urticaria, angioedema or bronchospasm.

4.3 Reye Syndrome

Do not use aspirin in children or teenagers with viral infections because of the risk of Reye syndrome.

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Bleeding

Aspirin and extended-release dipyridamole capsules increases the risk of bleeding. Risk factors for bleeding include the use of other drugs that increase the risk of bleeding (e.g., anticoagulants, antiplatelet agents, heparin, anagrelide, fibrinolytic therapy, and chronic use of NSAIDs) [see *Drug Interactions* (7.1)].

Intracranial Hemorrhage

In European Stroke Prevention Study-2 (ESPS2), the annualized event rate for intracranial hemorrhage was 0.39%/year in the aspirin and extended-release dipyridamole capsules group, 0.26%/year in the extended-release dipyridamole (ER-DP) group, 0.24%/year in the aspirin (ASA) group, and 0.29%/year in the placebo groups.

Gastrointestinal (GI) Side Effects

GI side effects include stomach pain, heartburn, nausea, vomiting, and gross GI bleeding. Although minor upper GI symptoms, such as dyspepsia, are common and can occur anytime during therapy, physicians should remain alert for signs of ulceration and bleeding, even in the absence of previous GI symptoms. Inform patients about the signs and symptoms of GI side effects and what steps to take if they occur.

In ESPS2, the annualized event rate for gastrointestinal bleeding was 2.97%/year in the aspirin and extended-release dipyridamole capsules group, 1.58%/year in the extended-release dipyridamole group, 2.06%/year in the aspirin group, and 1.40%/year in the placebo groups.

Peptic Ulcer Disease

Avoid using aspirin in patients with a history of active peptic ulcer disease, which can cause gastric mucosal irritation and bleeding.

Alcohol Warning

Because aspirin and extended-release dipyridamole capsules contains aspirin, counsel patients who consume three or more alcoholic drinks every day about the bleeding risks involved with chronic, heavy alcohol use while taking aspirin.

5.2 Renal Failure

Avoid aspirin in patients with severe renal failure (glomerular filtration rate less than 10 mL/minute) [see *Use in Specific Populations (8.6)* and *Clinical Pharmacology (12.3)*].

5.3 Hepatic Insufficiency

Elevations of hepatic enzymes and hepatic failure have been reported in association with dipyridamole administration [see *Use in Specific Populations (8.6)* and *Clinical Pharmacology (12.3)*].

5.4 Coronary Artery Disease

Dipyridamole has a vasodilatory effect. Chest pain may be precipitated or aggravated in patients with underlying coronary artery disease who are receiving dipyridamole.

For stroke or TIA patients for whom aspirin is indicated to prevent recurrent myocardial infarction (MI) or angina pectoris, the aspirin in this product may not provide adequate treatment for the cardiac indications.

5.5 Hypotension

Dipyridamole produces peripheral vasodilation, which can exacerbate pre-existing hypotension.

5.6 Stress Testing with Intravenous Dipyridamole and Other Adenosinergic

Agents

Clinical experience suggests that patients being treated with aspirin and extended-release dipyridamole who also require pharmacological stress testing with intravenous dipyridamole or other adenosinergic agents (e.g. adenosine, regadenoson) should interrupt aspirin and extended-release dipyridamole capsules for 48 hours prior to stress testing [see *Drug Interactions (7.1)*].

Intake of aspirin and extended-release dipyridamole capsules within 48 hours prior to stress testing with intravenous dipyridamole or other adenosinergic agents may increase the risk for cardiovascular side effects of these agents and may impair the sensitivity of the test.

5.7 General

Aspirin and extended-release dipyridamole capsules are not interchangeable with the individual components of aspirin and dipyridamole tablets.

6 ADVERSE REACTIONS

The following adverse reactions are discussed elsewhere in the labeling:

- Hypersensitivity [see *Contraindications (4.1)*]
- Allergy [see *Contraindications (4.2)*]
- Risk of Bleeding [see *Warnings and Precautions (5.1)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The efficacy and safety of aspirin and extended-release dipyridamole capsules was established in the European Stroke Prevention Study-2 (ESPS2). ESPS2 was a double-blind, placebo-controlled study that evaluated 6,602 patients over the age of 18 years who had a previous ischemic stroke or transient ischemic attack within ninety days prior to entry. Patients were randomized to either aspirin and extended-release dipyridamole capsules, aspirin, ER-DP, or placebo [see *Clinical Studies (14)*]; primary endpoints included stroke (fatal or nonfatal) and death from all causes.

This 24-month, multicenter, double-blind, randomized study (ESPS2) was conducted to compare the efficacy and safety of aspirin and extended-release dipyridamole capsules with placebo, extended-release dipyridamole alone and aspirin alone. The study was conducted in a total of 6,602 male and female patients who had experienced a previous ischemic stroke or transient ischemia of the brain within three months prior to randomization.

Table 1 presents the annualized event rate for adverse events that occurred in 1%/year or more of patients treated with aspirin and extended-release dipyridamole capsules where the incidence was also at least 1%/year greater than in those patients treated with placebo. There is no clear benefit of the dipyridamole/aspirin combination over aspirin with respect to safety.

Table 1 Incidence of Adverse Events in ESPS2^a

Body System/Preferred Term	Individual Treatment Group	ER-DP Alone n(%/year) ^b	ASA Alone n(%/year) ^b	Placebo n(%/year) ^b
	Aspirin and Extended-Release Dipyridamole Capsules n (%/year) ^b			
Total Number of Patients	1650	1654	1649	1649
Central and Peripheral Nervous System Disorders				
Headache	647 (28.25)	634 (27.91)	558 (22.10)	543 (22.29)
Gastrointestinal System Disorders				
Dyspepsia	303 (13.23)	288 (12.68)	299 (11.84)	275 (11.29)
Abdominal Pain	289 (12.62)	255 (11.22)	262 (10.38)	239 (9.81)
Nausea	264 (11.53)	254 (11.18)	210 (8.32)	232 (9.53)
Diarrhea	210 (9.17)	257 (11.31)	112 (4.44)	161 (6.61)
Vomiting	138 (6.03)	129 (5.68)	101 (4.00)	118 (4.84)
Platelet, Bleeding and Clotting Disorders				
Hemorrhage NOS	52 (2.27)	24 (1.06)	46 (1.82)	24 (0.99)

^aReported by $\geq 1\%$ /year of patients during aspirin and extended-release dipyridamole capsules treatment where the incidence was at least 1%/year greater than in those treated with placebo.

^bAnnual event rate per 100 pt-years = $100 \times$ number of subjects with event/subject-years. Subject-years is defined as cumulative number of days on treatment divided by 365.25.

Note: ER-DP = extended-release dipyridamole 200 mg; ASA = aspirin 25 mg. The dosage regimen for all treatment groups is BID

NOS = not otherwise specified.

Discontinuation due to adverse events in ESPS2 was 25% for aspirin and extended-release dipyridamole capsules, 25% for extended-release dipyridamole, 19% for aspirin, and 21% for placebo (refer to Table 2).

Table 2 Incidence of Adverse Events that Led to the Discontinuation of Treatment^a

	Treatment Groups			
	Aspirin and Extended-Release Dipyridamole Capsules^a (%)^b	ER-DP (%/year)^b	ASA n (%/year)^b	Placebo^a (%/year)^b
Total Number of Patients	1650	1654	1649	1649
Patients with at least one Adverse Event that led to treatment discontinuation	417 (18.21)	419 (18.44)	318 (12.59)	352 (14.45)
Headache	165 (7.20)	166 (7.31)	57 (2.26)	69 (2.83)
Nausea	91 (3.97)	95 (4.18)	51 (2.02)	53 (2.18)
Abdominal Pain	74 (3.23)	64 (2.82)	56 (2.22)	52 (2.13)
Vomiting	53 (2.31)	52 (2.29)	28 (1.11)	24 (0.99)

^a Reported by $\geq 1\%$ /year of patients during aspirin and extended-release dipyridamole capsules treatment where the incidence was at least 1%/year greater than in those treated with placebo.

^b Annual event rate per 100 pt-years = $100 \times$ number of subjects with event/subject-years. Subject-years is defined as cumulative number of days on treatment divided by 365.25.

Note: ER-DP = extended-release dipyridamole 200 mg; ASA = aspirin 25 mg. The dosage regimen for all treatment groups is BID.

Headache was most notable in the first month of treatment.

6.2 Post-Marketing Experience

The following is a list of additional adverse reactions that have been reported either in the literature or are from post-marketing spontaneous reports for either dipyridamole or aspirin. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to estimate reliably their frequency or establish a causal relationship to drug exposure. Decisions to include these reactions in labeling are typically based on one or more of the following factors: (1) seriousness of the reaction, (2) frequency of reporting, or (3) strength of causal connection to aspirin and extended-release dipyridamole capsules.

Body as a Whole: Hypothermia, chest pain, allergic reaction, syncope

Cardiovascular: Angina pectoris, hypotension

Central Nervous System: Cerebral edema, dizziness, cerebral hemorrhage, intracranial hemorrhage, subarachnoid hemorrhage

Fluid and Electrolyte: Hyperkalemia, metabolic acidosis, respiratory alkalosis, hypokalemia

Gastrointestinal: Pancreatitis, Reye syndrome, hematemesis, gastritis, ulceration and perforation, hemorrhage rectum, melena, GI hemorrhage

Hearing and Vestibular Disorders: Hearing loss
Heart Rate and Rhythm Disorders: Tachycardia, palpitation

Immune System Disorders: Hypersensitivity, acute anaphylaxis, laryngeal edema
Liver and Biliary System Disorders: Hepatitis, hepatic failure, cholelithiasis, jaundice, hepatic

function abnormal

Musculoskeletal: Rhabdomyolysis, myalgia

Metabolic and Nutritional Disorders: Hypoglycemia, dehydration

Platelet, Bleeding and Clotting Disorders: Prolongation of the prothrombin time, disseminated intravascular coagulation, coagulopathy, thrombocytopenia, hematoma, gingival bleeding, epistaxis, purpura

Psychiatric Disorders: Confusion, agitation

Respiratory: Tachypnea, dyspnea, hemoptysis

Skin and Appendages Disorders: Rash, alopecia, angioedema, Stevens-Johnson syndrome, skin hemorrhages such as bruising, ecchymosis, and hematoma, pruritus, urticaria, and drug reaction with eosinophilia and systemic symptoms (DRESS)

Urogenital: Interstitial nephritis, papillary necrosis, proteinuria, renal insufficiency and failure, hematuria

Vascular (Extracardiac) Disorders: Allergic vasculitis, flushing

Other Adverse Events: Anorexia, aplastic anemia, migraine, pancytopenia, thrombocytosis.

7 DRUG INTERACTIONS

7.1 Drug Interaction Study Information Obtained From Literature

Adenosinergic agents (e.g. adenosine, regadenoson)

Dipyridamole has been reported to increase the plasma levels and cardiovascular effects of adenosine. Adjustment of adenosine dosage may be necessary. Dipyridamole also increases the cardiovascular effects of regadenoson, an adenosine A_{2A}-receptor agonist. The potential risk of cardiovascular side effects with intravenous adenosinergic agents may be increased during the testing period when dipyridamole is not held 48 hours prior to stress testing.

Angiotensin Converting Enzyme (ACE) Inhibitors

Because of the indirect effect of aspirin on the renin-angiotensin conversion pathway, the hyponatremic and hypotensive effects of ACE inhibitors may be diminished by concomitant administration of aspirin.

Acetazolamide

Concurrent use of aspirin and acetazolamide can lead to high serum concentrations of acetazolamide (and toxicity) due to competition at the renal tubule for secretion.

Anticoagulants and Antiplatelets

Patients taking aspirin and extended-release dipyridamole capsules in combination with anticoagulants, antiplatelets, or any substance impacting coagulation are at increased risk for bleeding. Aspirin can displace warfarin from protein binding sites, leading to prolongation of both the prothrombin time and the bleeding time. Aspirin can increase the anticoagulant activity of heparin, increasing bleeding risk.

Anagrelide

Patients taking aspirin in combination with anagrelide are at an increased risk of bleeding.

Anticonvulsants

Salicylic acid can displace protein-bound phenytoin and valproic acid, leading to a decrease in the total concentration of phenytoin and an increase in serum valproic acid levels.

Beta Blockers

The hypotensive effects of beta blockers may be diminished by the concomitant administration of aspirin due to inhibition of renal prostaglandins, leading to decreased renal blood flow and salt and fluid retention.

Cholinesterase Inhibitors

Dipyridamole may counteract the anticholinesterase effect of cholinesterase inhibitors, thereby potentially aggravating myasthenia gravis.

Diuretics

The effectiveness of diuretics in patients with underlying renal or cardiovascular disease may be diminished by the concomitant administration of aspirin due to inhibition of renal prostaglandins, leading to decreased renal blood flow and salt and fluid retention.

Methotrexate

Salicylate can inhibit renal clearance of methotrexate, leading to bone marrow toxicity, especially in the elderly or renal impaired.

Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)

The concurrent use of aspirin with other NSAIDs may increase bleeding or lead to decreased renal function.

Oral Hypoglycemics

Moderate doses of aspirin may increase the effectiveness of oral hypoglycemic drugs, leading to hypoglycemia.

Uricosuric Agents (probenecid and sulfinpyrazone)

Salicylates antagonize the uricosuric action of uricosuric agents.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Available data from published studies and postmarketing experience with aspirin and extended-release dipyridamole capsules use during pregnancy have not identified a clear association between aspirin and extended-release dipyridamole capsules use and major birth defects, miscarriage, or adverse maternal or fetal outcomes (see Data). Aspirin and extended-release dipyridamole capsules contains low-dose aspirin which is an NSAID (see Clinical Considerations). In animal reproduction studies, there were adverse

developmental effects with administration of aspirin in rats and rabbits at doses about 66 and 44 times, respectively, the human exposure at the maximum recommended daily dose of aspirin-dipyridamole. Reproduction studies with dipyridamole in mice, rabbits, and rats have revealed no evidence of harm to the fetus up to doses about 25 times the maximum recommended daily human dose of aspirin-dipyridamole. Nonclinical data are suggestive of a possible potentiation of aspirin-related fetal toxicity when combined with dipyridamole (*see Data*).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4 and 15 to 20%, respectively.

Clinical Considerations

Labor and Delivery

Aspirin and extended-release dipyridamole capsules, which contains dipyridamole and low-dose aspirin, increases the risk for bleeding [*see Warnings and Precautions (5.1)*]. Maternal use of high-dose aspirin can result in excessive blood loss at delivery, prolonged gestation, prolonged labor, intracranial hemorrhage in premature infants, low birth weight, stillbirth, and neonatal death.

Data

Human Data

Published data from clinical trials, observational studies, case series, and case reports over several decades have not identified a clear association between aspirin-dipyridamole use in pregnancy and major birth defects, miscarriage, or adverse maternal or fetal outcomes. However, these studies cannot definitively establish the absence of any aspirin-dipyridamole associated risks. Methodological limitations of these studies include variability in the timing and dose of drug exposure (e.g., most exposures occurred beyond the first trimester) and the small sample sizes of individual studies.

Animal Data

Aspirin has been shown to be teratogenic in rats (spina bifida, exencephaly, microphthalmia and coelosomia) and rabbits (congested fetuses, agenesis of skull and upper jaw, generalized edema with malformation of the head, and diaphanous skin) at oral doses of 330 mg/kg/day and 110 mg/kg/day, respectively. These doses, which also resulted in a high resorption rate in rats (63% of implantations versus 5% in controls), are, on a mg/m² basis, about 66 and 44 times, respectively, the dose of aspirin contained in the maximum recommended daily human dose of aspirin-dipyridamole. Reproduction studies with dipyridamole have been performed in mice, rabbits and rats at oral doses of up to 125 mg/kg, 40 mg/kg, and 1,000 mg/kg, respectively (about 1½, 2, and 25 times the maximum recommended daily human oral dose, respectively, on a mg/m² basis) and have revealed no evidence of harm to the fetus due to dipyridamole. When 330 mg aspirin/kg/day was combined with 75 mg dipyridamole/kg/day in the rat at doses about 66 and 2 times, respectively, the maximum recommended daily human dose, the resorption rate approached 100%.

8.2 Lactation

Risk Summary

Based on data from a clinical lactation study in breastfeeding women taking low-dose aspirin, the metabolite salicylic acid is present in human milk in low levels (see Data). Dipyridamole is also present in human milk. There is no information on the effects of aspirin and extended-release dipyridamole capsules or dipyridamole on the breastfed infant or on milk production. There is insufficient information to determine the effects of aspirin on the breastfed infant and no information on the effects of aspirin on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for aspirin and extended-release dipyridamole capsules and any potential adverse effects on the breastfed child from aspirin and extended-release dipyridamole capsules or from the underlying maternal condition.

Data

A published clinical study involved six exclusively breastfeeding women at 1 to 8 months postpartum who were taking 81 mg aspirin daily. Milk samples were collected at steady state, at 0, 1, 2, 4, 8, 12, and 24 hours after taking a dose of aspirin. Aspirin was undetectable in human milk. Salicylic acid was present in milk at low levels (average concentration of 24 ng/mL). Based on an average milk consumption of 150 mL/kg/day, the calculated relative infant dose was 0.4%. No adverse effects on the breastfed infants were noted.

8.4 Pediatric Use

Safety and effectiveness of aspirin and extended-release dipyridamole capsules in pediatric patients have not been studied. Because of the aspirin component, use of this product in the pediatric population is not recommended [*see Contraindications (4.3)*].

8.5 Geriatric Use

Of the total number of subjects in ESPS2, 61% were 65 and over, while 27% were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out [*see Clinical Pharmacology (12.3)*].

8.6 Patients with Severe Hepatic or Severe Renal Dysfunction

Aspirin and extended-release dipyridamole capsules have not been studied in patients with hepatic or renal impairment. Avoid using aspirin containing products, such as aspirin and extended-release dipyridamole capsules in patients with severe hepatic or severe renal (glomerular filtration rate < 10 mL/min) dysfunction [*see Warnings and Precautions (5.2, 5.3) and Clinical Pharmacology (12.3)*].

10 OVERDOSAGE

Because of the dose ratio of dipyridamole to aspirin, overdosage of aspirin and extended-release dipyridamole capsules are likely to be dominated by signs and symptoms of dipyridamole overdose. In case of real or suspected overdose, seek

medical attention or contact a Poison Control Center immediately. Careful medical management is essential.

Based upon the known hemodynamic effects of dipyridamole, symptoms such as warm feeling, flushes, sweating, restlessness, feeling of weakness, and dizziness may occur. A drop in blood pressure and tachycardia might also be observed.

Salicylate toxicity may result from acute ingestion (overdose) or chronic intoxication. Severity of aspirin intoxication is determined by measuring the blood salicylate level. The early signs of salicylic overdose (salicylism), including tinnitus (ringing in the ears), occur at plasma concentrations approaching 200 mg/mL. In severe cases, hyperthermia and hypovolemia are the major immediate threats to life. Plasma concentrations of aspirin above 300 mg/mL are clearly toxic. Severe toxic effects are associated with levels above 400 mg/mL. A single lethal dose of aspirin in adults is not known with certainty but death may be expected at 30 grams.

Treatment of overdose consists primarily of supporting vital functions, increasing drug elimination, and correcting acid-base disturbances. Consider gastric emptying and/or lavage as soon as possible after ingestion, even if the patient has vomited spontaneously. After lavage and/or emesis, administration of activated charcoal as a slurry may be beneficial if less than 3 hours have passed since ingestion. Charcoal absorption should not be employed prior to emesis and lavage. Follow acid-base status closely with serial blood gas and serum pH measurements. Maintain fluid and electrolyte balance. Administer replacement fluid intravenously and augment with correction of acidosis. Treatment may require the use of a vasopressor. Infusion of glucose may be required to control hypoglycemia.

Administration of xanthine derivatives (e.g., aminophylline) may reverse the vasodilatory effects of dipyridamole overdose. Plasma electrolytes and pH should be monitored serially to promote alkaline diuresis of salicylate if renal function is normal. In patients with renal insufficiency or in cases of life-threatening intoxication, dialysis is usually required to treat salicylic overdose; however, since dipyridamole is highly protein bound, dialysis is not likely to remove dipyridamole. Exchange transfusion may be indicated in infants and young children.

11 DESCRIPTION


Aspirin and extended-release dipyridamole capsules are a combination of aspirin and dipyridamole, antiplatelet agents, intended for oral administration. Each hard gelatin capsule contains 200 mg dipyridamole in an extended-release form and 25 mg aspirin, as an immediate-release film-coated tablet. In addition, each capsule contains the following inactive ingredients: acacia, anhydrous lactose, colloidal silicon dioxide, dimethicone, hypromellose, hypromellose phthalate, lecithin, methacrylic acid copolymer, microcrystalline cellulose, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, tartaric acid, titanium dioxide, triacetin and xanthan gum.

Each capsule shell contains FD&C yellow 6, gelatin, sodium lauryl sulfate, titanium dioxide and yellow iron oxide.

Imprinting ink contains black iron oxide, potassium hydroxide, propylene glycol, shellac and strong ammonia solution.

Dipyridamole


Dipyridamole is an antiplatelet agent chemically described as 2,2',2'',2'''-[(4,8-Dipiperidinopyrimido[5,4-d]pyrimidine-2,6-diyl)dinitrilo]-tetraethanol. It has the following structural formula:



Dipyridamole USP is an intensely yellow, crystalline powder or needles. It is practically insoluble in water, sparingly soluble in ethyl alcohol, very slightly soluble in acetone and ethyl acetate.

Aspirin

The antiplatelet agent aspirin (acetylsalicylic acid) is chemically known as benzoic acid, 2-(acetyloxy)-, and has the following structural formula:



$C_9H_8O_4$

Mol. Wt. 180.16

Aspirin USP is a white crystals, commonly tabular or needle like, or white, crystalline powder. Slightly soluble in water, freely soluble in alcohol, soluble in chloroform and in ether, sparingly soluble in absolute ether.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The antithrombotic action of aspirin and extended-release dipyridamole capsules is the result of the additive antiplatelet effects of dipyridamole and aspirin.

Dipyridamole

Dipyridamole inhibits the uptake of adenosine into platelets, endothelial cells and erythrocytes *in vitro* and *in vivo*; the inhibition occurs in a dose-dependent manner at

therapeutic concentrations (0.5 to 1.9 mcg/mL). This inhibition results in an increase in local concentrations of adenosine which acts on the platelet A₂-receptor thereby stimulating platelet adenylate cyclase and increasing platelet cyclic-3',5'-adenosine monophosphate (cAMP) levels. Via this mechanism, platelet aggregation is inhibited in response to various stimuli such as platelet activating factor (PAF), collagen and adenosine diphosphate (ADP).

Dipyridamole inhibits phosphodiesterase (PDE) in various tissues. While the inhibition of cAMP-PDE is weak, therapeutic levels of dipyridamole inhibit cyclic-3',5'-guanosine monophosphate-PDE (cGMP-PDE), thereby augmenting the increase in cGMP produced by EDRF (endothelium-derived relaxing factor, now identified as nitric oxide).

Aspirin

Aspirin inhibits platelet aggregation by irreversible inhibition of platelet cyclooxygenase and thus inhibits the generation of thromboxane A₂, a powerful inducer of platelet aggregation and vasoconstriction.

12.2 Pharmacodynamics

The effect of either agent on the other's inhibition of platelet reactivity has not been evaluated.

12.3 Pharmacokinetics

There are no significant interactions between aspirin and dipyridamole. The kinetics of the components are unchanged by their co-administration as aspirin and extended-release dipyridamole capsules.

Absorption

Dipyridamole

Peak plasma levels of dipyridamole are achieved 2 hours (range 1 to 6 hours) after administration of a daily dose of 400 mg aspirin and extended-release dipyridamole capsules (given as 200 mg BID). The peak plasma concentration at steady-state is 1.98 mcg/mL (1.01 to 3.99 mcg/mL) and the steady-state trough concentration is 0.53 mcg/mL (0.18 to 1.01 mcg/mL).

Aspirin

Peak plasma levels of aspirin are achieved 0.63 hours (0.5 to 1 hour) after administration of a 50 mg aspirin daily dose from aspirin and extended-release dipyridamole capsules (given as 25 mg BID). The peak plasma concentration at steady-state is 319 ng/mL (175 to 463 ng/mL). Aspirin undergoes moderate hydrolysis to salicylic acid in the liver and the gastrointestinal wall, with 50% to 75% of an administered dose reaching the systemic circulation as intact aspirin.

Effect of Food

Dipyridamole

When aspirin and dipyridamole extended-release capsules were taken with a high fat meal, dipyridamole peak plasma levels (C_{max}) and total absorption (AUC) were decreased at steady-state by 20 to 30% compared to fasting. Due to the similar degree of inhibition of adenosine uptake at these plasma concentrations, this food effect is not considered

clinically relevant.

Aspirin

When aspirin and extended-release dipyridamole capsules were taken with a high fat meal, there was no difference for aspirin in AUC at steady-state, and the approximately 50% decrease in C_{max} was not considered clinically relevant based on a similar degree of cyclooxygenase inhibition comparing the fed and fasted state.

Distribution

Dipyridamole

Dipyridamole is highly lipophilic ($\log P=3.71$, $pH=7$); however, it has been shown that the drug does not cross the blood-brain barrier to any significant extent in animals. The steady-state volume of distribution of dipyridamole is about 92 L. Approximately 99% of dipyridamole is bound to plasma proteins, predominantly to alpha 1-acid glycoprotein and albumin.

Aspirin

Aspirin is poorly bound to plasma proteins and its apparent volume of distribution is low (10 L). Its metabolite, salicylic acid, is highly bound to plasma proteins, but its binding is concentration-dependent (nonlinear). At low concentrations (<100 mcg/mL), approximately 90% of salicylic acid is bound to albumin. Salicylic acid is widely distributed to all tissues and fluids in the body, including the central nervous system, breast milk, and fetal tissues. Early signs of salicylate overdose (salicylism), including tinnitus (ringing in the ears), occur at plasma concentrations approximating 200 mcg/mL [see Overdosage (10)].

Metabolism and Elimination

Dipyridamole

Dipyridamole is metabolized in the liver, primarily by conjugation with glucuronic acid, of which monoglucuronide which has low pharmacodynamic activity is the primary metabolite. In plasma, about 80% of the total amount is present as parent compound and 20% as monoglucuronide. Most of the glucuronide metabolite (about 95%) is excreted via bile into the feces, with some evidence of enterohepatic circulation. Renal excretion of parent compound is negligible and urinary excretion of the glucuronide metabolite is low (about 5%). With intravenous (i.v.) treatment of dipyridamole, a triphasic profile is obtained: a rapid alpha phase, with a half-life of about 3.4 minutes, a beta phase, with a half-life of about 39 minutes, (which, together with the alpha phase accounts for about 70% of the total area under the curve, AUC) and a prolonged elimination phase λ_z with a half-life of about 15.5 hours. Because of the extended absorption phase of the dipyridamole component, only the terminal phase is apparent from oral treatment with aspirin and extended-release dipyridamole capsules which was 13.6 hours.

Aspirin

Aspirin is rapidly hydrolyzed in plasma to salicylic acid, with a half-life of 20 minutes. Plasma levels of aspirin are essentially undetectable 2 to 2.5 hours after dosing and peak salicylic acid concentrations occur 1 hour (range: 0.5 to 2 hours) after administration of aspirin. Salicylic acid is primarily conjugated in the liver to form salicyluric acid, a phenolic glucuronide, an acyl glucuronide, and a number of minor metabolites. Salicylate

metabolism is saturable and total body clearance decreases at higher serum concentrations due to the limited ability of the liver to form both salicylic acid and phenolic glucuronide. Following toxic doses (10 to 20 grams), the plasma half-life may be increased to over 20 hours.

The elimination of acetylsalicylic acid follows first-order kinetics with aspirin and extended-release dipyridamole capsules and has a half-life of 0.33 hours. The half-life of salicylic acid is 1.71 hours. Both values correspond well with data from the literature at lower doses which state a resultant half-life of approximately 2 to 3 hours. At higher doses, the elimination of salicylic acid follows zero-order kinetics (i.e., the rate of elimination is constant in relation to plasma concentration), with an apparent half-life of 6 hours or higher. Renal excretion of unchanged drug depends upon urinary pH. As urinary pH rises above 6.5, the renal clearance of free salicylate increases from <5% to >80%. Alkalinization of the urine is a key concept in the management of salicylate overdose [see Overdosage (10)]. Following therapeutic doses, about 10% is excreted as salicylic acid and 75% as salicylic acid, as the phenolic and acyl glucuronides, in urine.

Specific Populations

Geriatric Patients:

Dipyridamole

In ESPS2 [see Clinical Studies (14)], plasma concentrations (determined as AUC) of dipyridamole in healthy elderly subjects (>65 years) were about 40% higher than in subjects younger than 55 years receiving treatment with aspirin and extended-release dipyridamole capsules.

Hepatic Dysfunction:

No study has been conducted with aspirin and extended-release dipyridamole capsules in patients with hepatic dysfunction.

Dipyridamole

In a study conducted with an intravenous formulation of dipyridamole, patients with mild to severe hepatic insufficiency showed no change in plasma concentrations of dipyridamole but showed an increase in the pharmacologically inactive monoglucuronide metabolite. Dipyridamole can be dosed without restriction as long as there is no evidence of hepatic failure.

Aspirin

Avoid aspirin in patients with severe hepatic insufficiency

Renal Dysfunction:

Dipyridamole

In ESPS2 patients [see Clinical Studies (14)], with creatinine clearances ranging from about 15 mL/min to >100 mL/min, no changes were observed in the pharmacokinetics of dipyridamole or its glucuronide metabolite if data were corrected for differences in age.

Aspirin

Avoid aspirin in patients with severe renal failure (glomerular filtration rate <10 mL/min).

Drug Interaction Studies

A dedicated drug interaction study was conducted in 60 healthy volunteers to evaluate the effects of omeprazole 80 mg administered once daily on the pharmacokinetics (PK) of dipyridamole and the pharmacodynamics (PD) of acetylsalicylic acid when co-administered with aspirin and extended-release dipyridamole capsules twice daily. Dipyridamole exposure (C_{max} and AUC) at steady-state were similar with or without omeprazole co-administration. The pharmacokinetics of acetylsalicylic acid was not characterized. However, the antiplatelet activity as measured by arachidonic acid induced platelet aggregation was similar between the treatment arms at steady-state.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment Of Fertility

In studies in which dipyridamole was administered in the feed to mice (up to 111 weeks in males and females) and rats (up to 128 weeks in males and up to 142 weeks in females), there was no evidence of drug-related carcinogenesis. The highest dose administered in these studies (75 mg/kg/day) was, on a mg/m^2 basis, about equivalent to the maximum recommended daily human oral dose (MRHD) in mice and about twice the MRHD in rats.

Combinations of dipyridamole and aspirin (1:5 ratio) tested negative in the Ames test, in vivo chromosome aberration tests (in mice and hamsters), oral micronucleus tests (in mice and hamsters) and oral dominant lethal test (in mice). Aspirin, alone, induced chromosome aberrations in cultured human fibroblasts. Mutagenicity tests of dipyridamole alone with bacterial and mammalian cell systems were negative.

Combinations of dipyridamole and aspirin have not been evaluated for effects on fertility and reproductive performance. There was no evidence of impaired fertility when dipyridamole was administered to male and female rats at oral doses up to 500 mg/kg/day (about 12 times the MRHD on a mg/m^2 basis). A significant reduction in number of corpora lutea with consequent reduction in implantations and live fetuses was, however, observed at 1,250 mg/kg (more than 30 times the MRHD on a mg/m^2 basis). Aspirin inhibits ovulation in rats.

14 CLINICAL STUDIES

ESPS2 (European Stroke Prevention Study-2) was a double-blind, placebo-controlled, 24-month study in which 6602 patients over the age of 18 years had an ischemic stroke (76%) or transient ischemic attack (TIA, 24%) within three months prior to entry. Patients were enrolled in 13 European countries between February 1989 and May 1995 and were randomized to one of four treatment groups: Aspirin/extended-release dipyridamole 25 mg/200 mg; extended-release dipyridamole (ER-DP) 200 mg alone; aspirin (ASA) 25 mg alone; or placebo. The mean age in this population was 66.7 years with 58% of them being males. Patients received one capsule twice daily (morning and evening). Efficacy assessments included analyses of stroke (fatal or nonfatal) and death (from all causes) as confirmed by a blinded morbidity and mortality assessment group. There were no differences with regard to efficacy based on age or gender; patients who were older had a trend towards more events.

Stroke Endpoint

Aspirin and extended-release dipyridamole capsules reduced the risk of stroke by 22.1% compared to aspirin 50 mg/day alone ($p = 0.008$) and reduced the risk of stroke by 24.4% compared to extended-release dipyridamole 400 mg/day alone ($p = 0.002$) (Table 3). Aspirin and extended-release dipyridamole capsules reduced the risk of stroke by 36.8% compared to placebo ($p < 0.001$).

Table 3 Summary of First Stroke (Fatal or Nonfatal): ESPS2: Intent-to-Treat Population

	Total Number of Patients n	Number of Patients With Stroke Within 2 Years n(%)	Kaplan- Meier Estimate of Survival at 2 Years (95% C.I.)	Gehan- Wilcoxon Test P- value	Risk Reduction at 2 Years	Odds Ratio (95% C.I.)
Individual Treatment Group						
Aspirin and extended-release dipyridamole capsules	1650	157 (9.5%)	89.9% (88.4%, 91.4%)			
ER-DP ASA	1654	211 (12.8%)	86.7% (85.0%, 88.4%)			
Placebo	1649	206 (12.5%)	87.1% (85.4%, 88.7%)	----	----	----
	1649	250 (15.2%)	84.1% (82.2%, 85.9%)			
Pairwise Treatment Group Comparisons						
Aspirin and extended- release dipyridamole capsules ER-DP	vs. -	-	-	0.002 ^b	24.4%	0.72(0.58, 0.90)
Aspirin and extended- release dipyridamole capsules ASA	vs. -	-	-	0.008 ^b	22.1%	0.74(0.59, 0.92)
Aspirin and extended- release dipyridamole capsules Placebo	vs. -	-	-	<0.001 ^b	36.8%	0.59(0.48, 0.73)

ER-DP vs. Placebo	-	-	0.036 ^a	16.5%	0.82(0.67, 1.00)
ASA vs. Placebo	-	-	0.009 ^b	18.9%	0.80(0.66, 0.97)


^a 0.010 < p-value ≤ 0.050;

^b p-value ≤ 0.010.

Note: ER-DP = extended-release dipyridamole 200 mg; ASA = aspirin 25 mg. The dosage regimen for all treatment groups is BID.

Figure 1 ESPS2: Cumulative Stroke Rate (Fatal or Nonfatal)

Over 24 months of Follow-Up



Note: ER-DP = extended-release dipyridamole 200 mg; ASA = aspirin 25 mg. The dosage regimen for all treatment groups is b.i.d.

Combined Stroke or Death Endpoint

In ESPS2, aspirin and extended-release dipyridamole capsules reduced the risk of stroke or death by 24.2% compared to placebo.

Aspirin and extended-release dipyridamole capsules reduced the risk of stroke or death by 12.1% compared to aspirin alone and by 10.3% compared to extended-release dipyridamole alone. These results were not statistically significant.

Death Endpoint

The incidence rate of all-cause mortality was 11.3% for aspirin and extended-release dipyridamole capsules, 11% for aspirin alone, 11.4% for extended-release dipyridamole

alone and 12.3% for placebo alone. The differences between the aspirin and extended-release dipyridamole capsules, aspirin alone and extended-release dipyridamole alone treatment groups were not statistically significant. These incidence rates for aspirin and extended-release dipyridamole capsules and aspirin alone are consistent with previous aspirin studies in stroke and TIA patients.

16 HOW SUPPLIED/STORAGE AND HANDLING

Aspirin and extended-release dipyridamole capsules are available as a hard gelatin capsule, with orange colored cap and white to off white colored body, containing yellow to dark yellow colored extended-release pellets incorporating dipyridamole and white to off white round biconvex film coated tablet incorporating immediate-release aspirin. The capsule body is imprinted with '339' on the body and 'RDY' on cap with black ink.

Aspirin and extended-release dipyridamole capsules are supplied in unit-of-use bottles of 60s capsules.

Bottles of 60's NDC 16714-964-01

Store at 20°C to 25°C (68°F to 77°F); [See USP Controlled Room Temperature]. Protect from excessive moisture.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

• Risk of Bleeding

Inform patients that as with other antiplatelet agents, there is a general risk of bleeding including intracranial and gastrointestinal bleeding. Inform patients about the signs and symptoms of bleeding, including occult bleeding. Tell patients to notify their physician if they are prescribed any drug which may increase risk of bleeding.

Counsel patients who consume three or more alcoholic drinks daily about the bleeding risks involved with chronic, heavy alcohol use while taking aspirin.

• Pregnancy

Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during treatment with aspirin and extended-release dipyridamole capsules [see *Use in Specific Populations* (8.1)].

• Headaches

Some patients may experience headaches upon treatment initiation; these are usually transient. In case of intolerable headaches, tell patients to contact their physician.

• Stress Test

Instruct patients who are scheduled to undergo a pharmacologic stress test to tell their healthcare provider that they are taking aspirin and extended-release dipyridamole capsules.

• Dosage and Administration

Tell patients that aspirin and extended-release dipyridamole capsules should be

swallowed whole, and not chewed or crushed. If you miss a dose, continue with your next dose on your regular schedule. Do not take a double dose.

- **Storage**

Inform patients to protect aspirin and extended-release dipyridamole capsules from moisture.

Rx only

Manufactured for:

Northstar Rx LLC

Memphis, TN 38141

Manufactured by

Dr. Reddy's Laboratories Limited

Bachupally - 500 090 INDIA

Revised: 0521

PATIENT INFORMATION

Aspirin and Extended-Release Dipyridamole Capsules

(as' pir in and dye'' pir id' a mole)

Read this Patient Information before you start taking aspirin and extended-release dipyridamole capsules and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

What is aspirin and extended-release dipyridamole capsules?

Aspirin and extended-release dipyridamole capsule is a prescription medicine that contains aspirin and a medicine that is slowly released in your body, called dipyridamole. Aspirin and extended-release dipyridamole capsules are used to lower the risk of stroke in people who have had a “mini-stroke” (transient ischemic attack or TIA) or stroke due to a blood clot.

It is not known if aspirin and extended-release dipyridamole capsules is safe and effective in children. See “**Who should not take aspirin and extended-release dipyridamole capsules?**”

Who should not take aspirin and extended-release dipyridamole capsules?

Do not take aspirin and extended-release dipyridamole capsules if you:

- are allergic to any of the ingredients in aspirin and extended-release dipyridamole capsules. See the end of this leaflet for a list of ingredients in aspirin and extended-release dipyridamole capsules.
- are allergic to non-steroidal anti-inflammatory drugs (NSAIDs)
- have asthma in combination with runny nose and nasal polyps

Do not give aspirin and extended-release dipyridamole capsules to a child or

teenager with a viral illness. Reye syndrome, a life-threatening condition, can happen when aspirin (an ingredient in aspirin and extended-release dipyridamole capsules) is used in children and teenagers who have certain viral illnesses.

What should I tell my doctor before using aspirin and extended-release dipyridamole capsules?

Before taking aspirin and extended-release dipyridamole capsules, tell your healthcare provider if you:

- have stomach ulcers
- have a history of bleeding problems
- have heart problems
- have kidney or liver problems
- have low blood pressure
- have myasthenia gravis
- have any other medical conditions
- are pregnant or plan to become pregnant. You should not take aspirin and extended-release dipyridamole capsules during pregnancy without first talking to your healthcare provider. Tell your healthcare provider right away if you become pregnant while taking aspirin and extended-release dipyridamole capsules.
- are breast-feeding or plan to breast-feed. Aspirin and dipyridamole can pass into your milk. Talk to your healthcare provider about the best way to feed your baby if you take aspirin and extended-release dipyridamole capsules.

Tell your doctor you are taking aspirin and extended-release dipyridamole capsules if you are scheduled to have a stress test for your heart.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins and herbal supplements. Aspirin and extended-release dipyridamole capsules and other medicines may affect each other causing side effects. Aspirin and extended-release dipyridamole capsules may affect the way other medicines work, and other medicines may affect how aspirin and extended-release dipyridamole capsules works. Especially tell your healthcare provider if you take:

- **a medicine for high blood pressure, irregular heart beat, or heart failure**
- **acetazolamide [Diamox®]**
- **any blood thinner medicines**
- **warfarin sodium [Coumadin®, Jantoven®]**
- **a heparin medicine**
- **anagrelide [Agrylin®]**
- **a seizure medicine**
- **a medicine for Alzheimer's disease**

- a water pill
- methotrexate sodium [Trexall®]
- aspirin or a non-steroidal anti-inflammatory drug (NSAIDs). You should not take NSAIDs during treatment with aspirin and extended-release dipyridamole capsules. Using these medicines with aspirin and extended-release dipyridamole capsules can increase your risk of bleeding.
- a medicine for diabetes
- probenecid [Probalan®, Col-Probenecid®]

Ask your healthcare provider or pharmacist if you are not sure if your medicine is one that is listed above.

Know the medicines you take. Keep a list of them and show your healthcare provider and pharmacist when you get a new medicine.

How should I take aspirin and extended-release dipyridamole capsules?

- Take aspirin and extended-release dipyridamole capsules exactly as prescribed. Your healthcare provider will tell you how many aspirin and extended-release dipyridamole capsules to take and when to take them.
- Headaches are not uncommon when you first start taking aspirin and extended-release dipyridamole capsules, but often lessen as treatment continues. Tell your healthcare provider if you have a severe headache. Your healthcare provider may change the instructions for taking aspirin and extended-release dipyridamole capsules.
- Swallow aspirin and extended-release dipyridamole capsules whole. Do not crush or chew the capsules.
- You can take aspirin and extended-release dipyridamole capsules with or without food.
- **If you miss a dose, take your next dose at the usual time. Do not take two doses at one time.**
- If you take more aspirin and extended-release dipyridamole capsules (overdose) than prescribed, call your healthcare provider or Poison Control Center, or get emergency help right away.

Symptoms of an overdose of aspirin and extended-release dipyridamole capsules include:

- a warm feeling or flushing
- sweating
- restlessness
- weakness or dizziness
- a fast heart rate
- ringing in the ears

What should I avoid while using aspirin and extended-release dipyridamole capsules?

- **heavy alcohol use.** People who drink three or more alcoholic drinks every day have a higher risk of bleeding during treatment with aspirin and extended-release dipyridamole capsules, because it contains aspirin.

What are the possible side effects of aspirin and extended-release dipyridamole capsules?

Aspirin and extended-release dipyridamole capsules may cause serious side effects, including:

- **increased risk of bleeding.** You may bleed more easily during aspirin and extended-release dipyridamole capsules treatment, and it may take longer than usual for bleeding to stop. This can include:

- **bleeding into your brain (intracranial hemorrhage).** This can be a medical emergency. Get medical help right away if you have any of these symptoms while taking aspirin and extended-release dipyridamole capsules:

- severe headache with drowsiness
- confusion or memory change
- pass out (become unconscious)

- **bleeding in your stomach or intestine.**

- stomach pain
- heartburn or nausea
- vomiting blood or vomit looks like “coffee grounds”
- red or bloody stools
- black stools that look like tar

- **new or worsening chest pain in some people with heart disease.** Tell your healthcare provider if you have new chest pain or have any change in your chest pain during treatment with aspirin and extended-release dipyridamole capsules.

- **liver problems,** including increased liver function tests and liver failure. Tell your healthcare provider if you have any of these symptoms of a liver problem while taking aspirin and extended-release dipyridamole capsules:

- loss of appetite
- pale colored stool
- stomach area (abdomen) pain
- yellowing of your skin or whites of your eyes
- dark urine
- itching

Call your healthcare provider right away if you have any of the symptoms listed above.

The most common side effects of aspirin and extended-release dipyridamole capsules include:

- headache
- upset stomach
- diarrhea

These are not all the possible side effects of aspirin and extended-release dipyridamole capsules. Tell your healthcare provider or pharmacist if you have any side effect that bothers you or that does not go away.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA1088.

How should I store aspirin and extended-release dipyridamole capsules?

- Store aspirin and dipyridamole extended-release capsules at 20°C to 25°C (68°F to 77°F).
- Keep aspirin and dipyridamole extended-release capsules dry.

Keep aspirin and extended-release dipyridamole capsules and all medicines out of the reach of children.

General information about aspirin and extended-release dipyridamole capsules

Medicines are sometimes prescribed for purposes other than those listed in the Patient Information. Do not use aspirin and extended-release dipyridamole capsules for a condition for which it was not prescribed. Do not give aspirin and extended-release dipyridamole capsules to other people, even if they have the same symptoms that you have. It may harm them.

This Patient Information summarizes the most important information about aspirin and extended-release dipyridamole capsules. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about aspirin and extended-release dipyridamole capsules that is written for health professionals.

For more information, call 1-800-206-7821

What are the ingredients in aspirin and extended-release dipyridamole capsules?

Active Ingredients: dipyridamole in an extended-release form and aspirin

Inactive Ingredients: acacia, anhydrous lactose, colloidal silicon dioxide, dimethicone, hypromellose, hypromellose phthalate, lecithin, methacrylic acid copolymer, microcrystalline cellulose, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, tartaric acid, titanium dioxide, triacetin and xanthan gum.

Each capsule shell contains FD&C yellow 6, gelatin, sodium lauryl sulfate, titanium dioxide and yellow iron oxide.

Imprinting ink contains black iron oxide, potassium hydroxide, propylene glycol, shellac and strong ammonia solution.

The other brands listed are trademarks of their respective owners and are not trademarks of Dr. Reddy's Laboratories Inc.

Rx only

Manufactured for:

Northstar Rx LLC

Memphis, TN 38141

Manufactured by:

Dr. Reddy's Laboratories Limited


Bachupally - 500 090 INDIA

Revised: 0521

PACKAGE LABEL PRINCIPAL DISPLAY PANEL SECTION

Container: 60s count

Unvarnished Area Consists of 2D Barcode, Lot Number, Expiry Date and Serial Number



ASPIRIN AND DIPYRIDAMOLE

aspirin and dipyridamole capsule, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:16714-964
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Aspirin (UNII: R16CO5Y76E) (Aspirin - UNII:R16CO5Y76E)	Aspirin	25 mg
Dipyridamole (UNII: 64ALC7F90C) (Dipyridamole - UNII:64ALC7F90C)	Dipyridamole	200 mg

Inactive Ingredients

Ingredient Name	Strength
Acacia (UNII: 5C5403N26O)	
Anhydrous Lactose (UNII: 3SY5LH9PMK)	
Silicon Dioxide (UNII: ETJ7Z6XBU4)	
Dimethicone (UNII: 92RU3N3Y1O)	
Hypromelloses (UNII: 3NXW29V3WO)	
Hypromellose Phthalate (24% Phthalate, 55 Cst) (UNII: 87Y6436BKR)	
Lecithin, Soybean (UNII: 1DI56QDM62)	
Cellulose, Microcrystalline (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
Polyvinyl Alcohol, Unspecified (UNII: 532B59J990)	
Povidone (UNII: FZ989GH94E)	
Stearic Acid (UNII: 4ELV7Z65AP)	
Talc (UNII: 7SEV7J4R1U)	
Tartaric Acid (UNII: W4888I119H)	
Titanium Dioxide (UNII: 15FIX9V2JP)	
Triacetin (UNII: XHX3C3X673)	
Xanthan Gum (UNII: TTV12P4NEE)	
Gelatin (UNII: 2G86QN327L)	
Sodium Lauryl Sulfate (UNII: 368GB5141J)	
Fd&C Yellow No. 6 (UNII: H77VEI93A8)	
Ferric Oxide Yellow (UNII: EX438O2MRT)	
Methacrylic Acid - Methyl Methacrylate Copolymer (1:2) (UNII: 5KY68S2577)	
Ferrosoferric Oxide (UNII: XM0M87F357)	
Potassium Hydroxide (UNII: WZH3C48M4T)	
Propylene Glycol (UNII: 6DC9Q167V3)	
Shellac (UNII: 46N107B71O)	
Ammonia (UNII: 5138Q19F1X)	

Product Characteristics

Color	ORANGE (orange colored cap and white to off white colored body)	Score	no score
Shape	CAPSULE	Size	25mm
Flavor		Imprint Code	RDY;339
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16714-964-01	60 in 1 BOTTLE; Type 0: Not a Combination Product	04/03/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209048	04/03/2019	

Labeler - NorthStar Rx LLC (830546433)

Establishment

Name	Address	ID/FEI	Business Operations
Dr. Reddy's Laboratories Limited-FTO III		918608162	analysis(16714-964) , manufacture(16714-964)

Revised: 12/2023

NorthStar Rx LLC