

**TOPCARE CHILDRENS COLD COUGH AND SORE THROAT- acetaminophen,
dextromethorphan hydrobromide suspension**

Topco Associates LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topco Associates LLC. Children's Cold, Cough & Sore Throat Drug Facts

Active Ingredients (in each 5 mL)

Acetaminophen 160 mg

Dextromethorphan HBr 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Uses

- temporarily relieves the following cold/flu symptoms:
- minor aches and pains
- headache
- sore throat
- cough
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

- liver disease
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dosage (see overdose warning)**

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours, while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)
under 36	under 4 years	do not use
36-47	4 to 5 years	5 mL
48-95	6 to 11 years	10 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

Inactive Ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, edetate disodium, FD&C red #40, flavor, glycerin, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, tribasic sodium phosphate, xanthan gum

Questions or comments?

1-888-423-0139

Package/Label Principal Display Panel

COMPARE TO CHILDREN'S TYLENOL® COLD + COUGH + SORE THROAT ORAL
SUSPENSION ACTIVE INGREDIENTS

children's

Cold, Cough & Sore Throat

ORAL SUSPENSION

PAIN RELIEVER – FEVER REDUCER – ACETAMINOPHEN

COUGH SUPPRESSANT – DEXTROMETHORPHAN HBr

RELIEVES:

Fever

Sore Throat

Cough

OUR PHARMACISTS RECOMMEND

Ages 4 to 11 Years

BUBBLE GUM FLAVOR

4 FL OZ (118 mL)



TOPCARE CHILDRENS COLD COUGH AND SORE THROAT

acetaminophen, dextromethorphan hydrobromide suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-893
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCROSE (UNII: C151H8M554)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	PINK	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-893-26	1 in 1 CARTON	08/13/2019	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/13/2019	

Labeler - Topco Associates LLC (006935977)