

BISACODYL 5MG- bisacodyl 5mg tablet, coated
Allegiant Health

232 - Health A2Z Womans Laxative

Active Ingredient (in each tablet)

Bisacodyl USP 5mg

Purpose

Stimulant Laxative

Uses

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

- if you are unable to swallow without chewing
- within 1 hour after taking an antacid or milk

Ask a doctor before use if you have

- abdominal pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that persists over a period of 2 weeks

When using this product

- do not chew or crush tablets
- abdominal discomfort, faintness, and cramps may occur

Stop use and ask a doctor if

- you need to use more than 1 week
- rectal bleeding or failure to have a bowel movement after use of a laxative.

These may be signs of a serious condition.

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children

Keep out of reach of children. In case of accidental overdose, get medical help or

contact a doctor or Poison Control Center (1-800-222-1222) right away.

Directions

- take with water

Adults and children 12 years of age and older: take 1 to 3 tablets in a single dose, once daily

Children 6 to 12 years of age: take 1 tablet once a day

Children under 6 years of age: consult a doctor

Other information

- store between 68° to 77°F (20° to 25°C)
- do not use blister unit is broken or torn

Inactive ingredients

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C red #27 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow # 6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polyvinyl acetate phthalate, polydextrose, polyethylene glycol, propylparaben, propylene glycol, povidone, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

Principal Display Panel

HealthA2Z®

Woman's Gentle Laxative

HealthA2Z®

Compare to Dulcolax Pink® active ingredient*

Woman's Gentle Laxative



BISACODYL STIMULANT LAXATIVE 5 mg

For Gentle, Reliable Overnight Relief



25 Tablets

Lot:

Exp:

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark Dulcolax Pink®.

polyethylene glycol, propylparaben, propylene glycol, povidone, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, tracefin, triethyl citrate

Drug Facts (continued)

Inactive ingredients acacia, ammonium hydroxide, calcium carbonate, camellia wax, colloidal anhydrous silica, corn starch, D&C red #27 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polyvinyl acetate phthalate, polydextrose, ▲

Other information ■ store between 68° to 77°F (20° to 25°C) ■ do not use if blister unit is broken or torn

Directions ■ take with water
Adults and children 12 years of age and older: take 1 to 3 tablets in a single dose, once daily
Children 6 to 12 years of age: take 1 tablet once a day
Children under 6 years of age: consult a doctor

Keep out of reach of children. In case of accidental overdose, get medical help or contact a doctor or Poison Control Center (1-800-222-1222) right away.
If pregnant or breast-feeding, ask a health professional before use.

Stop use and ask a doctor if ■ you need to use more than 1 week ■ rectal bleeding or failure to have a bowel movement after use of a laxative. These may be signs of a serious condition.
When using this product ■ do not chew or crush tablets ■ abdominal discomfort, faintness, and cramps may occur

Ask a doctor before use if you have ■ abdominal pain ■ nausea ■ vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
Do not use ■ if you are unable to swallow without chewing ■ within 1 hour after taking an antacid or milk

Warnings

Uses ■ for relief of occasional constipation (irregularity) ■ this product generally produces bowel movement in 6 to 12 hours

Active ingredient (in each tablet) Bisacodyl USP 5mg.....Stimulant Laxative

Purpose

Distributed by:
Allegiant Health
Deer Park, NY 11729

LB1638
R1218



Bisacodyl 5mg Pink

BISACODYL 5MG

bisacodyl 5mg tablet, coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:69168-398 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y) | BISACODYL | 5 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ACACIA (UNII: 5C5403N26O) | |
| CALCIUM CARBONATE (UNII: H0G9379FGK) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| D&C RED NO. 27 (UNII: 2LRS185U6K) | |
| FD&C BLUE NO. 2--ALUMINUM LAKE (UNII: 4AQJ3LG584) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| METHYLPARABEN (UNII: A2I8C7HI9T) | |
| Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |
| POLYETHYLENE GLYCOL 1000 (UNII: U076Q6Q621) | |
| PROPYLPARABEN (UNII: Z8IX2SC1OH) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| POVIDONE (UNII: FZ989GH94E) | |
| SHELLAC (UNII: 46N107B71O) | |
| SODIUM ALGINATE (UNII: C269C4G2ZQ) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| SUCROSE (UNII: C151H8M554) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| TRIACETIN (UNII: XHX3C3X673) | |
| TRIETHYL CITRATE (UNII: 8Z96QXD6UM) | |

Product Characteristics

| | | | |
|---------------|-------|---------------------|----------|
| Color | pink | Score | no score |
| Shape | ROUND | Size | 5mm |
| Flavor | | Imprint Code | |

Contains**Packaging**

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:69168-398-92 | 1 in 1 CARTON | 12/23/2014 | |
| 1 | | 25 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |
| 2 | NDC:69168-398-02 | 150 in 1 BOTTLE; Type 0: Not a Combination Product | 01/16/2024 | |
| 3 | NDC:69168-398-03 | 250 in 1 BOTTLE; Type 0: Not a Combination Product | 03/07/2025 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M007 | 12/23/2014 | |

Labeler - Allegiant Health (079501930)

Revised: 3/2019

Allegiant Health