

**PROVON ANTIMICROBIAL FOAM HANDWASH- chlorhexidine gluconate  
2% solution  
Xttrium Laboratories, Inc.**

-----  
**2% GOJO ADX/LTX**

**Active ingredient**

chlorhexidine gluconate 2% solution

**Purpose**

Antiseptic

**Uses**

- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease
- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care

**Warnings**

**For external use only**

**Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- if you are allergic to chlorhexidine gluconate or any other ingredient in this product

**When using this product**

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- do not use routinely if you have wounds which involve more than the superficial

layers of the skin

**Stop use and ask a doctor if** irritation, sensitization or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

- Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

### **Healthcare personnel handwash:**

- wet hands with water
- dispense 5 mL of product (6 full strokes) into cupped hands and wash in a vigorous manner for 15 seconds and rinse under running water and dry thoroughly

### **Surgical hand scrub:**

- wet hands and forearms with water
- scrub for 3 minutes with 5 mL of product (6 full strokes), with a wet brush, paying close attention to the nails, cuticles, and skin between the fingers
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product (6 full strokes) and rinse under running water and dry thoroughly

### **Other information**

- store at 20-25° C (68-77° F)
- avoid excessive heat above 40° C (104° F)

### **Inactive ingredients**

citric add. cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

### **Questions or comments?**

call 1-800-321-9647 Monday through Friday 8:00 AM to 5:00 PM EST

PROVON® Antimicrobial Foam Handwash

Chlorhexidine Gluconate 2% Solution - Antiseptic

GOJOADXFLBLC

8842-643-F

**PROVON® Antimicrobial Foam Handwash**  
**Chlorhexidine Gluconate 2% Solution - Antiseptic**

**Drug Facts**

<b>Active ingredient</b>	<b>Purpose</b>
Chlorhexidine gluconate 2% solution .....	Antiseptic

**Uses** • **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease  
• **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care

**Warnings**

For external use only

**Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include:  
• wheezing/difficulty breathing • shock • facial swelling • hives • rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use** • if you are allergic to chlorhexidine gluconate or any other ingredient in this product

**When using this product**

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

**Stop use and ask a doctor** if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

**Healthcare personnel handwash:** • wet hands with water  
• dispense 5 mL of product (6 full strokes) into cupped hands and wash in a vigorous manner for 15 seconds  
• rinse and dry thoroughly

**Surgical hand scrub:** • wet hands and forearms with water  
• scrub for 3 minutes with 5 mL of product (6 full strokes), with a wet brush, paying close attention to the nails, cuticles, and skin between the fingers  
• a separate nail cleaner may be used • rinse thoroughly  
• wash for an additional 3 minutes with 5 mL of product (6 full strokes) and rinse under running water and dry thoroughly

**Other information** • store at 20-25°C (68-77°F)  
• avoid excessive heat above 40°C (104°F)

**Inactive ingredients** citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

GOJADXBLEB | 8842-643-F

8842

NDC 21749-850-90

Brought to you by GOJO

PROVON®

Antimicrobial Foam Handwash with 2% CHG

Chlorhexidine Gluconate 2% Solution - Antiseptic

42 fl oz (1250 mL)

GOJADXBLEB

8842-642-B

8842 NDC 21749-850-90

Brought to you by GOJO

**PROVON**<sup>®</sup>  
BRAND

Antimicrobial Foam  
Handwash with 2% CHG  
*Chlorhexidine Gluconate 2% Solution – Antiseptic*

0 73852 06917 4

GOJOADXBLLB 8842-642-B

42 fl oz (1250 mL)

Peel here for more Drug Facts

gojo

Distributed by:  
GOJO Industries, Inc.  
Akron, OH 44309  
800-321-9647 • 330-255-6000  
www.GOJO.com  
©2019, GOJO Industries, Inc.  
All rights reserved.  
Made in U.S.A.

**Drug Facts**

<b>Active ingredient</b>	<b>Purpose</b>
Chlorhexidine gluconate 2% solution	Antiseptic

**Uses**

- healthcare personnel handwash: helps reduce bacteria that potentially can cause disease
- surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care

**Warnings**

For external use only

**Allergy alert**

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- if you are allergic to chlorhexidine gluconate or any other ingredient in this product

**When using this product**

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

**Drug Facts** (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

**Healthcare personnel handwash:**

- wet hands with water
- dispense 5 mL of product (6 full strokes) into cupped hands and wash in a vigorous manner for 15 seconds and rinse under running water and dry thoroughly

**Surgical hand scrub:**

- wet hands and forearms with water
- scrub for 3 minutes with 5 mL of product (6 full strokes), with a wet brush, paying close attention to the nails, cuticles, and skin between the fingers
- a separate nail cleaner may be used • rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product (6 full strokes) and rinse under running water and dry thoroughly

**Other information** • store at 20-25°C (68-77°F)

- avoid excessive heat above 40°C (104°F)

**Inactive ingredients** citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

**Questions or comments?**

call 1-800-321-9647 Monday through Friday 8:00 AM to 5:00 PM EST

PROVON<sup>®</sup> Antimicrobial Foam Handwash

Chlorhexidine Gluconate 2% Solution - Antiseptic

GOJOLTXFLBLC

1922-643-F

**PROVON® Antimicrobial Foam Handwash**  
**Chlorhexidine Gluconate 2% Solution - Antiseptic**

**Drug Facts**

<b>Active ingredient</b>	<b>Purpose</b>
Chlorhexidine gluconate 2% solution	Antiseptic

**Uses** • healthcare personnel handwash: helps reduce bacteria that potentially can cause disease  
• surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care

**Warnings**

For external use only

**Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include:  
• wheezing/difficulty breathing • shock • facial swelling • hives • rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use** • if you are allergic to chlorhexidine gluconate or any other ingredient in this product

**When using this product**

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

**Stop use and ask a doctor** if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

**Healthcare personnel handwash:** • wet hands with water

- dispense 5 mL of product (6 full strokes) into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

**Surgical hand scrub:** • wet hands and forearms with water

- scrub for 3 minutes with 5 mL of product (6 full strokes), with a wet brush, paying close attention to the nails, cuticles, and skin between the fingers • a separate nail cleaner may be used • rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product (6 full strokes) and rinse under running water and dry thoroughly

**Other information** • store at 20-25°C (68-77°F)  
• avoid excessive heat above 40°C (104°F)

**Inactive ingredients** citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

GOJOLTXBLBLB; 1922-642-F

1922

NDC 21749-850-89

Brought to you by GOJO

PROVON®

Antimicrobial Foam Handwash with 2% CHG

Chlorhexidine Gluconate 2% Solution - Antiseptic

40.5 fl oz (1200 mL)

GOJOLTXBLBLB

1922-642-B

1922

NDC 21749-650-89

Peel here for more Drug Facts

Brought to you by GOJO



PROVON<sup>®</sup>  
BRAND

Distributed by:  
GOJO Incentives, Inc.  
Aston, OH 44209  
800-421-4647 • 330-255-6000  
www.GOJO.com  
©2019, GOJO Incentives, Inc.  
All rights reserved.  
Made in the USA with U.S. and  
foreign components

Antimicrobial Foam  
Handwash with 2% CHG  
Chlorhexidine Gluconate 2% Solution –  
Antiseptic



GOJOLTXLBLB 1922-642-B

40.5 fl oz (1200 mL)

**Drug Facts**

Active ingredient	Purpose
Chlorhexidine gluconate 2% solution	Antiseptic

**Uses**

- healthcare personnel handwash: helps reduce bacteria that potentially can cause disease
- surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care

**Warnings**  
For external use only

**Allergy alert**  
This product may cause a severe allergic reaction. Symptoms may include:  
• wheezing/difficulty breathing • shock • facial swelling • hives • rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**  
• if you are allergic to chlorhexidine gluconate or any other ingredient in this product

**When using this product**

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶



**Drug Facts (continued)**

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

**Healthcare personnel handwash:**

- wet hands with water
- dispense 5 mL of product (6 full strokes) into cupped hands and wash in a vigorous manner for 15 seconds and rinse under running water and dry thoroughly

**Surgical hand scrub:**

- wet hands and forearms with water
- scrub for 3 minutes with 5 mL of product (6 full strokes), with a wet brush, paying close attention to the nails, cuticles, and skin between the fingers
- a separate nail cleaner may be used • rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product (6 full strokes) and rinse under running water and dry thoroughly

**Other information** • store at 20-25°C (68-77°F)  
• avoid excessive heat above 40°C (104°F)

**Inactive ingredients** citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

**Questions or comments?**

call 1-800-321-9647 Monday through Friday 8:00 AM to 5:00 PM EST

**PROVON ANTIMICROBIAL FOAM HANDWASH**

chlorhexidine gluconate 2% solution

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0116-8500
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>CHLORHEXIDINE GLUCONATE</b> (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	20 mg in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>COCO DIETHANOLAMIDE</b> (UNII: 92005F972D)	
<b>HYDROXYETHYL CELLULOSE (2000 MPA.S AT 1%)</b> (UNII: S38J6RZN16)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>LAURAMINE OXIDE</b> (UNII: 4F6FC4MI8W)	
<b>WATER</b> (UNII: 059QF0KO0R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0116-8500-89	1200 mL in 1 BOTTLE, DISPENSING; Type 0: Not a Combination Product	07/01/2016	
2	NDC:0116-8500-90	1250 mL in 1 BOTTLE, DISPENSING; Type 0: Not a Combination Product	07/01/2016	



## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019422	07/01/2016	

**Labeler** - Xttrium Laboratories, Inc. (007470579)

**Registrant** - Xttrium Laboratories, Inc. (007470579)

## Establishment

Name	Address	ID/FEI	Business Operations
Xttrium Laboratories, Inc.		007470579	manufacture(0116-8500)

Revised: 12/2024

Xttrium Laboratories, Inc.