

C-LAX LAXATIVE- bisacodyl tablet, delayed release
CVS Pharmacy

CVS 44-676

Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

Purpose

Stimulant laxative

Uses

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness, and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- use by expiration date on package

Inactive ingredients

carmines, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal display panel

♥CVS
Health®

C-Lax
Laxative

BISACODYL USP, 5 mg
Stimulant laxative

Enteric coated
For gentle, dependable
constipation relief

Actual Size

250 TABLETS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

50844

REV0923D67613

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895


© 2019 CVS/pharmacy

CVS.com® 1-800-SHOP CVS

V-14236

✓ **CVS®** Quality


Money Back Guarantee



C-Lax Laxative

BISACODYL USP, 5 mg
Stimulant laxative

Enteric coated
For gentle, dependable
constipation relief




Actual Size
250 TABLETS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING


Drug Facts	Purpose Bisacodyl USP, 5 mg Stimulant laxative
Active ingredient (in each tablet) Bisacodyl USP, 5 mg	
Uses ■ for relief of occasional constipation (irregularity) ■ this product generally produces bowel movement in 6 to 12 hours	
Warnings Do not use if you cannot swallow without chewing. Ask a doctor before use if you have ■ a sudden change in bowel habits that lasts more than 2 weeks ■ stomach pain, nausea or vomiting	

No print/No varnish
Lot & Exp date



#965891
50844 REV0923D67613

Distributed by: **CVS Pharmacy, Inc.**
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-14236



PEEL HERE FOR MORE DRUG FACTS

STOP PEELING

Drug Facts (continued)	<p>When using this product</p> <ul style="list-style-type: none"> ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk ■ it may cause stomach discomfort, faintness, and cramps <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition. ■ you need to use a laxative for more than 1 week <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ■ do not take more than directed</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%;">take with a glass of water</td> <td style="width: 50%;">adults and children 12 years and over</td> </tr> <tr> <td></td> <td style="text-align: center;">take 1 to 3 tablets in a single daily dose</td> </tr> <tr> <td></td> <td style="text-align: center;">children 6 to under 12 years</td> </tr> <tr> <td></td> <td style="text-align: center;">take 1 tablet in a single daily dose</td> </tr> <tr> <td></td> <td style="text-align: center;">children under 6 years</td> </tr> <tr> <td></td> <td style="text-align: center;">ask a doctor</td> </tr> </table> <p>Other information</p> <ul style="list-style-type: none"> ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ avoid excessive humidity ■ use by expiration date on package <p>Inactive ingredients</p> <p>carminic acid, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shealac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate</p> <p>Questions or comments? 1-800-426-9391</p>	take with a glass of water	adults and children 12 years and over		take 1 to 3 tablets in a single daily dose		children 6 to under 12 years		take 1 tablet in a single daily dose		children under 6 years		ask a doctor
take with a glass of water	adults and children 12 years and over												
	take 1 to 3 tablets in a single daily dose												
	children 6 to under 12 years												
	take 1 tablet in a single daily dose												
	children under 6 years												
	ask a doctor												

CVS 44-676A

C-LAX LAXATIVE

bisacodyl tablet, delayed release

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:59779-676

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	B
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-676-52	6 in 1 CARTON	09/16/2015	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:59779-676-13	250 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/16/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	09/16/2015	

Labeler - CVS Pharmacy (062312574)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(59779-676)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(59779-676) , pack(59779-676)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(59779-676)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(59779-676)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(59779-676)

Revised: 1/2024

CVS Pharmacy