

# LEDIPASVIR AND SOFOSBUVIR- ledipasvir and sofosbuvir tablet, film coated

## Asegua Therapeutics LLC

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LEDIPASVIR and SOFOSBUVIR TABLETS safely and effectively. See full prescribing information for LEDIPASVIR and SOFOSBUVIR TABLETS.

LEDIPASVIR and SOFOSBUVIR tablets, for oral use, 90 mg/400 mg (Authorized generic of HARVONI®)  
Initial U.S. Approval: 2014

#### **WARNING: RISK OF HEPATITIS B VIRUS REACTIVATION IN PATIENTS COINFECTED WITH HCV AND HBV**

*See full prescribing information for complete boxed warning.*

**Hepatitis B virus (HBV) reactivation has been reported, in some cases resulting in fulminant hepatitis, hepatic failure, and death. (5.1)**

### INDICATIONS AND USAGE

Ledipasvir and sofosbuvir is a fixed-dose combination of ledipasvir, a hepatitis C virus (HCV) NS5A inhibitor, and sofosbuvir, an HCV nucleotide analog NS5B polymerase inhibitor, and is indicated for the treatment of chronic hepatitis C virus (HCV) in adults and pediatric patients 3 years of age and older:

- Genotype 1, 4, 5, or 6 infection without cirrhosis or with compensated cirrhosis
- Genotype 1 infection with decompensated cirrhosis, in combination with ribavirin
- Genotype 1 or 4 infection who are liver transplant recipients without cirrhosis or with compensated cirrhosis, in combination with ribavirin. (1)

### DOSAGE AND ADMINISTRATION

- Testing prior to the initiation of therapy: Test all patients for HBV infection by measuring HBsAg and anti-HBc. (2.1)
- Recommended treatment regimen and duration in patients 3 years of age and older: (2.2)

HCV Genotype	Patient Population	Regimen and Duration
Genotype 1	Treatment-naïve without cirrhosis or with compensated cirrhosis (Child-Pugh A)	Ledipasvir and sofosbuvir 12 weeks
	Treatment-experienced without cirrhosis	Ledipasvir and sofosbuvir 12 weeks
	Treatment-experienced with compensated cirrhosis (Child-Pugh A)	Ledipasvir and sofosbuvir 24 weeks
	Treatment-naïve and treatment-experienced with decompensated cirrhosis (Child-Pugh B or C)	Ledipasvir and sofosbuvir + ribavirin 12 weeks
Genotype 1 or 4	Treatment-naïve and treatment-experienced liver transplant recipients without cirrhosis, or with compensated cirrhosis (Child-Pugh A)	Ledipasvir and sofosbuvir + ribavirin 12 weeks
Genotype 4, 5, or 6	Treatment-naïve and treatment-experienced without cirrhosis or with compensated cirrhosis (Child-Pugh A)	Ledipasvir and sofosbuvir 12 weeks

- Recommended dosage in adults: One tablet (90 mg of ledipasvir and 400 mg of sofosbuvir) taken orally once daily with or without food. (2.3)
- Recommended dosage in pediatric patients 3 years and older: Recommended dosage of ledipasvir and sofosbuvir in pediatric patients 3 years of age and older is based on weight. Refer to Table 2 of the full prescribing information for specific dosing guidelines based on body weight. (2.4)
- Instructions for Use should be followed for preparation and administration of ledipasvir and sofosbuvir (HARVONI) oral pellets. (2.5)
- HCV/HIV-1 coinfection: For adult and pediatric patients with HCV/HIV-1 coinfection, follow the dosage recommendations in the tables in the full prescribing information. (2.3, 2.4)

- If used in combination with ribavirin, follow the recommendations for ribavirin dosing and dosage modifications. (2.3, 2.4)
- For patients with any degree of renal impairment, including end stage renal disease on dialysis, no ledipasvir and sofosbuvir dosage adjustment is recommended. (2.6)

#### ----- **DOSAGE FORMS AND STRENGTHS** -----

Tablets: 90 mg of ledipasvir and 400 mg of sofosbuvir. (3)

#### ----- **CONTRAINDICATIONS** -----

If used in combination with ribavirin, all contraindications to ribavirin also apply to ledipasvir and sofosbuvir combination therapy. (4)

#### ----- **WARNINGS AND PRECAUTIONS** -----

- Risk of Hepatitis B Virus Reactivation: Test all patients for evidence of current or prior HBV infection before initiation of HCV treatment. Monitor HCV/HBV coinfecting patients for HBV reactivation and hepatitis flare during HCV treatment and post-treatment follow-up. Initiate appropriate patient management for HBV infection as clinically indicated. (5.1)
- Bradycardia with amiodarone coadministration: Serious symptomatic bradycardia may occur in patients taking amiodarone, particularly in patients also receiving beta blockers, or those with underlying cardiac comorbidities and/or advanced liver disease. Coadministration of amiodarone with ledipasvir and sofosbuvir is not recommended. In patients without alternative, viable treatment options, cardiac monitoring is recommended. (5.2, 6.2, 7.2)

#### ----- **ADVERSE REACTIONS** -----

- The most common adverse reactions (incidence greater than or equal to 10%, all grades) observed with treatment with ledipasvir and sofosbuvir were fatigue, headache, and asthenia. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Asegua Therapeutics at 1-800-445-3235 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

#### ----- **DRUG INTERACTIONS** -----

- Coadministration with amiodarone may result in serious symptomatic bradycardia. Use of ledipasvir and sofosbuvir with amiodarone is not recommended. (5.2, 6.2, 7.2)
- P-gp inducers (e.g., rifampin, St. John's wort): May alter concentrations of ledipasvir and sofosbuvir. Use of ledipasvir and sofosbuvir with P-gp inducers is not recommended. (5.3, 7, 12.3)
- Consult the full prescribing information prior to use for potential drug interactions. (5.2, 5.3, 7, 12.3)
- Clearance of HCV infection with direct acting antivirals may lead to changes in hepatic function, which may impact safe and effective use of concomitant medications. Frequent monitoring of relevant laboratory parameters (INR or blood glucose) and dose adjustments of certain concomitant medications may be necessary. (7.2)

#### ----- **USE IN SPECIFIC POPULATIONS** -----

- Pediatric Use: No data are available regarding the safety of ledipasvir and sofosbuvir in pediatric patients with renal impairment. (8.4)

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 12/2024**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **WARNING: RISK OF HEPATITIS B VIRUS REACTIVATION IN PATIENTS COINFECTED WITH HCV AND HBV**

#### **1 INDICATIONS AND USAGE**

#### **2 DOSAGE AND ADMINISTRATION**

2.1 Testing Prior to the Initiation of Therapy

2.2 Recommended Treatment Regimen and Duration in Patients 3 Years of Age and Older with Genotype 1, 4, 5, or 6 HCV

2.3 Recommended Dosage in Adults

- 2.4 Recommended Dosage in Pediatric Patients 3 Years of Age and Older
- 2.5 Preparation and Administration of Ledipasvir and Sofosbuvir (HARVONI) Oral Pellets
- 2.6 Renal Impairment

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Risk of Hepatitis B Virus Reactivation in Patients Coinfected with HCV and HBV
- 5.2 Serious Symptomatic Bradycardia When Coadministered with Amiodarone
- 5.3 Risk of Reduced Therapeutic Effect Due to Use with P-gp Inducers
- 5.4 Risks Associated with Ribavirin Combination Treatment

### **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

### **7 DRUG INTERACTIONS**

- 7.1 Potential for Drug Interaction
- 7.2 Established and Potentially Significant Drug Interactions
- 7.3 Drugs without Clinically Significant Interactions with Ledipasvir and Sofosbuvir

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.4 Microbiology

### **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **14 CLINICAL STUDIES**

- 14.1 Description of Clinical Trials
- 14.2 Clinical Trials in Subjects with Genotype 1 HCV
- 14.3 Clinical Trials in Subjects with Genotype 4, 5, or 6 HCV
- 14.4 Clinical Trials in Subjects Coinfected with HCV and HIV-1
- 14.5 Clinical Trials in Liver Transplant Recipients and/or Subjects with Decompensated Cirrhosis
- 14.6 Clinical Trials in Adults with Severe Renal Impairment, Including those Requiring Dialysis
- 14.7 Clinical Trial in Pediatric Subjects

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### **WARNING: RISK OF HEPATITIS B VIRUS REACTIVATION IN PATIENTS COINFECTED WITH HCV AND HBV**

**Test all patients for evidence of current or prior hepatitis B virus (HBV) infection before initiating treatment with ledipasvir and sofosbuvir. HBV reactivation has been reported in HCV/HBV coinfecting patients who were undergoing or had completed treatment with HCV direct acting antivirals and were not receiving HBV antiviral therapy. Some cases have resulted in fulminant hepatitis, hepatic failure, and death. Monitor HCV/HBV coinfecting patients for hepatitis flare or HBV reactivation during HCV treatment and post-treatment follow-up. Initiate appropriate patient management for HBV infection as clinically indicated [see *Warnings and Precautions (5.1)*].**

## 1 INDICATIONS AND USAGE

Ledipasvir and sofosbuvir is indicated for the treatment of adults and pediatric patients 3 years of age and older with chronic hepatitis C virus (HCV) [see *Dosage and Administration (2.2 and 2.3) and Clinical Studies (14)*]:

- genotype 1, 4, 5, or 6 infection without cirrhosis or with compensated cirrhosis
- genotype 1 infection with decompensated cirrhosis, for use in combination with ribavirin
- genotype 1 or 4 infection who are liver transplant recipients without cirrhosis or with compensated cirrhosis, for use in combination with ribavirin

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Testing Prior to the Initiation of Therapy

Test all patients for evidence of current or prior HBV infection by measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) before initiating HCV treatment with ledipasvir and sofosbuvir [see *Warnings and Precautions (5.1)*].

### 2.2 Recommended Treatment Regimen and Duration in Patients 3 Years of Age and Older with Genotype 1, 4, 5, or 6 HCV

Table 1 shows the recommended ledipasvir and sofosbuvir treatment regimen and duration based on patient population. Relapse rates are affected by baseline host and viral factors and differ between treatment durations for certain subgroups [see *Clinical Studies (14)*].

For patients with HCV/HIV-1 coinfection, follow the dosage recommendations in Table 1 [see *Clinical Studies (14)*]. Refer to *Drug Interactions (7)* for dosage recommendations for concomitant HIV-1 antiviral drugs.

#### **Table 1 Recommended Treatment Regimen and Duration**

**for Ledipasvir and Sofosbuvir in Patients 3 Years of Age and Older with Genotype 1, 4, 5, or 6 HCV**

<b>HCV Genotype</b>	<b>Patient Population</b>	<b>Treatment Regimen and Duration</b>
Genotype 1	Treatment-naïve without cirrhosis or with compensated cirrhosis (Child-Pugh A)	Ledipasvir and sofosbuvir 12 weeks*
	Treatment-experienced <sup>†</sup> without cirrhosis	Ledipasvir and sofosbuvir 12 weeks
	Treatment-experienced <sup>†</sup> with compensated cirrhosis (Child-Pugh A)	Ledipasvir and sofosbuvir 24 weeks <sup>‡</sup>
	Treatment-naïve and treatment-experienced <sup>†</sup> with decompensated cirrhosis (Child-Pugh B or C)	Ledipasvir and sofosbuvir + ribavirin <sup>§</sup> 12 weeks
Genotype 1 or 4	Treatment-naïve and treatment-experienced <sup>†</sup> liver transplant recipients without cirrhosis, or with compensated cirrhosis (Child-Pugh A)	Ledipasvir and sofosbuvir + ribavirin <sup>§</sup> 12 weeks
Genotype 4, 5, or 6	Treatment-naïve and treatment-experienced <sup>†</sup> , without cirrhosis or with compensated cirrhosis (Child-Pugh A)	Ledipasvir and sofosbuvir 12 weeks

\* Ledipasvir and sofosbuvir for 8 weeks can be considered in treatment-naïve genotype 1 patients without cirrhosis who have pretreatment HCV RNA less than 6 million IU/mL [see *Clinical Studies (14.2)*].

† Treatment-experienced adult and pediatric subjects have failed a peginterferon alfa +/- ribavirin based regimen with or without an HCV protease inhibitor.

‡ Ledipasvir and sofosbuvir + ribavirin for 12 weeks can be considered in treatment-experienced genotype 1 patients with cirrhosis who are eligible for ribavirin [see *Dosage and Administration (2.3 and 2.4)* and *Clinical Studies (14.2)*].

§ See *Dosage and Administration 2.3 and 2.4* for ribavirin dosage recommendations.

### **2.3 Recommended Dosage in Adults**

The recommended dosage of ledipasvir and sofosbuvir in adults with genotype 1, 4, 5 or 6 HCV is one tablet (90 mg ledipasvir and 400 mg sofosbuvir) taken orally once daily with or without food [see *Clinical Pharmacology (12.3)*].

The daily dosage of ribavirin is weight-based (1000 mg for patients <75 kg and 1200 mg for those ≥75 kg) administered orally in two divided doses with food.

In patients with decompensated cirrhosis, the starting dosage of ribavirin is 600 mg and can be titrated up to 1000 mg for patients <75 kg and 1200 mg for those ≥75 kg in two divided doses with food. If the starting dosage of ribavirin is not well tolerated, the dosage should be reduced as clinically indicated based on hemoglobin levels.

For further information on ribavirin dosing and dosage modifications, refer to the ribavirin prescribing information [see *Dosage and Administration (2.4)*, *Use in Specific Populations (8.6)*, and *Clinical Studies (14.5)*].

## 2.4 Recommended Dosage in Pediatric Patients 3 Years of Age and Older

The recommended dosage of ledipasvir and sofosbuvir in pediatric patients 3 years of age and older with genotype 1, 4, 5, or 6 HCV is based on weight (Table 2). Table 3 provides the weight-based dosage of ribavirin when used in combination with ledipasvir and sofosbuvir for pediatric patients. Take ledipasvir and sofosbuvir (with or without food) once daily [see *Dosage and Administration (2.5)*, *Clinical Pharmacology (12.3)*, and *Clinical Studies (14.7)*]. Ledipasvir and sofosbuvir (HARVONI) oral pellets can be taken in pediatric patients who cannot swallow the tablet formulation.

**Table 2 Dosing for Pediatric Patients 3 Years and Older**

<b>Body Weight (kg)</b>	<b>Dosing of Ledipasvir and Sofosbuvir</b>	<b>Ledipasvir and Sofosbuvir Daily Dose</b>
at least 35	one Ledipasvir and Sofosbuvir 90 mg/400 mg tablet once daily or two Ledipasvir and Sofosbuvir (HARVONI) 45 mg/200 mg tablets once daily or two 45 mg/200 mg packets of Ledipasvir and Sofosbuvir (HARVONI) oral pellets once daily	90 mg/400 mg per day
17 to less than 35	one Ledipasvir and Sofosbuvir (HARVONI) 45 mg/200 mg tablet once daily or one 45 mg/200 mg packet of Ledipasvir and Sofosbuvir (HARVONI) oral pellets once daily	45 mg/200 mg per day
less than 17	one 33.75 mg/150 mg packet of Ledipasvir and Sofosbuvir (HARVONI) oral pellets once daily	33.75 mg/150 mg per day

**Table 3 Recommended Dosing for Ribavirin in Combination Therapy with Ledipasvir and Sofosbuvir for Pediatric Patients 3 Years and Older**

<b>Body Weight (kg)</b>	<b>Oral Ribavirin Daily Dosage*</b>
less than 47	15 mg per kg per day (divided dose AM and PM)
47-49	600 mg per day (1 × 200 mg AM, 2 × 200 mg PM)
50-65	800 mg per day (2 × 200 mg AM, 2 × 200 mg PM)
66-80	1000 mg per day (2 × 200 mg AM, 3 × 200 mg PM)
greater than 80	1200 mg per day (3 × 200 mg AM, 3 × 200 mg PM)

\* The daily dosage of ribavirin is weight-based and is administered orally in two divided doses with food.

## **2.5 Preparation and Administration of Ledipasvir and Sofosbuvir (HARVONI) Oral Pellets**

See the ledipasvir and sofosbuvir (HARVONI) oral pellets full Instructions for Use for details on the preparation and administration of ledipasvir and sofosbuvir (HARVONI) pellets.

Do not chew ledipasvir and sofosbuvir (HARVONI) oral pellets. If ledipasvir and sofosbuvir (HARVONI) oral pellets are administered with food, sprinkle the pellets on one or more spoonfuls of non-acidic soft food at or below room temperature. Examples of non-acidic foods include pudding, chocolate syrup, mashed potato, and ice cream. Take ledipasvir and sofosbuvir (HARVONI) oral pellets within 30 minutes of gently mixing with food and swallow the entire contents without chewing to avoid a bitter aftertaste.

## **2.6 Renal Impairment**

No dosage adjustment of ledipasvir and sofosbuvir tablets (90 mg/400 mg) is recommended in patients with any degree of renal impairment, including end stage renal disease (ESRD) on dialysis [see *Dosage and Administration (2.3)*]. Take ledipasvir and sofosbuvir tablets (90 mg/400 mg) with or without ribavirin according to the recommendations in Table 1 [see *Adverse Reactions (6.1)*, *Use in Specific Populations (8.6)* and *Clinical Studies (14.6)*]. Refer to ribavirin tablet prescribing information for ribavirin dosage modification for patients with CrCl less than or equal to 50 mL per minute.

## **3 DOSAGE FORMS AND STRENGTHS**

Ledipasvir and sofosbuvir tablets (90 mg/400 mg) are available as orange, diamond-shaped, film-coated tablets debossed with "ASE" on one side and "9875" on the other side of the tablet. Each tablet contains 90 mg ledipasvir and 400 mg sofosbuvir.

## **4 CONTRAINDICATIONS**

If ledipasvir and sofosbuvir is administered with ribavirin, the contraindications to ribavirin also apply to this combination regimen. Refer to the ribavirin prescribing information for a list of contraindications for ribavirin [see *Dosage and Administration* (2.2)].

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Risk of Hepatitis B Virus Reactivation in Patients Coinfected with HCV and HBV**

Hepatitis B virus (HBV) reactivation has been reported in HCV/HBV coinfecting patients who were undergoing or had completed treatment with HCV direct acting antivirals, and who were not receiving HBV antiviral therapy. Some cases have resulted in fulminant hepatitis, hepatic failure, and death. Cases have been reported in patients who are HBsAg positive and also in patients with serologic evidence of resolved HBV infection (i.e., HBsAg negative and anti-HBc positive). HBV reactivation has also been reported in patients receiving certain immunosuppressants or chemotherapeutic agents; the risk of HBV reactivation associated with treatment with HCV direct-acting antivirals may be increased in these patients.

HBV reactivation is characterized as an abrupt increase in HBV replication manifesting as a rapid increase in serum HBV DNA level. In patients with resolved HBV infection, reappearance of HBsAg can occur. Reactivation of HBV replication may be accompanied by hepatitis, i.e., increases in aminotransferase levels and, in severe cases, increases in bilirubin levels, liver failure, and death can occur.

Test all patients for evidence of current or prior HBV infection by measuring HBsAg and anti-HBc before initiating HCV treatment with ledipasvir and sofosbuvir. In patients with serologic evidence of HBV infection, monitor for clinical and laboratory signs of hepatitis flare or HBV reactivation during HCV treatment with ledipasvir and sofosbuvir and during post-treatment follow-up. Initiate appropriate patient management for HBV infection as clinically indicated.

### **5.2 Serious Symptomatic Bradycardia When Coadministered with Amiodarone**

Postmarketing cases of symptomatic bradycardia, as well as fatal cardiac arrest and cases requiring pacemaker intervention, have been reported when amiodarone is coadministered with ledipasvir and sofosbuvir. Bradycardia has generally occurred within hours to days, but cases have been observed up to 2 weeks after initiating HCV treatment. Patients also taking beta blockers, or those with underlying cardiac comorbidities and/or advanced liver disease, may be at increased risk for symptomatic bradycardia with coadministration of amiodarone. Bradycardia generally resolved after discontinuation of HCV treatment. The mechanism for this effect is unknown.

Coadministration of amiodarone with ledipasvir and sofosbuvir is not recommended. For patients taking amiodarone who have no other alternative, viable treatment options and who will be coadministered ledipasvir and sofosbuvir:

- Counsel patients about the risk of serious symptomatic bradycardia
- Cardiac monitoring in an in-patient setting for the first 48 hours of coadministration is recommended, after which outpatient or self-monitoring of the heart rate should occur on a daily basis through at least the first 2 weeks of treatment.

Patients who are taking ledipasvir and sofosbuvir who need to start amiodarone therapy due to no other alternative, viable treatment options should undergo similar cardiac monitoring as outlined above.

Due to amiodarone's long half-life, patients discontinuing amiodarone just prior to starting ledipasvir and sofosbuvir should also undergo similar cardiac monitoring as outlined above.

Patients who develop signs or symptoms of bradycardia should seek medical evaluation immediately. Symptoms may include near-fainting or fainting, dizziness or lightheadedness, malaise, weakness, excessive tiredness, shortness of breath, chest pains, confusion, or memory problems [see *Adverse Reactions (6.2)*, *Drug Interactions (7.2)*].

### **5.3 Risk of Reduced Therapeutic Effect Due to Use with P-gp Inducers**

The concomitant use of ledipasvir and sofosbuvir and P-gp inducers may significantly decrease ledipasvir and sofosbuvir plasma concentrations and may lead to a reduced therapeutic effect of ledipasvir and sofosbuvir. Therefore, the use of ledipasvir and sofosbuvir with P-gp inducers (e.g., rifampin, St. John's wort) is not recommended [see *Drug Interactions (7.2)*].

### **5.4 Risks Associated with Ribavirin Combination Treatment**

If ledipasvir and sofosbuvir is administered with ribavirin, the warnings and precautions for ribavirin, in particular the pregnancy avoidance warning, apply to this combination regimen. Refer to the ribavirin prescribing information for a full list of the warnings and precautions for ribavirin [see *Dosage and Administration (2.2)*].

## **6 ADVERSE REACTIONS**

The following serious adverse reactions are described below and elsewhere in labeling:

- Serious Symptomatic Bradycardia When Coadministered with Amiodarone [see *Warnings and Precautions (5.2)*].

### **6.1 Clinical Trials Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

If ledipasvir and sofosbuvir is administered with ribavirin to adults, refer to the prescribing information for ribavirin for a description of ribavirin-associated adverse reactions.

#### Clinical Trials in Adult Subjects

The safety assessment of ledipasvir and sofosbuvir was based on pooled data from three randomized, open-label Phase 3 clinical trials (ION-3, ION-1, and ION-2) of subjects with genotype 1 HCV with compensated liver disease (with and without cirrhosis) including 215, 539, and 326 subjects who received ledipasvir and sofosbuvir tablets (90 mg/400 mg) once daily by mouth for 8, 12, and 24 weeks, respectively [see *Clinical Studies (14)*].

The proportion of subjects who permanently discontinued treatment due to adverse

events was 0%, less than 1%, and 1% for subjects receiving ledipasvir and sofosbuvir for 8, 12, and 24 weeks, respectively.

The most common adverse reactions (at least 10%) were fatigue and headache in subjects treated with 8, 12, or 24 weeks of ledipasvir and sofosbuvir.

Table 4 lists adverse reactions (adverse events assessed as causally related by the investigator, all grades) observed in at least 5% of subjects receiving 8, 12, or 24 weeks of treatment with ledipasvir and sofosbuvir in clinical trials. The majority of adverse reactions presented in Table 4 occurred at severity of grade 1. The side-by-side tabulation is to simplify presentation; direct comparison across trials should not be made due to differing trial designs.

**Table 4 Adverse Reactions (All Grades) Reported in  $\geq$ 5% of Subjects Receiving 8, 12, or 24 Weeks of Treatment with Ledipasvir and Sofosbuvir**

	<b>Ledipasvir and Sofosbuvir 8 weeks (N=215)</b>	<b>Ledipasvir and Sofosbuvir 12 weeks (N=539)</b>	<b>Ledipasvir and Sofosbuvir 24 weeks (N=326)</b>
Fatigue	16%	13%	18%
Headache	11%	14%	17%
Nausea	6%	7%	9%
Diarrhea	4%	3%	7%
Insomnia	3%	5%	6%

The safety assessment of ledipasvir and sofosbuvir was also based on pooled data from three open-label trials (Study 1119, ION-4, and ELECTRON-2) in 118 subjects with chronic HCV genotype 4, 5, or 6 infection with compensated liver disease (with or without cirrhosis) [see *Clinical Studies (14.3)*]. The subjects received ledipasvir and sofosbuvir tablets (90 mg/400 mg) once daily by mouth for 12 weeks. The safety profile in subjects with chronic HCV genotype 4, 5, or 6 infection with compensated liver disease was similar to that observed in subjects with chronic HCV genotype 1 infection with compensated liver disease. The most common adverse reactions occurring in at least 10% of subjects were asthenia (18%), headache (14%), and fatigue (10%).

#### Adverse Reactions in Subjects with Cirrhosis

The safety assessment of ledipasvir and sofosbuvir with or without ribavirin was based on a randomized, double-blind and placebo-controlled trial in treatment-experienced genotype 1 subjects with compensated cirrhosis and was compared to placebo in the SIRIUS trial. Subjects were randomized to receive 24 weeks of ledipasvir and sofosbuvir tablets (90 mg/400 mg) once daily by mouth without ribavirin or 12 weeks of placebo followed by 12 weeks of ledipasvir and sofosbuvir tablets (90 mg/400 mg) once daily by mouth + ribavirin [see *Clinical Studies (14.2)*]. Table 5 presents the adverse reactions, as defined above, that occurred with at least 5% greater frequency in subjects treated with 24 weeks of ledipasvir and sofosbuvir or 12 weeks of ledipasvir and sofosbuvir + ribavirin, compared to those reported for 12 weeks of placebo. The majority of the adverse reactions presented in Table 5 were Grade 1 or 2 in severity.

**Table 5 Adverse Reactions with  $\geq 5\%$  Greater Frequency Reported in Treatment-Experienced Subjects with Cirrhosis Receiving Ledipasvir and Sofosbuvir for 24 Weeks or Ledipasvir and Sofosbuvir + Ribavirin for 12 Weeks Compared to Placebo for 12 weeks**

	<b>Ledipasvir and Sofosbuvir 24 weeks (N=78)</b>	<b>Ledipasvir and Sofosbuvir + RBV 12 weeks (N=76)</b>	<b>Placebo 12 weeks (N=77)</b>
Asthenia	31%	36%	23%
Headache	29%	13%	16%
Fatigue	18%	4%	1%
Cough	5%	11%	1%
Myalgia	9%	4%	0
Dyspnea	3%	9%	1%
Irritability	8%	7%	1%
Dizziness	5%	1%	0

RBV=ribavirin

#### Adverse Reactions in Subjects Coinfected with HIV-1

The safety assessment of ledipasvir and sofosbuvir was based on an open-label clinical trial in 335 genotype 1 or 4 subjects with HCV/HIV-1 coinfection who were on stable antiretroviral therapy in Study ION-4 [see *Clinical Studies (14.4)*]. The safety profile in HCV/HIV-1 coinfecting subjects was similar to that observed in HCV mono-infected subjects. The most common adverse reactions occurring in at least 10% of subjects were headache (20%) and fatigue (17%).

#### Adverse Reactions in Liver Transplant Recipients and/or Subjects with Decompensated Cirrhosis

The safety assessment of ledipasvir and sofosbuvir with ribavirin in liver transplant recipients and/or those who had decompensated liver disease was based on pooled data from two Phase 2 open-label clinical trials including 336 subjects who received ledipasvir and sofosbuvir tablets (90 mg/400 mg) plus ribavirin for 12 weeks. Subjects with Child-Pugh-Turcotte (CPT) scores greater than 12 were excluded from the trials [see *Clinical Studies (14.5)*].

The adverse events observed were consistent with the expected clinical sequelae of liver transplantation and/or decompensated liver disease, or the known safety profile of ledipasvir and sofosbuvir and/or ribavirin.

Decreases in hemoglobin to less than 10 g/dL and 8.5 g/dL during treatment were observed in 38% and 13% of subjects treated with ledipasvir and sofosbuvir plus ribavirin for 12 weeks, respectively. Ribavirin was permanently discontinued in 11% of subjects treated with ledipasvir and sofosbuvir plus ribavirin for 12 weeks.

#### *Liver Transplant Recipients with Compensated Liver Disease:*

Among the 174 liver transplant recipients with compensated liver disease who received ledipasvir and sofosbuvir tablets (90 mg/400 mg) with ribavirin for 12 weeks, 2 (1%)

subjects permanently discontinued ledipasvir and sofosbuvir due to an adverse event.

#### *Subjects with Decompensated Liver Disease:*

Among the 162 subjects with decompensated liver disease (pre- or post-transplant) who received ledipasvir and sofosbuvir tablets (90 mg/400 mg) with ribavirin for 12 weeks, 7 (4%) subjects died, 4 (2%) subjects underwent liver transplantation, and 1 subject (<1%) underwent liver transplantation and died during treatment or within 30 days after discontinuation of treatment. Because these events occurred in patients with advanced liver disease who are at risk of progression of liver disease including liver failure and death, it is not possible to reliably assess the contribution of drug effect to outcomes. A total of 4 (2%) subjects permanently discontinued ledipasvir and sofosbuvir due to an adverse event.

Less Common Adverse Reactions Reported in Clinical Trials (less than 5%): The following adverse reactions occurred in less than 5% of subjects receiving ledipasvir and sofosbuvir tablets (90 mg/400 mg) in any one trial. These events have been included because of their seriousness or assessment of potential causal relationship.

*Psychiatric disorders:* depression (including in subjects with pre-existing history of psychiatric illness).

Depression (particularly in subjects with pre-existing history of psychiatric illness) occurred in subjects receiving sofosbuvir containing regimens. Suicidal ideation and suicide have occurred in less than 1% of subjects treated with sofosbuvir in combination with ribavirin or pegylated interferon/ribavirin in other clinical trials.

#### Laboratory Abnormalities

*Bilirubin Elevations:* Bilirubin elevations of greater than 1.5×ULN were observed in 3%, less than 1%, and 2% of subjects treated with ledipasvir and sofosbuvir tablets (90 mg/400 mg) for 8, 12, and 24 weeks, respectively. Bilirubin elevations of greater than 1.5×ULN were observed in 3%, 11%, and 3% of subjects with compensated cirrhosis treated with placebo, ledipasvir and sofosbuvir + ribavirin for 12 weeks, and ledipasvir and sofosbuvir for 24 weeks, respectively, in the SIRIUS trial.

*Lipase Elevations:* Transient, asymptomatic lipase elevations of greater than 3×ULN were observed in less than 1%, 2%, and 3% of subjects treated with ledipasvir and sofosbuvir tablets (90 mg/400 mg) for 8, 12, and 24 weeks, respectively. Transient, asymptomatic lipase elevations of greater than 3×ULN were observed in 1%, 3%, and 9% of subjects with compensated cirrhosis treated with placebo, ledipasvir and sofosbuvir + ribavirin for 12 weeks, and ledipasvir and sofosbuvir for 24 weeks, respectively, in the SIRIUS trial.

*Creatine Kinase:* Creatine kinase was not assessed in Phase 3 trials ION-3, ION-1, or ION-2 of ledipasvir and sofosbuvir. Creatine kinase was assessed in the ION-4 trial. Isolated, asymptomatic creatine kinase elevations of greater than or equal to 10×ULN was observed in 1% of subjects treated with ledipasvir and sofosbuvir tablets (90 mg/400 mg) for 12 weeks in the ION-4 trial and has also been previously reported in subjects treated with sofosbuvir in combination with ribavirin or peginterferon/ribavirin in other clinical trials.

#### Adverse Reactions in Adults with Severe Renal Impairment, Including those on Dialysis

In an open-label trial (Trial 0154) in which adults with HCV with compensated liver disease (with or without cirrhosis) and severe renal impairment received ledipasvir and

sofosbuvir tablets (90 mg/400 mg) for 12 weeks (N=18), the most common adverse reaction was fatigue (17%) [see *Clinical Studies (14.6)*].

In an open-label clinical trial, Trial 4063, a total of 95 adults with HCV with compensated liver disease (with or without cirrhosis) and ESRD requiring dialysis received ledipasvir and sofosbuvir tablets (90 mg/400 mg) for 8 (n=45), 12 (n=31), or 24 (n=19) weeks. The most common adverse reactions were insomnia and headache (each reported in 4% of subjects overall) [see *Clinical Studies (14.6)*].

#### Adverse Reactions in Pediatric Subjects 3 Years of Age and Older

The safety assessment of ledipasvir and sofosbuvir tablets (90 mg/400 mg), ledipasvir and sofosbuvir (HARVONI) tablets (45 mg/200 mg), or ledipasvir and sofosbuvir (HARVONI) oral pellets in pediatric subjects 3 years of age and older is based on data from a Phase 2, open-label clinical trial (Study 1116). In total, 226 subjects were enrolled, which included 223 subjects without cirrhosis or with compensated cirrhosis who were treated with ledipasvir and sofosbuvir for 12 weeks; one genotype 1 treatment-experienced subject with cirrhosis who was treated with ledipasvir and sofosbuvir for 24 weeks; and two genotype 3 subjects who were treated with ledipasvir and sofosbuvir + ribavirin for 24 weeks. The adverse reactions observed were consistent with those observed in clinical studies of ledipasvir and sofosbuvir tablets (90 mg/400 mg) in adults. Limited safety data are available in pediatric subjects receiving ledipasvir and sofosbuvir for 24 weeks. No Grade 3 or 4 adverse reactions or discontinuation due to an adverse reaction was observed in those pediatric subjects receiving ledipasvir and sofosbuvir for 24 weeks [see *Clinical Studies (14.7)*].

In a 5-year follow-up study, 178 of the 226 subjects from the Phase 2 open-label clinical trial (Study 1116) were followed for a median (Q1, Q3) duration of 239 (143, 244) weeks. No notable effects on growth as assessed by changes from baseline through end of study were observed for height, weight, BMI percentiles, and Z-scores for any age group. No notable effects were observed on the development of secondary sexual characteristics of subjects as assessed by changes from baseline through end of study in Tanner pubertal stages [see *Use in Specific Populations (8.4)*].

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post approval use of ledipasvir and sofosbuvir. Because postmarketing reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

### Cardiac Disorders

Serious symptomatic bradycardia has been reported in patients taking amiodarone who initiate treatment with ledipasvir and sofosbuvir [see *Warnings and Precautions (5.2)*, *Drug Interactions (7.2)*].

### Skin and Subcutaneous Tissue Disorders

Skin rashes, sometimes with blisters or angioedema-like swelling  
Angioedema

## **7 DRUG INTERACTIONS**

## 7.1 Potential for Drug Interaction

Any interactions that have been identified with ledipasvir or sofosbuvir individually may occur with ledipasvir and sofosbuvir.

After oral administration of ledipasvir and sofosbuvir, sofosbuvir is rapidly absorbed and subject to extensive first-pass hepatic extraction. In clinical pharmacology studies, both sofosbuvir and the inactive metabolite GS-331007 were monitored for purposes of pharmacokinetic analyses.

Ledipasvir is an inhibitor of the drug transporters P-gp and breast cancer resistance protein (BCRP) and may increase intestinal absorption of coadministered substrates for these transporters.

Ledipasvir and sofosbuvir are substrates of drug transporters P-gp and BCRP while GS-331007 is not. P-gp inducers (e.g., rifampin, St. John's wort) may decrease ledipasvir and sofosbuvir plasma concentrations, leading to reduced therapeutic effect of ledipasvir and sofosbuvir, and the use with P-gp inducers is not recommended with ledipasvir and sofosbuvir [see *Warnings and Precautions (5.3)*].

## 7.2 Established and Potentially Significant Drug Interactions

Clearance of HCV infection with direct acting antivirals may lead to changes in hepatic function, which may impact the safe and effective use of concomitant medications. For example, altered blood glucose control resulting in serious symptomatic hypoglycemia has been reported in diabetic patients in postmarketing case reports and published epidemiological studies. Management of hypoglycemia in these cases required either discontinuation or dose modification of concomitant medications used for diabetes treatment.

Frequent monitoring of relevant laboratory parameters (e.g., International Normalized Ratio [INR] in patients taking warfarin, blood glucose levels in diabetic patients) or drug concentrations of concomitant medications such as cytochrome P450 substrates with a narrow therapeutic index (e.g., certain immunosuppressants) is recommended to ensure safe and effective use. Dose adjustments of concomitant medications may be necessary.

Table 6 provides a listing of established or potentially clinically significant drug interactions. The drug interactions described are based on studies conducted with either ledipasvir and sofosbuvir tablets (90 mg/400 mg) or ledipasvir and sofosbuvir as individual agents, or are predicted drug interactions that may occur with ledipasvir and sofosbuvir [see *Warnings and Precautions (5.2, 5.3)* and *Clinical Pharmacology (12.3)*].

**Table 6 Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interaction\***

<b>Concomitant Drug Class: Drug Name</b>	<b>Effect on Concentration†</b>	<b>Clinical Comment</b>
<b>Acid Reducing Agents:</b>		Ledipasvir solubility decreases as pH increases. Drugs that increase gastric pH are expected to decrease concentration of ledipasvir.
Antacids (e.g., aluminum		It is recommended to separate antacid

and magnesium hydroxide)		and ledipasvir and sofosbuvir administration by 4 hours.
H <sub>2</sub> -receptor antagonists <sup>‡</sup> (e.g., famotidine)	↓ ledipasvir	H <sub>2</sub> -receptor antagonists may be administered simultaneously with or 12 hours apart from ledipasvir and sofosbuvir at a dose that does not exceed doses comparable to famotidine 40 mg twice daily.
Proton-pump inhibitors <sup>‡</sup> (e.g., omeprazole)		Proton-pump inhibitor doses comparable to omeprazole 20 mg or lower can be administered simultaneously with ledipasvir and sofosbuvir under fasted conditions.
<b>Antiarrhythmics:</b> amiodarone	Effect on amiodarone, ledipasvir, and sofosbuvir concentrations unknown	Coadministration of amiodarone with ledipasvir and sofosbuvir may result in serious symptomatic bradycardia. The mechanism of this effect is unknown. Coadministration of amiodarone with ledipasvir and sofosbuvir is not recommended; if coadministration is required, cardiac monitoring is recommended [ <i>see Warnings and Precautions (5.2), Adverse Reactions (6.2)</i> ].
digoxin	↑ digoxin	Coadministration of ledipasvir and sofosbuvir with digoxin may increase the concentration of digoxin. Therapeutic concentration monitoring of digoxin is recommended when coadministered with ledipasvir and sofosbuvir.
<b>Anticonvulsants:</b> carbamazepine <sup>‡</sup> phenytoin phenobarbital	↓ ledipasvir ↓ sofosbuvir	Coadministration of ledipasvir and sofosbuvir with carbamazepine, phenytoin, or phenobarbital is expected to decrease the concentration of ledipasvir and sofosbuvir, leading to reduced therapeutic effect of ledipasvir and sofosbuvir. Coadministration is not recommended.
<b>Antimycobacterials:</b> rifabutin <sup>‡</sup> rifampin <sup>‡</sup> rifapentine	↓ ledipasvir ↓ sofosbuvir	Coadministration of ledipasvir and sofosbuvir with rifampin, rifabutin, or rifapentine is not recommended [ <i>see Warnings and Precautions (5.3)</i> ].
<b>HIV Antiretrovirals:</b>		
Regimens containing tenofovir DF without an HIV protease inhibitor/ritonavir or cobicistat	↑ tenofovir	Monitor for tenofovir-associated adverse reactions in patients receiving ledipasvir and sofosbuvir concomitantly with a regimen containing tenofovir DF without an HIV protease inhibitor/ritonavir or cobicistat. Refer to Viread <sup>®</sup> or Truvada <sup>®</sup> prescribing information for

		recommendations on renal monitoring.
<p>Regimens containing tenofovir DF and an HIV protease inhibitor/ritonavir or cobicistat</p> <ul style="list-style-type: none"> <li>• atazanavir/ritonavir or cobicistat + emtricitabine/tenofovir DF<sup>‡</sup></li> <li>• darunavir/ritonavir or cobicistat + emtricitabine/tenofovir DF<sup>‡</sup></li> <li>• lopinavir/ritonavir + emtricitabine/tenofovir DF</li> </ul>	↑ tenofovir	<p>The safety of increased tenofovir concentrations in the setting of ledipasvir and sofosbuvir and an HIV protease inhibitor/ritonavir or cobicistat has not been established. Consider alternative HCV or antiretroviral therapy to avoid increases in tenofovir exposures. If coadministration is necessary, monitor for tenofovir-associated adverse reactions. Refer to Viread or Truvada prescribing information for recommendations on renal monitoring.</p>
elvitegravir, cobicistat, emtricitabine, tenofovir DF	↑ tenofovir	The safety of increased tenofovir concentrations in the setting of ledipasvir and sofosbuvir and the combination of elvitegravir, cobicistat, emtricitabine, and tenofovir DF has not been established. Coadministration is not recommended.
tipranavir/ritonavir	↓ ledipasvir ↓ sofosbuvir	Coadministration of ledipasvir and sofosbuvir with tipranavir/ritonavir is expected to decrease the concentration of ledipasvir and sofosbuvir, leading to reduced therapeutic effect of ledipasvir and sofosbuvir. Coadministration is not recommended.
<b>HCV Products:</b> simeprevir <sup>‡</sup>	↑ ledipasvir ↑ simeprevir	Concentrations of ledipasvir and simeprevir are increased when simeprevir is coadministered with ledipasvir. Coadministration of ledipasvir and sofosbuvir with simeprevir is not recommended.
<b>Herbal Supplements:</b> St. John's wort ( <i>Hypericum perforatum</i> )	↓ ledipasvir ↓ sofosbuvir	Coadministration of ledipasvir and sofosbuvir with St. John's wort, a P-gp inducer, is not recommended [see <i>Warnings and Precautions (5.3)</i> ].
<b>HMG-CoA Reductase Inhibitors:</b> rosuvastatin	↑ rosuvastatin	Coadministration of ledipasvir and sofosbuvir with rosuvastatin may significantly increase the concentration of rosuvastatin, which is associated with increased risk of myopathy, including rhabdomyolysis. Coadministration of ledipasvir and sofosbuvir with rosuvastatin is not recommended.

atorvastatin	↑ atorvastatin	Coadministration of ledipasvir and sofosbuvir with atorvastatin may be associated with increased risk of myopathy, including rhabdomyolysis. Monitor closely for HMG-CoA reductase inhibitor-associated adverse reactions, such as myopathy and rhabdomyolysis.
--------------	----------------	---

tenofovir DF = tenofovir disoproxil fumarate

\* This table is not all inclusive.

† ↓ = decrease, ↑ = increase

‡ These interactions have been studied in healthy adults.

### 7.3 Drugs without Clinically Significant Interactions with Ledipasvir and Sofosbuvir

Based on drug interaction studies conducted with ledipasvir, sofosbuvir, or ledipasvir and sofosbuvir tablets (90 mg/400 mg) no clinically significant drug interactions have been either observed or are expected when ledipasvir and sofosbuvir is used with the following drugs [see *Clinical Pharmacology (12.3)*]: abacavir, atazanavir/ritonavir, cyclosporine, darunavir/ritonavir, dolutegravir, efavirenz, elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide, emtricitabine, lamivudine, methadone, midazolam, oral contraceptives, pravastatin, raltegravir, rilpivirine, tacrolimus, or verapamil. See Table 6 for use of ledipasvir and sofosbuvir with certain HIV antiretroviral regimens [see *Drug Interactions (7.2)*].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

If ledipasvir and sofosbuvir is administered with ribavirin, the combination regimen is contraindicated in pregnant women and in men whose female partners are pregnant. Refer to the ribavirin prescribing information for more information on ribavirin-associated risks of use during pregnancy.

No adequate human data are available to establish whether or not ledipasvir and sofosbuvir pose a risk to pregnancy outcomes. In animal reproduction studies, no evidence of adverse developmental outcomes was observed with ledipasvir or sofosbuvir at exposures greater than those in humans at the recommended human dose (RHD) [see *Data*]. During organogenesis in the rat and rabbit, systemic exposures (AUC) to ledipasvir were approximately 4 (rats) and 2 (rabbits) times the exposure in humans at the RHD, while exposures to the predominant circulating metabolite of sofosbuvir (GS-331007) were  $\geq 3$  (rats) and 7 (rabbits) times the exposure in humans at the RHD. In rat pre/postnatal development studies, maternal systemic exposures (AUC) to ledipasvir and GS-331007 were approximately 5 and 7 times, respectively, the exposure in humans at the RHD.

The background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2–4% and 15–20%, respectively.

## Data

### *Animal Data*

*Ledipasvir*: Ledipasvir was administered orally to pregnant rats (up to 100 mg/kg/day) and rabbits (up to 180 mg/kg/day) on gestation days 6 to 18 and 7 to 20, respectively, and also to rats (oral doses up to 100 mg/kg/day) on gestation day 6 to lactation/postpartum day 20. No significant effects on embryo-fetal (rats and rabbits) or pre/postnatal (rats) development were observed at the highest doses tested. Systemic exposures (AUC) to ledipasvir were  $\geq 4$  (rats) and 2 (rabbits) times the exposure in humans at the RHD.

*Sofosbuvir*: Sofosbuvir was administered orally to pregnant rats (up to 500 mg/kg/day) and rabbits (up to 300 mg/kg/day) on gestation days 6 to 18 and 6 to 19, respectively, and also to rats (oral doses up to 500 mg/kg/day) on gestation day 6 to lactation/postpartum day 20. No significant effects on embryo-fetal (rats and rabbits) or pre/postnatal (rats) development were observed at the highest doses tested. Systemic exposures (AUC) to the predominant circulating metabolite of sofosbuvir (GS-331007) were  $\geq 3$  (rats) and 7 (rabbits) times the exposure in humans at the RHD, with exposures increasing during gestation from approximately 3 to 6 (rats) and 7 to 17 (rabbits) times the exposure in humans at the RHD.

## **8.2 Lactation**

### Risk Summary

It is not known whether ledipasvir, sofosbuvir, or their metabolites are present in human breast milk, affect human milk production or have effects on the breastfed infant. When administered to lactating rats, ledipasvir was detected in the plasma of nursing pups likely due to the presence of ledipasvir in milk, without clear effects on nursing pups [see *Data*]. The predominant circulating metabolite of sofosbuvir (GS-331007) was the primary component observed in the milk of lactating rats, without effect on nursing pups.

The development and health benefits of breastfeeding should be considered along with the mother's clinical need for ledipasvir and sofosbuvir and any potential adverse effects on the breastfed child from ledipasvir and sofosbuvir or from the underlying maternal condition.

If ledipasvir and sofosbuvir is administered with ribavirin, the nursing mother's information for ribavirin also applies to this combination regimen. Refer to the ribavirin prescribing information for more information on use during lactation.

### Data

*Ledipasvir*: No effects of ledipasvir on growth and postnatal development were observed in nursing pups at the highest dose tested in rats. Maternal systemic exposure (AUC) to ledipasvir was approximately 5 times the exposure in humans at the RHD. Although not measured directly, ledipasvir was likely present in the milk of lactating rats, since systemic exposure (AUC) to ledipasvir of approximately 25% that of maternal exposure was observed in nursing pups on lactation day 10.

*Sofosbuvir*: No effects of sofosbuvir on growth and postnatal development were observed in nursing pups at the highest dose tested in rats. Maternal systemic exposure (AUC) to the predominant circulating metabolite of sofosbuvir (GS-331007) was approximately 7 times the exposure in humans at the RHD, with exposure of

approximately 2% that of maternal exposure observed in nursing pups on lactation day 10. In a lactation study, sofosbuvir metabolites (primarily GS-331007) were excreted into the milk of lactating rats following administration of a single oral dose of sofosbuvir (20 mg/kg) on lactation day 2, with milk concentrations of approximately 10% that of maternal plasma concentrations observed 1 hour post-dose.

### **8.3 Females and Males of Reproductive Potential**

If ledipasvir and sofosbuvir is administered with ribavirin, the information for ribavirin with regard to pregnancy testing, contraception, and infertility also applies to this combination regimen. Refer to ribavirin prescribing information for additional information.

### **8.4 Pediatric Use**

The safety, pharmacokinetics, and efficacy of ledipasvir and sofosbuvir for treatment of HCV genotype 1 and 4 infection in treatment-naïve and treatment-experienced pediatric patients 3 years of age and older without cirrhosis or with compensated cirrhosis have been established in an open-label, multicenter clinical trial (Study 1116, N=226; 186 treatment-naïve, 40 treatment-experienced) and are comparable to that observed in adults.

The safety and efficacy of ledipasvir and sofosbuvir for treatment of HCV genotypes 5 or 6 infection in pediatric patients 3 years of age and older are supported by comparable ledipasvir, sofosbuvir, and GS-331007 exposures between adults and pediatric patients [see *Dosage and Administration (2.2 and 2.4)*, *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)*, and *Clinical Studies (14.3, 14.6, 14.7)*]. Similar rationale is used to support dosing recommendations for pediatric patients with HCV genotype 1 infection who have decompensated cirrhosis (Child-Pugh B or C) and for pediatric patients with HCV genotype 1 and 4 infection who are liver transplant recipients without cirrhosis or with compensated cirrhosis.

In patients with severe renal impairment, including those requiring dialysis, exposures of GS-331007, the inactive metabolite of sofosbuvir, are *increased* [see *Clinical Pharmacology (12.3)*]. No data are available regarding the safety of ledipasvir and sofosbuvir in pediatric patients with renal *impairment* [see *Use in Specific Populations (8.6)*].

The safety and efficacy of ledipasvir and sofosbuvir have not been established in pediatric patients less than 3 years of age.

In a 5-year follow-up study, the long-term effects of ledipasvir and sofosbuvir on pediatric growth were assessed in 178 pediatric subjects 3 years of age and older treated with ledipasvir and sofosbuvir in Study 1116. No notable effects on growth from baseline through end of study were observed [see *Adverse Reactions (6.1)*]. All subjects who had achieved SVR12 maintained SVR through end of study.

### **8.5 Geriatric Use**

Clinical trials of ledipasvir and sofosbuvir included 225 subjects aged 65 and over (9% of total number of subjects in the clinical studies). No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be

ruled out. No dosage adjustment of ledipasvir and sofosbuvir is warranted in geriatric patients [see *Clinical Pharmacology (12.3)*].

## 8.6 Renal Impairment

No dosage adjustment of ledipasvir and sofosbuvir is recommended for patients with mild, moderate, or severe renal impairment, including ESRD requiring dialysis [see *Dosage and Administration (2.4)*, *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)*, and *Clinical Studies (14.6)*]. No safety data are available in subjects with both decompensated cirrhosis and severe renal impairment, including those on dialysis. Additionally, no safety data are available in pediatric patients with renal impairment [see *Use in Specific Populations (8.4)*]. Refer to ribavirin tablet prescribing information regarding use in patients with renal impairment.

## 8.7 Hepatic Impairment

No dosage adjustment of ledipasvir and sofosbuvir is recommended for patients with mild, moderate, or severe hepatic impairment (Child-Pugh Class A, B, or C) [see *Dosage and Administration (2.3)*, *Clinical Pharmacology (12.3)* and *Clinical Studies (14.5)*].

Clinical and hepatic laboratory monitoring, as clinically indicated, is recommended for patients with decompensated cirrhosis receiving treatment with ledipasvir and sofosbuvir and ribavirin [see *Adverse Reactions (6.1)*].

## 10 OVERDOSAGE

No specific antidote is available for overdose with ledipasvir and sofosbuvir. If overdose occurs, the patient must be monitored for evidence of toxicity. Treatment of overdose with ledipasvir and sofosbuvir consists of general supportive measures including monitoring of vital signs as well as observation of the clinical status of the patient. Hemodialysis is unlikely to result in significant removal of ledipasvir since ledipasvir is highly bound to plasma protein. Hemodialysis can efficiently remove the predominant circulating metabolite of sofosbuvir, GS-331007, with an extraction ratio of 53%.

## 11 DESCRIPTION


Ledipasvir and sofosbuvir tablets (90 mg/400 mg) are fixed-dose combination tablets containing ledipasvir and sofosbuvir for oral administration. Ledipasvir is an HCV NS5A inhibitor and sofosbuvir is a nucleotide analog inhibitor of HCV NS5B polymerase.

Each tablet contains 90 mg ledipasvir and 400 mg sofosbuvir. The tablets include the following inactive ingredients: colloidal silicon dioxide, copovidone, croscarmellose sodium, lactose monohydrate, magnesium stearate, and microcrystalline cellulose. The tablets are film-coated with a coating material containing the following inactive ingredients: FD&C yellow #6/sunset yellow FCF aluminum lake, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide.

*Ledipasvir*: The IUPAC name for ledipasvir is methyl [(2S)-1-{(6S)-6-[5-(9,9-difluoro-7-{2-[(1R,3S,4S)-2-{(2S)-2-[(methoxycarbonyl)amino]-3-methylbutanoyl]}-2-azabicyclo[2.2.1]hept-3-yl]-1H-benzimidazol-6-yl]-9H-fluoren-2-yl)-1H-imidazol-2-yl]-5-azaspiro[2.4]hept-5-yl}-3-methyl-1-oxobutan-2-yl]carbamate.


It has a molecular formula of C<sub>49</sub>H<sub>54</sub>F<sub>2</sub>N<sub>8</sub>O<sub>6</sub> and a molecular weight of 889.00. It has the

following structural formula:



Ledipasvir is practically insoluble (less than 0.1 mg/mL) across the pH range of 3.0–7.5 and is slightly soluble below pH 2.3 (1.1 mg/mL).

*Sofosbuvir*: The IUPAC name for sofosbuvir is (S)-isopropyl 2-((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)-(phenoxy)phosphorylamino)propanoate. It has a molecular formula of  $C_{22}H_{29}FN_3O_9P$  and a molecular weight of 529.45. It has the following structural formula:



Sofosbuvir is a white to off-white crystalline solid with a solubility of at least 2 mg/mL across the pH range of 2–7.7 at 37°C and is slightly soluble in water.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Ledipasvir and sofosbuvir tablets (90 mg/400 mg) are a fixed-dose combination of ledipasvir and sofosbuvir, which are direct-acting antiviral agents against the hepatitis C virus [see *Microbiology* (12.4)].

### 12.2 Pharmacodynamics

#### Cardiac Electrophysiology

Thorough QT studies have been conducted for ledipasvir and sofosbuvir.

The effect of ledipasvir 120 mg twice daily (2.67 times the maximum recommended dosage) for 10 days on QTc interval was evaluated in a randomized, multiple-dose, placebo-, and active-controlled (moxifloxacin 400 mg) three-period crossover thorough QT trial in 59 healthy subjects. At the dose of 120 mg twice daily (2.67 times the maximum recommended dosage), ledipasvir does not prolong QTc interval to any clinically relevant extent.

The effect of sofosbuvir 400 mg (maximum recommended dosage) and 1200 mg (three times the maximum recommended dosage) on QTc interval was evaluated in a randomized, single-dose, placebo-, and active-controlled (moxifloxacin 400 mg) four-period crossover thorough QT trial in 59 healthy subjects. At a dose three times the

maximum recommended dose, sofosbuvir does not prolong QTc to any clinically relevant extent.

## 12.3 Pharmacokinetics

### Absorption

The pharmacokinetic properties of ledipasvir, sofosbuvir, and the predominant circulating metabolite GS-331007 have been evaluated in healthy adult subjects and in subjects with chronic hepatitis C. Following oral administration of ledipasvir and sofosbuvir tablets (90 mg/400 mg), ledipasvir median peak concentrations were observed 4 to 4.5 hours post-dose. Sofosbuvir was absorbed quickly and the peak median plasma concentration was observed ~0.8 to 1 hour post-dose. Median peak plasma concentration of GS-331007 was observed between 3.5 to 4 hours post-dose.

Based on the population pharmacokinetic analysis in HCV-infected subjects, geometric mean steady-state  $AUC_{0-24}$  for ledipasvir (N=2113), sofosbuvir (N=1542), and GS-331007 (N=2113) were 7290, 1320, and 12,000 ng•hr/mL, respectively. Steady-state  $C_{max}$  for ledipasvir, sofosbuvir, and GS-331007 were 323, 618, and 707 ng/mL, respectively. Sofosbuvir and GS-331007  $AUC_{0-24}$  and  $C_{max}$  were similar in healthy adult subjects and subjects with HCV infection. Relative to healthy subjects (N=191), ledipasvir  $AUC_{0-24}$  and  $C_{max}$  were 24% lower and 32% lower, respectively, in HCV-infected subjects.

### Effect of Food

Relative to fasting conditions, the administration of a single dose of ledipasvir and sofosbuvir tablets (90 mg/400 mg) with a moderate fat (~600 kcal, 25% to 30% fat) or high fat (~1000 kcal, 50% fat) meal increased sofosbuvir  $AUC_{0-inf}$  by approximately 2-fold, but did not significantly affect sofosbuvir  $C_{max}$ . The exposures of GS-331007 and ledipasvir were not altered in the presence of either meal type. The response rates in Phase 3 trials were similar in HCV-infected subjects who received ledipasvir and sofosbuvir tablets (90 mg/400 mg) with food or without food. Ledipasvir and sofosbuvir can be administered without regard to food.

### Distribution

Ledipasvir is greater than 99.8% bound to human plasma proteins. After a single 90 mg dose of [ $^{14}C$ ]-ledipasvir in healthy subjects, the blood to plasma ratio of  $^{14}C$ -radioactivity ranged between 0.51 and 0.66.

Sofosbuvir is approximately 61–65% bound to human plasma proteins and the binding is independent of drug concentration over the range of 1 microgram/mL to 20 microgram/mL. Protein binding of GS-331007 was minimal in human plasma. After a single 400 mg dose of [ $^{14}C$ ]-sofosbuvir in healthy subjects, the blood to plasma ratio of  $^{14}C$ -radioactivity was approximately 0.7.

### Metabolism

In vitro, no detectable metabolism of ledipasvir was observed by human CYP1A2, CYP2C8, CYP2C9, CYP2C19, CYP2D6, and CYP3A4. Evidence of slow oxidative metabolism via an unknown mechanism has been observed. Following a single dose of 90 mg [ $^{14}C$ ]-ledipasvir, systemic exposure was almost exclusively to the parent drug (greater than 98%). Unchanged ledipasvir is the major species present in feces.

Sofosbuvir is extensively metabolized in the liver to form the pharmacologically active

nucleoside analog triphosphate GS-461203. The metabolic activation pathway involves sequential hydrolysis of the carboxyl ester moiety catalyzed by human cathepsin A (CatA) or carboxylesterase 1 (CES1) and phosphoramidate cleavage by histidine triad nucleotide-binding protein 1 (HINT1) followed by phosphorylation by the pyrimidine nucleotide biosynthesis pathway. Dephosphorylation results in the formation of nucleoside metabolite GS-331007 that cannot be efficiently rephosphorylated and lacks anti-HCV activity in vitro. After a single 400 mg oral dose of [<sup>14</sup>C]-sofosbuvir, GS-331007 accounted for approximately greater than 90% of total systemic exposure.

### Elimination

Following a single 90 mg oral dose of [<sup>14</sup>C]-ledipasvir, mean total recovery of the [<sup>14</sup>C]-radioactivity in feces and urine was approximately 87%, with most of the radioactive dose recovered from feces (approximately 86%). Unchanged ledipasvir excreted in feces accounted for a mean of 70% of the administered dose and the oxidative metabolite M19 accounted for 2.2% of the dose. These data indicate that biliary excretion of unchanged ledipasvir is a major route of elimination, with renal excretion being a minor pathway (approximately 1%). The median terminal half-life of ledipasvir following administration of ledipasvir and sofosbuvir tablets (90 mg/400 mg) was 47 hours.

Following a single 400 mg oral dose of [<sup>14</sup>C]-sofosbuvir, mean total recovery of the dose was greater than 92%, consisting of approximately 80%, 14%, and 2.5% recovered in urine, feces, and expired air, respectively. The majority of the sofosbuvir dose recovered in urine was GS-331007 (78%) while 3.5% was recovered as sofosbuvir. These data indicate that renal clearance is the major elimination pathway for GS-331007. The median terminal half-lives of sofosbuvir and GS-331007 following administration of ledipasvir and sofosbuvir tablets (90 mg/400 mg) were 0.5 and 27 hours, respectively.

### Specific Populations

*Race:* Population pharmacokinetics analysis in HCV-infected subjects indicated that race had no clinically relevant effect on the exposure of ledipasvir, sofosbuvir, and GS-331007.

*Gender:* Population pharmacokinetics analysis in HCV-infected subjects indicated that gender had no clinically relevant effect on the exposure of sofosbuvir and GS-331007. AUC and C<sub>max</sub> of ledipasvir were 77% and 58% higher, respectively, in females than males; however, the relationship between gender and ledipasvir exposures was not considered clinically relevant, as high response rates (SVR12 >90%) were achieved in male and female subjects across the Phase 3 studies and the safety profiles are similar in females and males.

*Pediatric Patients:* The pharmacokinetics of ledipasvir, sofosbuvir, and GS-331007 were determined in HCV genotype 1, 3, or 4 infected pediatric subjects 3 years of age and older receiving a daily dose of ledipasvir and sofosbuvir as described below in Table 7. Exposures in pediatric subjects were similar to those observed in adults.

**Table 7 Pharmacokinetic Properties of Ledipasvir, Sofosbuvir, and GS-331007 in Pediatric Subjects 3 Years of Age and Older\***

Weight Group	Dose	PK Parameter	Geometric Mean (%CV)		
			Ledipasvir	Sofosbuvir	GS-331007

≥35 kg <sup>†</sup>	90/400 mg	AUC <sub>tau</sub> (ng•hr/mL)	11200 (45.7)	1350 (45.2)	13600 (18.9)
		C <sub>max</sub> (ng/mL)	550 (44.2)	660 (51.1)	921 (17.8)
17 to <35 kg <sup>‡</sup>	45/200 mg	AUC <sub>tau</sub> (ng•hr/mL)	8750 (46.6)	1420 (34.2)	10700 (30.9)
		C <sub>max</sub> (ng/mL)	440 (42.7)	690 (24.8)	958 (26.1)
<17 kg <sup>§</sup>	33.75/150 mg	AUC <sub>tau</sub> (ng•hr/mL)	7460 (31.0)	1720 (23.2)	12200 (15.2)
		C <sub>max</sub> (ng/mL)	405 (25.7)	791 (16.6)	1070 (13.0)

\* Population PK derived parameters

† Ledipasvir N=100; Sofosbuvir N=72; GS-331007 N=100

‡ Ledipasvir N=86; Sofosbuvir N=66; GS-331007 N=86

§ Ledipasvir N=9; Sofosbuvir N=9; GS-331007 N=9

The pharmacokinetics of ledipasvir, sofosbuvir, and GS-331007 have not been established in pediatric subjects less than 3 years of age [see *Use in Specific Populations (8.4) and Clinical Studies (14.7)*].

**Geriatric Patients:** Population pharmacokinetic analysis in HCV-infected subjects showed that within the age range (18 to 80 years) analyzed, age did not have a clinically relevant effect on the exposure to ledipasvir, sofosbuvir, and GS-331007 [see *Use in Specific Populations (8.5)*].

**Patients with Renal Impairment:** The pharmacokinetics of ledipasvir were studied with a single dose of 90 mg ledipasvir in HCV negative subjects with severe renal impairment (eGFR less than 30 mL/min by Cockcroft-Gault). No clinically relevant differences in ledipasvir pharmacokinetics were observed between healthy subjects and subjects with severe renal impairment.

The pharmacokinetics of sofosbuvir were studied in HCV negative subjects with mild (eGFR between 50 to less than 80 mL/min/1.73 m<sup>2</sup>), moderate (eGFR between 30 to less than 50 mL/min/1.73 m<sup>2</sup>), severe renal impairment (eGFR less than 30 mL/min/1.73 m<sup>2</sup>), and subjects with ESRD requiring hemodialysis following a single 400 mg dose of sofosbuvir. Relative to subjects with normal renal function (eGFR greater than 80 mL/min/1.73 m<sup>2</sup>), the sofosbuvir AUC<sub>0-inf</sub> was 61%, 107%, and 171% higher in mild, moderate, and severe renal impairment, while the GS-331007 AUC<sub>0-inf</sub> was 55%, 88%, and 451% higher, respectively. In subjects with ESRD, relative to subjects with normal renal function, sofosbuvir and GS-331007 AUC<sub>0-inf</sub> was 28% and 1280% higher when sofosbuvir was dosed 1 hour before hemodialysis compared with 60% and 2070% higher when sofosbuvir was dosed 1 hour after hemodialysis, respectively. A 4-hour hemodialysis session removed approximately 18% of administered dose [see *Dosage and Administration (2.6) and Use in Specific Populations (8.6)*].

The pharmacokinetics of ledipasvir, sofosbuvir, and GS-331007 were studied in HCV-infected subjects with severe renal impairment or ESRD requiring dialysis treated with ledipasvir and sofosbuvir tablets (90 mg/400 mg) for 8, 12, or 24 weeks. The results were generally consistent with those observed in HCV-negative subjects with ESRD requiring dialysis.

**Patients with Hepatic Impairment:** The pharmacokinetics of ledipasvir were studied with a single dose of 90 mg ledipasvir in HCV negative subjects with severe hepatic impairment (Child-Pugh Class C). Ledipasvir plasma exposure (AUC<sub>0-inf</sub>) was similar in subjects with

severe hepatic impairment and control subjects with normal hepatic function. Population pharmacokinetics analysis in HCV-infected subjects indicated that cirrhosis (including decompensated cirrhosis) had no clinically relevant effect on the exposure of ledipasvir [see *Use in Specific Populations (8.7)*].

The pharmacokinetics of sofosbuvir were studied following 7-day dosing of 400 mg sofosbuvir in HCV-infected subjects with moderate and severe hepatic impairment (Child-Pugh Class B and C). Relative to subjects with normal hepatic function, the sofosbuvir AUC<sub>0-24</sub> were 126% and 143% higher in moderate and severe hepatic impairment, while the GS-331007 AUC<sub>0-24</sub> were 18% and 9% higher, respectively. Population pharmacokinetics analysis in HCV-infected subjects indicated that cirrhosis (including decompensated cirrhosis) had no clinically relevant effect on the exposure of sofosbuvir and GS-331007 [see *Use in Specific Populations (8.7)* and *Clinical Studies (14.5)*].

### Drug Interaction Studies

Ledipasvir and sofosbuvir are substrates of drug transporters P-gp and BCRP while GS-331007 is not. P-gp inducers (e.g., rifampin or St. John's wort) may decrease ledipasvir and sofosbuvir plasma concentrations, leading to reduced therapeutic effect of ledipasvir and sofosbuvir, and the use with P-gp inducers is not recommended with ledipasvir and sofosbuvir [see *Warnings and Precautions (5.3)*]. Coadministration with drugs that inhibit P-gp and/or BCRP may increase ledipasvir and sofosbuvir plasma concentrations without increasing GS-331007 plasma concentration; ledipasvir and sofosbuvir may be coadministered with P-gp and/or BCRP inhibitors. Neither ledipasvir nor sofosbuvir is a substrate for hepatic uptake transporters OCT1, OATP1B1, or OATP1B3. GS-331007 is not a substrate for renal transporters, including organic anion transporter OAT1 or OAT3, or organic cation transporter OCT2.

Ledipasvir is subject to slow oxidative metabolism via an unknown mechanism. In vitro, no detectable metabolism of ledipasvir by CYP enzymes has been observed. Biliary excretion of unchanged ledipasvir is a major route of elimination. Sofosbuvir is not a substrate for CYP and UGT1A1 enzymes. Clinically significant drug interactions with ledipasvir and sofosbuvir mediated by CYP or UGT1A1 enzymes are not expected.

The effects of coadministered drugs on the exposure of ledipasvir, sofosbuvir, and GS-331007 are shown in Table 8 [see *Drug Interactions (7)*].

**Table 8 Drug Interactions: Changes in Pharmacokinetic Parameters for Ledipasvir, Sofosbuvir, and the Predominant Circulating Metabolite GS-331007 in the Presence of the Coadministered Drug\***

Coadministered Drug	Dose of Coadministered Drug (mg)	Ledipasvir Dose (mg)	Sofosbuvir Dose (mg)	N	Mean Ratio (90% CI) of Ledipasvir, Sofosbuvir, and GS-331007 PK With/Without Coadministered Drug No Effect=1.00			
					C <sub>max</sub>	AUC	C <sub>min</sub>	
Atazanavir/	300/1000				ledipasvir	1.68 (1.54, 1.84)	1.96 (1.74, 2.21)	2.18 (1.91, 2.50)

ritonavir + emtricitabine/tenofovir DF†,‡	500/100 + 200/300 once daily	90 once daily	400 once daily	24	sofosbuvir	1.01 (0.88, 1.15)	1.11 (1.02, 1.21)	NA
					GS-331007	1.17 (1.12, 1.23)	1.31 (1.25, 1.36)	1.42 (1.34, 1.49)
Carbamazepine	300 twice daily	ND	400 single dose	24	sofosbuvir	0.52 (0.43, 0.62)	0.52 (0.46, 0.59)	NA
					GS-331007	1.04 (0.97, 1.11)	0.99 (0.94, 1.04)	NA
Cyclosporine	600 single dose	ND	400 single dose	19	sofosbuvir	2.54 (1.87, 3.45)	4.53 (3.26, 6.30)	NA
					GS-331007	0.60 (0.53, 0.69)	1.04 (0.90, 1.20)	NA
Darunavir/ritonavir	800/100 once daily	90 once daily	ND	23	ledipasvir	1.45 (1.34, 1.56)	1.39 (1.28, 1.49)	1.39 (1.29, 1.51)
		ND	400 single dose	18	sofosbuvir	1.45 (1.10, 1.92)	1.34 (1.12, 1.59)	NA
Darunavir/ritonavir + emtricitabine/tenofovir DF†	800/100 + 200/300 once daily	90 once daily	400 once daily	23	ledipasvir	1.11 (0.99, 1.24)	1.12 (1.00, 1.25)	1.17 (1.04, 1.31)
					sofosbuvir	0.63 (0.52, 0.75)	0.73 (0.65, 0.82)	NA
Efavirenz/emtricitabine/tenofovir DF§	600/200/300 once daily	90 once daily	400 once daily	14	GS-331007	1.10 (1.04, 1.16)	1.20 (1.16, 1.24)	1.26 (1.20, 1.32)
					ledipasvir	0.66 (0.59, 0.75)	0.66 (0.59, 0.75)	0.66 (0.57, 0.76)
Elvitegravir/cobicistat/emtricitabine/	150/150/200/10	90 once	400 once	20	sofosbuvir	1.03 (0.87, 1.23)	0.94 (0.81, 1.10)	NA
					ledipasvir	1.65 (1.53, 1.78)	1.79 (1.64, 1.96)	1.93 (1.74, 2.15)
					sofosbuvir	1.28 (1.12, 1.44)	1.47 (1.25, 1.69)	NA

emtricitabine/ tenofovir alafenamide	once daily	daily	daily	30	sofosbuvir	(1.15, 1.47)	(1.55, 1.59)	NA
					GS-331007	1.29 (1.24, 1.35)	1.48 (1.44, 1.53)	1.66 (1.60, 1.73)
Famotidine	40 single dose simultaneously with ledipasvir and sofosbuvir tablets	90 single dose	400 single dose	12	ledipasvir	0.80 (0.69, 0.93)	0.89 (0.76, 1.06)	NA
					sofosbuvir	1.15 (0.88, 1.50)	1.11 (1.00, 1.24)	NA
					GS-331007	1.06 (0.97, 1.14)	1.06 (1.02, 1.11)	NA
	40 single dose 12 hours prior to ledipasvir and sofosbuvir tablets			12	ledipasvir	0.83 (0.69, 1.00)	0.98 (0.80, 1.20)	NA
					sofosbuvir	1.00 (0.76, 1.32)	0.95 (0.82, 1.10)	NA
					GS-331007	1.13 (1.07, 1.20)	1.06 (1.01, 1.12)	NA
Methadone	30 to 130 daily	ND	400 once daily	14	sofosbuvir	0.95 (0.68, 1.33)	1.30 (1.00, 1.69)	NA
					GS-331007	0.73 (0.65, 0.83)	1.04 (0.89, 1.22)	NA
Omeprazole	20 once daily simultaneously with ledipasvir and sofosbuvir tablets	90 single dose	400 single dose	16	ledipasvir	0.89 (0.61, 1.30)	0.96 (0.66, 1.39)	NA
					sofosbuvir	1.12 (0.88, 1.42)	1.00 (0.80, 1.25)	NA
					GS-331007	1.14 (1.01, 1.29)	1.03 (0.96, 1.12)	NA
	20 once daily 2 hours prior to ledipasvir	30 single dose	ND	17	ledipasvir	0.52 (0.41, 0.66)	0.58 (0.48, 0.71)	NA
Rifabutin	300 once daily	ND	400 single dose	20	sofosbuvir	0.64 (0.53, 0.77)	0.76 (0.63, 0.91)	NA
					GS-331007	1.15 (1.03, 1.27)	1.03 (0.95, 1.12)	NA
		90 single dose <sup>¶</sup>	ND	31	ledipasvir	0.65 (0.56, 0.76)	0.41 (0.36, 0.48)	NA

Rifampin	600 once daily	ND	400 single dose	17	sofosbuvir	0.23 (0.19, 0.29)	0.28 (0.24, 0.32)	NA
					GS-331007	1.23 (1.14, 1.34)	0.95 (0.88, 1.03)	NA
Simeprevir	150 once daily	30 once daily	ND	22	ledipasvir	1.81 (1.69, 2.94)	1.92 (1.77, 2.07)	NA
Tacrolimus	5 single dose	ND	400 single dose	16	sofosbuvir	0.97 (0.65, 1.43)	1.13 (0.81, 1.57)	NA
					GS-331007	0.97 (0.83, 1.14)	1.00 (0.87, 1.13)	NA

NA = not available/not applicable, ND = not dosed.

tenofovir DF = tenofovir disoproxil fumarate

\* All interaction studies conducted in healthy volunteers.

† Data generated from simultaneous dosing with ledipasvir and sofosbuvir. Staggered administration (12 hours apart) of atazanavir/ritonavir + emtricitabine/tenofovir DF or darunavir/ritonavir + emtricitabine/tenofovir DF and ledipasvir and sofosbuvir provided similar results.

‡ The effects of atazanavir/ritonavir on ledipasvir and sofosbuvir are similar with or without the presence of emtricitabine/tenofovir DF.

§ Administered as Atripla<sup>®</sup> (efavirenz, emtricitabine, tenofovir DF).

¶ This study was conducted in the presence of two other investigational HCV direct-acting agents.

No effect on the pharmacokinetic parameters of ledipasvir, sofosbuvir, and GS-331007 was observed with raltegravir and the combination of abacavir and lamivudine; emtricitabine, rilpivirine, and tenofovir disoproxil fumarate; or dolutegravir, emtricitabine, and tenofovir disoproxil fumarate.

Ledipasvir is an inhibitor of drug transporter P-gp and breast cancer resistance protein (BCRP) and may increase intestinal absorption of coadministered substrates for these transporters. Ledipasvir is an inhibitor of transporters OATP1B1, OATP1B3, and BSEP only at concentrations exceeding those achieved in clinic. Ledipasvir is not an inhibitor of transporters MRP2, MRP4, OCT2, OAT1, OAT3, MATE1, and OCT1. The drug-drug interaction potential of ledipasvir is primarily limited to the intestinal inhibition of P-gp and BCRP. Clinically relevant transporter inhibition by ledipasvir in the systemic circulation is not expected due to its high protein binding. Sofosbuvir and GS-331007 are not inhibitors of drug transporters P-gp, BCRP, MRP2, BSEP, OATP1B1, OATP1B3, and OCT1, and GS-331007 is not an inhibitor of OAT1, OCT2, and MATE1. Ledipasvir, sofosbuvir, and GS-331007 are not inhibitors or inducers of CYP or UGT1A1 enzymes.

The effects of ledipasvir or sofosbuvir on the exposure of coadministered drugs are shown in Table 9 [see *Drug Interactions (7)*].

**Table 9 Drug Interactions: Changes in Pharmacokinetic Parameters for Coadministered Drug in the Presence of Ledipasvir, Sofosbuvir, or Ledipasvir and Sofosbuvir\***

								Mean Ratio (90% CI) of
--	--	--	--	--	--	--	--	---------------------------

Coadministered Drug	Dose of Coadministered Drug (mg)	Ledipasvir Dose (mg)	Sofosbuvir Dose (mg)	N	Coadministered Drug PK With/Without Ledipasvir, Sofosbuvir, or Ledipasvir and Sofosbuvir No Effect=1.00		
					C <sub>max</sub>	AUC	C <sub>min</sub>
Atazanavir/ ritonavir + emtricitabine /tenofovir DF <sup>†,‡,§</sup>	atazanavir 300 once daily	90 once daily	400 once daily	24	1.07 (0.99, 1.14)	1.27 (1.18, 1.37)	1.63 (1.45, 1.84)
	ritonavir 100 once daily				0.86 (0.79, 0.93)	0.97 (0.89, 1.05)	1.45 (1.27, 1.64)
	tenofovir DF 300 once daily				1.47 (1.37, 1.58)	1.35 (1.29, 1.42)	1.47 (1.38, 1.57)
Darunavir/ ritonavir + emtricitabine/ tenofovir DF <sup>†,§</sup>	darunavir 800 once daily	90 once daily	400 once daily	23	1.01 (0.96, 1.06)	1.04 (0.99, 1.08)	1.08 (0.98, 1.20)
	ritonavir 100 once daily				1.17 (1.01, 1.35)	1.25 (1.15, 1.36)	1.48 (1.34, 1.63)
	tenofovir DF 300 once daily				1.64 (1.54, 1.74)	1.50 (1.42, 1.59)	1.59 (1.49, 1.70)
Elvitegravir/ cobicistat/ emtricitabine/ tenofovir alafenamide	elvitegravir 150 once daily	90 once daily	400 once daily	30	0.98 (0.90, 1.07)	1.11 (1.02, 1.20)	1.46 (1.28, 1.66)
	cobicistat 150 once daily				1.23 (1.15, 1.32)	1.53 (1.45, 1.62)	3.25 (2.88, 3.67)
	tenofovir alafenamide 10 once daily				0.90 (0.73, 1.11)	0.86 (0.78, 0.95)	NA
Norelgestromin		90 once daily	ND		1.02 (0.89, 1.16)	1.03 (0.90, 1.18)	1.09 (0.91, 1.31)
		ND	400 once daily		1.07 (0.94, 1.22)	1.06 (0.92, 1.21)	1.07 (0.89, 1.28)
Norgestrel	norgestimate 0.180/0.215/0.25/ethinyl estradiol 0.025 once daily	90 once daily	ND	15	1.03 (0.87, 1.23)	0.99 (0.82, 1.20)	1.00 (0.81, 1.23)
		ND	400 once daily		1.18 (0.99, 1.41)	1.19 (0.98, 1.45)	1.23 (1.00, 1.51)
		90 once			1.40	1.20	0.98

Ethinyl estradiol		90 once daily	ND		(1.18, 1.66)	(1.04, 1.39)	(0.79, 1.22)
		ND	400 once daily		1.15 (0.97, 1.36)	1.09 (0.94, 1.26)	0.99 (0.80, 1.23)
Midazolam	2.5 single dose	90 single dose	ND	30	1.07 (1.00, 1.14)	0.99 (0.95, 1.04)	NA
					0.95 (0.87, 1.04)	0.89 (0.84, 0.95)	NA
Raltegravir	400 twice daily	90 once daily	ND	28	0.82 (0.66, 1.02)	0.85 (0.70, 1.02)	1.15 (0.90, 1.46)
		ND	400 single dose	19	0.57 (0.44, 0.75)	0.73 (0.59, 0.91)	0.95 (0.81, 1.12)
Simeprevir	150 once daily	30 once daily	ND	22	2.61 (2.39, 2.86)	2.69 (2.44, 2.96)	NA
Tacrolimus	5 single dose	ND	400 single dose	16	0.73 (0.59, 0.90)	1.09 (0.84, 1.40)	NA
Tenofovir DF	300 once daily <sup>¶</sup>	90 once daily	400 once daily	15	1.79 (1.56, 2.04)	1.98 (1.77, 2.23)	2.63 (2.32, 2.97)

NA = not available/not applicable, ND = not dosed.

tenofovir DF = tenofovir disoproxil fumarate

\* All interaction studies conducted in healthy volunteers.

† Data generated from simultaneous dosing with ledipasvir and sofosbuvir. Staggered administration (12 hours apart) of atazanavir/ritonavir + emtricitabine/tenofovir DF or darunavir/ritonavir + emtricitabine/tenofovir DF and ledipasvir and sofosbuvir provided similar results.

‡ The effects of ledipasvir and sofosbuvir on atazanavir and ritonavir are similar with or without the presence of emtricitabine/tenofovir DF.

§ This magnitude of change in tenofovir exposure does not reflect the approximately 60–80% increase caused by the effects of an HIV PI/ritonavir and the effect of food. Therefore, tenofovir exposure is approximately 130% higher when administered as tenofovir DF + atazanavir/ritonavir + ledipasvir and sofosbuvir or tenofovir DF + darunavir/ritonavir + ledipasvir and sofosbuvir and with food as compared to the tenofovir exposure observed following fasted administration of tenofovir DF-based regimens that do not contain an HIV PI/ritonavir and ledipasvir and sofosbuvir.

¶ Administered as Atripla (efavirenz, emtricitabine, tenofovir DF). The effects of ledipasvir and sofosbuvir on tenofovir exposures are similar when tenofovir is administered as Atripla, Complera<sup>®</sup>, or Truvada + dolutegravir.

No effect on the pharmacokinetic parameters of the following coadministered drugs was observed with ledipasvir or sofosbuvir: abacavir, cyclosporine, darunavir/ritonavir, dolutegravir, efavirenz, emtricitabine, lamivudine, methadone, or rilpivirine.

## 12.4 Microbiology

### Mechanism of Action

Ledipasvir is an inhibitor of the HCV NS5A protein, which is required for viral replication. Resistance selection in cell culture and cross-resistance studies indicate ledipasvir targets NS5A as its mode of action.

Sofosbuvir is an inhibitor of the HCV NS5B RNA-dependent RNA polymerase, which is required for viral replication. Sofosbuvir is a nucleotide prodrug that undergoes intracellular metabolism to form the pharmacologically active uridine analog triphosphate (GS-461203), which can be incorporated into HCV RNA by the NS5B polymerase and acts as a chain terminator. In a biochemical assay, GS-461203 inhibited the polymerase activity of the recombinant NS5B from HCV genotypes 1b and 4a with  $IC_{50}$  values of 3.3 and 2.7  $\mu$ M, respectively. GS-461203 is neither an inhibitor of human DNA and RNA polymerases nor an inhibitor of mitochondrial RNA polymerase.

### Antiviral Activity

In HCV replicon assays, the  $EC_{50}$  values of ledipasvir against full-length replicons from genotypes 1a and 1b were 0.031 nM and 0.004 nM, respectively. The median  $EC_{50}$  values of ledipasvir against chimeric replicons encoding NS5A sequences from clinical isolates from treatment-naïve HCV-infected subjects were 0.02 nM for genotype 1a (range 0.007–1.0 nM; N=23) and 0.006 nM for genotype 1b (range 0.002–1.0 nM; N=34). Ledipasvir had median  $EC_{50}$  values ranging between 0.002 nM to 0.16 nM against 11 genotype 4 subtypes (4a, 4d, 4n, 4r, 4o, 4c, 4f, 4k, 4l, 4m, and 4t). The median  $EC_{50}$  value for subtype 4b was 199.6 nM (range 0.66–1799 nM; N=3); the two 4b isolates with  $EC_{50}$  values greater than 100 nM had NS5A resistance-associated polymorphisms L30S+M31M+P58S+Y93H. The median  $EC_{50}$  value of ledipasvir was 0.03 nM against genotype 5a isolates (range 0.008–0.081 nM; N=35). For genotype 6, the  $EC_{50}$  values for ledipasvir varied by subtype. Subtypes 6a and 6h had median  $EC_{50}$  values of 0.55 and 0.17 nM, respectively. For subtypes 6e, 6l, 6n, 6q, 6k, and 6m, the median  $EC_{50}$  values ranged from 60.6 nM to 430.1 nM.

In HCV replicon assays, the  $EC_{50}$  values of sofosbuvir against full-length replicons from genotypes 1a, 1b, and 4a, and chimeric 1b replicons encoding NS5B from genotypes 5a or 6a ranged from 14–110 nM. The median  $EC_{50}$  value of sofosbuvir against chimeric replicons encoding NS5B sequences from clinical isolates was 62 nM for genotype 1a (range 29–128 nM; N=67) and 102 nM for genotype 1b (range 45–170 nM; N=29). In replication competent virus assays, the  $EC_{50}$  value of sofosbuvir against genotype 1a was 30 nM. Evaluation of sofosbuvir in combination with ledipasvir showed no antagonistic effect in reducing HCV RNA levels in replicon cells.

### Resistance

#### *In Cell Culture*

HCV replicons with reduced susceptibility to ledipasvir have been selected in cell culture for genotypes 1a and 1b. Reduced susceptibility to ledipasvir was associated with the primary NS5A amino acid substitution Y93H in both genotypes 1a and 1b. Additionally, a Q30E substitution emerged in genotype 1a replicons. Site-directed mutagenesis of the Y93H in both genotypes 1a and 1b, as well as the Q30E substitution in genotype 1a, conferred high levels of reduced susceptibility to ledipasvir (fold change in  $EC_{50}$  greater than 1000-fold).

HCV replicons with reduced susceptibility to sofosbuvir have been selected in cell culture for multiple genotypes including 1b, 4a, 5a, and 6a. Reduced susceptibility to sofosbuvir was associated with the NS5B substitution S282T in all replicon genotypes examined. An M289L substitution developed along with the S282T substitution in genotype 5 and 6

replicons. Site-directed mutagenesis of the S282T substitution in replicons of 8 genotypes conferred 2- to 18-fold reduced susceptibility to sofosbuvir.

## *In Clinical Trials*

### Genotype 1

In a pooled analysis of subjects who received ledipasvir and sofosbuvir tablets in Phase 3 trials (ION-3, ION-1, and ION-2), 37 subjects (29 with genotype 1a HCV and 8 with genotype 1b HCV) qualified for resistance analysis due to virologic failure (35 with virologic relapse, 2 with breakthrough on-treatment due to documented non-adherence). Post-baseline NS5A and NS5B deep nucleotide sequence analysis data (assay sensitivity of 1%) were available for 37/37 and 36/37 subjects' viruses, respectively.

Of the 29 genotype 1a virologic failure subjects, 55% (16/29) of subjects had virus with emergent NS5A resistance-associated substitutions K24R, M28T/V, Q30R/H/K/L, L31M, or Y93H/N at failure. Five of these 16 subjects' viruses also had baseline NS5A polymorphisms at resistance-associated amino acid positions. The most common substitutions detected at failure were Q30R, Y93H or N, and L31M.

Of the 8 genotype 1b virologic failure subjects, 88% (7/8) had virus with emergent NS5A resistance-associated substitutions L31V/M/I or Y93H at failure. Virus from three of these 7 subjects also had baseline NS5A polymorphisms at resistance-associated positions. The most common substitution detected at failure was Y93H.

At failure, 38% (14/37) of virologic failure subjects' viruses had 2 or more NS5A substitutions at resistance-associated positions.

In the SOLAR-1 and SOLAR-2 trials (liver transplant recipients or subjects with decompensated liver disease), there were 24 virologic failures with genotype 1 infection (20 relapsers and 4 subjects who discontinued treatment prior to achieving HCV RNA <LLOQ). Treatment-emergent NS5A resistance-associated substitutions K24R, M28T, Q30R/H/K, L31V, H58D/P, and/or Y93H/C were detected in 14/17 (82%) genotype 1a virologic failure subjects, and R30Q, L31M, and/or Y93H/N were detected in 6/7 (86%) genotype 1b virologic failure subjects.

In phenotypic analyses, post-baseline isolates from subjects who harbored NS5A resistance-associated substitutions at failure showed 20- to >243-fold reduced susceptibility to ledipasvir.

Treatment-emergent NS5B substitutions L159 (n=1) and V321 (n=2) previously associated with sofosbuvir failure were detected in the Phase 3 trials (ION-3, ION-1, and ION-2). In addition, treatment-emergent NS5B substitutions at highly conserved positions D61G (n=3), A112T (n=2), E237G (n=2), and S473T (n=1) were detected at low frequency by next generation sequencing in treatment failure subjects infected with HCV genotype 1a. The D61G substitution was previously described in subjects infected with HCV genotype 1a in a liver pre-transplant trial. The E237G substitution was detected in 3 subjects infected with HCV GT1a in the SOLAR-1 and SOLAR-2 trials. The clinical significance of these substitutions is currently unknown. The sofosbuvir-associated resistance substitution S282T in NS5B was not detected in any failure isolate from the Phase 3 trials. NS5B substitutions S282T, L320V/I, and V321I in combination with NS5A substitutions L31M, Y93H, and Q30L were detected in one subject at failure following 8 weeks of treatment with ledipasvir and sofosbuvir tablets (90 mg/400 mg) in a Phase 2 trial.

## Genotype 4, 5 or 6

Resistance analysis was performed for 6 relapse subjects infected with HCV genotype 4 (Study 1119 and ION-4, N=3), genotype 5 (Study 1119, N=2) or genotype 6 (ELECTRON-2, N=1) and treated with ledipasvir and sofosbuvir tablets (90 mg/400 mg) for 12 weeks. All the relapse subjects with NS5A sequencing data (5 of 6) had pretreatment NS5A resistance-associated polymorphisms (single or combinations at positions 24, 28, 30, 31, and 58). NS5A resistance substitutions (Y93C or L28V) emerged in two of the genotype 4 relapse subjects post-treatment who also had NS5A polymorphisms pretreatment that were retained post-treatment. Two of the relapsers with genotype 4 HCV infection had an NS5B V321I substitution pretreatment, which was retained post-treatment. Three of the relapse subjects (1 each for genotype 4, 5, and 6) had virus with emergent sofosbuvir resistance-associated substitution S282T at relapse; the genotype 5 relapse subject also had emergent nucleoside inhibitor substitution M289I.

## Persistence of Resistance-Associated Substitutions

Certain NS5A inhibitor resistance-associated substitutions have been found to persist for >1 year in some patients. The long-term clinical impact of the emergence or persistence of virus containing ledipasvir or sofosbuvir resistance-associated substitutions is unknown.

## Effect of Baseline HCV Polymorphisms on Treatment Response

### *Adults*

#### *Genotype 1*

Analyses were conducted to explore the association between pre-existing baseline NS5A polymorphisms at resistance-associated positions and relapse rates. In the pooled analysis of the Phase 3 trials, 23% (370/1589) of subjects' virus had baseline NS5A polymorphisms at resistance-associated positions (any change from reference at NS5A amino acid positions 24, 28, 30, 31, 58, 92, or 93) identified by population or analysis of deep nucleotide sequences with a 15% frequency threshold.

In treatment-naïve subjects whose virus had baseline NS5A polymorphisms at resistance-associated positions in Studies ION-1 and ION-3, relapse rates were 6% (3/48) after 8 weeks and 1% (1/113) after 12 weeks of treatment with ledipasvir and sofosbuvir tablets (90 mg/400 mg). Relapse rates among subjects without baseline NS5A polymorphisms at resistance-associated positions were 5% (8/167) after 8 weeks and 1% (3/306) after 12 weeks of treatment with ledipasvir and sofosbuvir tablets.

In treatment-experienced subjects in Study ION-2 whose virus had baseline NS5A polymorphisms at resistance-associated positions, relapse rates were 22% (5/23) after 12 weeks and 0% (0/19) after 24 weeks of treatment with ledipasvir and sofosbuvir tablets (90 mg/400 mg). In another study in treatment-experienced subjects (SIRIUS), 0/15 (0%) subjects with NS5A polymorphisms at resistance-associated positions relapsed after 12 weeks of treatment with ledipasvir and sofosbuvir tablets (90 mg/400 mg) + ribavirin compared to 2/15 (13%) subjects treated with 24 weeks of ledipasvir and sofosbuvir tablets (90 mg/400 mg).

SVR was achieved in all 24 subjects (N=20 with L159F+C316N; N=1 with L159F; and N=3 with N142T) who had baseline polymorphisms associated with resistance to sofosbuvir and/or other NS5B nucleoside inhibitors. The NS5B S282T substitution

associated with resistance to sofosbuvir was not detected in the baseline NS5B sequence of any subject in Phase 3 trials by population or deep nucleotide sequence analysis.

In the SOLAR-1 and SOLAR-2 trials (liver transplant recipients or subjects with decompensated liver disease), after 12 weeks of treatment with ledipasvir and sofosbuvir tablets (90 mg/400 mg) and ribavirin, relapse rates were 7% (5/71) and 5% (10/217) in genotype 1 subjects with and without baseline NS5A polymorphisms at resistance-associated positions, respectively.

In the Phase 3 trials and SOLAR trials, the specific baseline NS5A resistance-associated polymorphisms observed among subjects who relapsed were M28T/V, Q30H/R, L31M, H58D/P, and Y93H/N in genotype 1a, and L28M, L31M, A92T, and Y93H in genotype 1b. Subjects with multiple NS5A polymorphisms at resistance-associated positions appeared to have higher relapse rates.

#### *Genotype 4, 5 or 6*

Phylogenetic analysis of HCV sequences from genotype 4-infected subjects in Study 1119 (N=44) and ION-4 (N=8) identified 7 HCV genotype 4 subtypes (4a, 4b, 4d, 4f, 4m, 4o, and 4r). Most subjects were infected with subtype 4a (N=32; 62%) or 4d (N=11; 21%); 1 to 3 subjects were infected with each of the other genotype 4 subtypes. There were 3 subjects with subtype 4r, 2 of whom experienced virologic relapse, and both had a combination of 2 pretreatment NS5A resistance-associated polymorphisms (L28M/V+L30R).

Phylogenetic analysis of HCV sequences from genotype 5-infected subjects in Study 1119 showed almost all were subtype 5a (N=39) with one subject not having a subtype identified at screening or by analysis.

Phylogenetic analysis of HCV sequences from genotype 6-infected subjects in ELECTRON-2 identified 7 HCV genotype 6 subtypes (6a, 6e, 6l, 6m, 6p, 6q, and 6r). Thirty-two percent of the subjects had subtype 6a and 24% had subtype 6e. One to three subjects were infected with the other subtypes 6l, 6m, 6p, 6q, or 6r. The one subject who did not achieve SVR12 had subtype 6l.

Although the data are limited, baseline HCV NS5A resistance-associated polymorphisms are not expected to impact the likelihood of achieving SVR when ledipasvir and sofosbuvir is used as recommended to treat HCV genotype 4, 5, or 6-infected patients, based on the low virologic failure rate observed in Study 1119 and ELECTRON-2. The specific baseline polymorphisms observed in subjects with virologic failure were L28M/V, L30R, and P58T for genotype 4; L31M for genotype 5; and Q24K, F28V, R30A, and T58P for genotype 6.

Relapse occurred in 2 of 3 genotype 4 subjects who had baseline NS5B V321I, a polymorphism at a position associated with treatment failure to sofosbuvir and other nucleoside inhibitors; these two subjects also had baseline NS5A resistance-associated polymorphisms. For genotype 5 and 6, SVR12 was achieved in subjects who had baseline NS5B polymorphisms at positions associated with resistance to sofosbuvir and other nucleoside inhibitors (N=1 with N142T in genotype 5; N=1 with M289I in genotype 5; N=15 with M289L/I in genotype 6). The sofosbuvir resistance-associated substitution S282T was not detected in the baseline NS5B sequence of any subject with genotype 4, 5, or 6 HCV in clinical trials by population or deep nucleotide sequence analysis.

#### *Pediatrics*

In Study 1116, the presence of NS5A and NS5B resistance-associated polymorphisms did not impact treatment outcome; all pediatric subjects 3 years of age and older with baseline NS5A or NS5B nucleoside inhibitor resistance-associated polymorphisms (14%; 32/223) achieved SVR following 12 weeks treatment with ledipasvir and sofosbuvir tablets.

### Cross Resistance

Based on resistance patterns observed in cell culture replicon studies and HCV-infected subjects, cross-resistance between ledipasvir and other NS5A inhibitors is expected. Both sofosbuvir and ledipasvir were fully active against substitutions associated with resistance to other classes of direct-acting antivirals with different mechanisms of action, such as NS5B non-nucleoside inhibitors and NS3 protease inhibitors. The efficacy of ledipasvir/sofosbuvir has not been established in patients who have previously failed treatment with other regimens that include an NS5A inhibitor.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis and Mutagenesis

*Ledipasvir:* Ledipasvir was not genotoxic in a battery of in vitro or in vivo assays, including bacterial mutagenicity, chromosome aberration using human peripheral blood lymphocytes, and in vivo rat micronucleus assays.

Ledipasvir was not carcinogenic in a 6-month rasH2 transgenic mouse study (up to 300 mg/kg/day). Similarly, ledipasvir was not carcinogenic in a 2-year rat study (up to 100 mg/kg/day in males and 30 mg/kg/day in females), resulting in exposures approximately 10 and 4 times, respectively, higher than the exposure in humans at the recommended human dose (RHD).

*Sofosbuvir:* Sofosbuvir was not genotoxic in a battery of in vitro or in vivo assays, including bacterial mutagenicity, chromosome aberration using human peripheral blood lymphocytes, and in vivo mouse micronucleus assays.

Sofosbuvir was not carcinogenic in a 2-year mouse study (up to 200 mg/kg/day in males and 600 mg/kg/day in females) and in a 2-year rat study (up to 750 mg/kg/day), resulting in exposures of the predominant circulating metabolite GS-331007 of approximately 4 and 18 times (in mice) and 8 and 10 times (in rats), in males and females respectively, the exposure in humans at the RHD.

#### Impairment of Fertility

*Ledipasvir:* Ledipasvir had no adverse effects on mating and fertility. In female rats, the mean number of corpora lutea and implantation sites were reduced slightly at maternal exposures approximately 3 times the exposure in humans at the RHD. At the highest dose levels without effects, exposures of ledipasvir were approximately 5 and 2 times, in males and females, respectively, the exposure in humans at the RHD.

*Sofosbuvir:* Sofosbuvir had no effects on embryo-fetal viability or on fertility when evaluated in rats. At the highest dose tested, exposure to the predominant circulating metabolite GS-331007 was approximately 5 times the exposure in humans at the RHD.

## 14 CLINICAL STUDIES

### 14.1 Description of Clinical Trials

The efficacy and safety of ledipasvir and sofosbuvir were evaluated in four trials in genotype 1 HCV mono-infected subjects including one trial exclusively in treatment-experienced subjects with compensated cirrhosis (Child-Pugh A); one trial in genotype 1 or 4 HCV/HIV-1 coinfecting subjects; two trials in genotype 4, 5, or 6 HCV mono-infected subjects; two trials in genotype 1 or 4 HCV infected pretransplant subjects with decompensated cirrhosis (Child-Pugh B and C) or post-transplant with Metavir F0-F3 fibrosis, compensated cirrhosis, decompensated cirrhosis, or fibrosing cholestatic hepatitis (FCH); two trials in subjects with severe renal impairment (one of which included subjects requiring dialysis); and one trial in genotype 1 or 4 HCV pediatric subjects 3 years of age and older without cirrhosis or with compensated cirrhosis, as summarized in Table 10 [see *Clinical Studies (14.2, 14.3, 14.4, 14.5, 14.6, and 14.7)*]:

**Table 10 Trials Conducted with Ledipasvir and Sofosbuvir with or without Ribavirin in Subjects with Chronic HCV Genotype 1, 4, 5, or 6 Infection**

<b>Trial</b>	<b>Population</b>	<b>Study Arms (Number of Subjects Treated)</b>
ION-3 * (NCT01851330)	GT1, TN without cirrhosis	Ledipasvir and sofosbuvir 8 weeks (215) Ledipasvir and sofosbuvir + RBV 8 weeks (216) Ledipasvir and sofosbuvir 12 weeks (216)
ION-1 * (NCT01701401)	GT1, TN with or without cirrhosis	Ledipasvir and sofosbuvir 12 weeks (214) Ledipasvir and sofosbuvir + RBV 12 weeks (217) Ledipasvir and sofosbuvir 24 weeks (217) Ledipasvir and sofosbuvir + RBV 24 weeks (217)
ION-2 * (NCT01768286)	GT1, TE <sup>†</sup> with or without cirrhosis	Ledipasvir and sofosbuvir 12 weeks (109) Ledipasvir and sofosbuvir + RBV 12 weeks (111) Ledipasvir and sofosbuvir 24 weeks (109) Ledipasvir and sofosbuvir + RBV 24 weeks (111)
SIRIUS ‡ (NCT01965535)	GT1, TE <sup>†</sup> with cirrhosis	Ledipasvir and sofosbuvir + RBV 12 Weeks (77) Ledipasvir and sofosbuvir 24 weeks (77)
ION-4 * (NCT02073656)	GT1 and GT4 HCV/HIV-1 coinfecting TN and TE <sup>†</sup> with or without cirrhosis	Ledipasvir and sofosbuvir 12 Weeks (N=327 for GT1; N=8 for GT4)
1119 * (NCT02081079)	GT4 and GT5, TN and TE <sup>†</sup> with or without cirrhosis	Ledipasvir and sofosbuvir 12 Weeks (N=44 for GT4; N=41 for GT5)
ELECTRON-2 * (NCT01826981)	GT6, TN and TE <sup>†</sup> with or without cirrhosis	Ledipasvir and sofosbuvir 12 Weeks (25)
SOLAR-1 * and SOLAR-2 *	GT1 and GT4 pre-transplant with decompensated cirrhosis or post-transplant	Ledipasvir and sofosbuvir + RBV 12

SOLAR-2 (NCT01938430 and NCT02010255)	Cirrhosis of post-transplant with Metavir F0–F3 fibrosis, compensated cirrhosis, decompensated cirrhosis, or FCH	Weeks (336) Ledipasvir and sofosbuvir + RBV 24 weeks (334)
1116 * (NCT02249182)	GT1 or 4 TN and TE with or without cirrhosis in pediatric subjects 3 years of age and older	Ledipasvir and sofosbuvir 12 Weeks (223) Ledipasvir and sofosbuvir 24 Weeks (1)
0154 * (NCT01958281)	GT1 TN and TE <sup>†</sup> with severe RI without dialysis	Ledipasvir and sofosbuvir 12 weeks (18)
4063 * (NCT03036839)	GT1, 5, or 6 TN and TE <sup>§</sup> with or without compensated cirrhosis, with ESRD requiring dialysis	Ledipasvir and sofosbuvir 8 Weeks (45) Ledipasvir and sofosbuvir 12 Weeks (12) Ledipasvir and sofosbuvir 24 Weeks (6)

ESRD = End stage renal disease; RBV = ribavirin; RI = Renal impairment; TN = Treatment-naïve subjects

\* Open-label.

† TE = Treatment-experienced subjects including those who have failed a peginterferon alfa + RBV based regimen with or without an HCV protease inhibitor.

‡ Double-blind, placebo-controlled.

§ TE = Treatment experienced subjects including those who have failed either interferon/peginterferon alfa/ribavirin based regimens or HCV-specific direct-acting antiviral regimens that do not include an NS5A polymerase inhibitor.

Ledipasvir and sofosbuvir was administered once daily by mouth in these trials. For subjects without cirrhosis or with compensated cirrhosis who received ribavirin, the ribavirin dosage was 1000 mg per day for subjects weighing less than 75 kg or 1200 mg per day for subjects weighing at least 75 kg. For subjects with decompensated cirrhosis in SOLAR-1 and SOLAR-2 studies, the starting ribavirin dosage was 600 mg per day regardless of transplantation status. Ribavirin dose adjustments were performed according to the ribavirin labeling.

Serum HCV RNA values were measured during the clinical trials using the COBAS TaqMan HCV test (version 2.0), for use with the High Pure System in ION-3, ION-1, ION-2, SIRIUS, and ION-4 studies or the COBAS AmpliPrep/COBAS Taqman HCV test (version 2.0) in ELECTRON-2, 1119, SOLAR-1, SOLAR-2, and 1116 studies. The COBAS TaqMan HCV test (version 2.0) for use with the High Pure System has a lower limit of quantification (LLOQ) of 25 IU per mL and the COBAS AmpliPrep/COBAS Taqman HCV test (version 2.0) has a LLOQ of 15 IU per mL. Sustained virologic response (SVR12), defined as HCV RNA less than LLOQ at 12 weeks after the cessation of treatment, was the primary endpoint in studies in adults and the key efficacy endpoint in the study in pediatric subjects 12 years of age and older. Relapse was a secondary endpoint, which was defined as HCV RNA greater than or equal to LLOQ with 2 consecutive values or last available post-treatment measurement during the post-treatment period after achieving HCV RNA less than LLOQ at end of treatment.

## 14.2 Clinical Trials in Subjects with Genotype 1 HCV

### Treatment-Naïve Adults without Cirrhosis – ION-3 (Study 0108)

ION-3 was a randomized, open-label trial in treatment-naïve non-cirrhotic subjects with genotype 1 HCV. Subjects were randomized in a 1:1:1 ratio to one of the following three

treatment groups and stratified by HCV genotype (1a vs 1b): ledipasvir and sofosbuvir tablets (90 mg/400 mg) for 8 weeks, ledipasvir and sofosbuvir tablets (90 mg/400 mg) for 12 weeks, or ledipasvir and sofosbuvir tablets (90 mg/400 mg) + ribavirin for 8 weeks.

Demographics and baseline characteristics were balanced across the treatment groups. Of the 647 treated subjects, the median age was 55 years (range: 20 to 75); 58% of the subjects were male; 78% were White; 19% were Black; 6% were Hispanic or Latino; mean body mass index was 28 kg/m<sup>2</sup> (range: 18 to 56 kg/m<sup>2</sup>); 81% had baseline HCV RNA levels greater than or equal to 800,000 IU per mL; 80% had genotype 1a HCV infection; 73% had non-C/C IL28B alleles (CT or TT).

Table 11 presents the SVR12 for the ledipasvir and sofosbuvir treatment groups in the ION-3 trial after 8 and 12 weeks of ledipasvir and sofosbuvir treatment. Ribavirin was not shown to increase the SVR12 observed with ledipasvir and sofosbuvir. Therefore, the ledipasvir and sofosbuvir + ribavirin arm is not presented in Table 11.

**Table 11 Study ION-3: SVR12 after 8 and 12 Weeks of Treatment in Treatment-Naïve Non-Cirrhotic Subjects with Genotype 1 HCV**

	<b>Ledipasvir and Sofosbuvir 8 Weeks (N=215)</b>	<b>Ledipasvir and Sofosbuvir 12 Weeks (N=216)</b>
SVR12	94% (202/215)	96% (208/216)
Outcome for Subjects without SVR		
On-Treatment Virologic Failure	0/215	0/216
Relapse*	5% (11/215)	1% (3/216)
Other†	1% (2/215)	2% (5/216)
SVR by Genotype‡		
Genotype 1a	93% (159/171)	96% (165/172)
Genotype 1b	98% (42/43)	98% (43/44)

\* The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

† Other includes subjects who did not achieve SVR and did not meet virologic failure criteria (e.g., lost to follow-up).

‡ One subject without a confirmed subtype for genotype 1 infection was excluded from this subgroup analysis.

The treatment difference between the 8-week treatment of ledipasvir and sofosbuvir and 12-week treatment of ledipasvir and sofosbuvir was -2.3% (97.5% confidence interval - 7.2% to 2.5%). Among subjects with a baseline HCV RNA less than 6 million IU per mL, the SVR12 was 97% (119/123) with 8-week treatment of ledipasvir and sofosbuvir and 96% (126/131) with 12-week treatment of ledipasvir and sofosbuvir.

Relapse rates by baseline viral load are presented in Table 12.

**Table 12 Study ION-3: Relapse Rates by Baseline Viral Load after 8 and 12 Weeks of Treatment in Treatment-Naïve Non-Cirrhotic Subjects with Genotype 1 HCV**

	<b>Ledipasvir and Sofosbuvir 8 Weeks (N=215)</b>	<b>Ledipasvir and Sofosbuvir 12 Weeks (N=216)</b>
Number of Responders at End of Treatment	215	216
Baseline HCV RNA*		
HCV RNA <6 million IU/mL	2% (2/123)	2% (2/131)
HCV RNA ≥6 million IU/mL	10% (9/92)	1% (1/85)

\* HCV RNA values were determined using the Roche TaqMan Assay; a subject's HCV RNA may vary from visit to visit.

### Treatment-Naïve Adults with or without Cirrhosis – ION-1 (Study 0102)

ION-1 was a randomized, open-label trial that evaluated 12 and 24 weeks of treatment with ledipasvir and sofosbuvir tablets (90 mg/400 mg) with or without ribavirin in 865 treatment-naïve subjects with genotype 1 HCV including those with cirrhosis. Subjects were randomized in a 1:1:1:1 ratio to receive ledipasvir and sofosbuvir for 12 weeks, ledipasvir and sofosbuvir + ribavirin for 12 weeks, ledipasvir and sofosbuvir for 24 weeks, or ledipasvir and sofosbuvir + ribavirin for 24 weeks. Randomization was stratified by the presence or absence of cirrhosis and HCV genotype (1a vs 1b).

Demographics and baseline characteristics were balanced across the treatment groups. Of the 865 treated subjects, the median age was 54 years (range: 18 to 80); 59% of the subjects were male; 85% were White; 12% were Black; 12% were Hispanic or Latino; mean body mass index was 27 kg/m<sup>2</sup> (range: 18 to 48 kg/m<sup>2</sup>); 79% had baseline HCV RNA levels greater than or equal to 800,000 IU per mL; 67% had genotype 1a HCV infection; 70% had non-C/C IL28B alleles (CT or TT); and 16% had cirrhosis.

Table 13 presents the SVR12 for the treatment group of ledipasvir and sofosbuvir for 12 weeks in the ION-1 trial. Ribavirin was not shown to increase SVR12 observed with ledipasvir and sofosbuvir. Therefore, the ledipasvir and sofosbuvir + ribavirin arm is not presented in Table 13.

**Table 13 Study ION-1: SVR12 after 12 Weeks of Treatment in Treatment-Naïve Subjects with Genotype 1 HCV with and without Cirrhosis**

	<b>Ledipasvir and Sofosbuvir 12 Weeks (N=214)</b>
SVR12*	99% (210/213)
Outcome for Subjects without SVR	
On-Treatment Virologic Failure*	0/213
Relapse*,†	<1% (1/212)
Other*,‡	1% (2/213)

\* Excluding one subject with genotype 4 infection.

† The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

‡ Other includes subjects who did not achieve SVR12 and did not meet virologic failure criteria (e.g., lost to follow-up).

SVR12 for selected subgroups are presented in Table 14.

**Table 14 Study ION-1: SVR12 for Selected Subgroups after 12 Weeks of Treatment in Treatment-Naïve Subjects with Genotype 1 HCV with and without Cirrhosis**

	<b>Ledipasvir and Sofosbuvir 12 Weeks (N=214)</b>
Genotype*	
Genotype 1a	98% (142/145)
Genotype 1b	100% (67/67)
Cirrhosis†	
No	99% (176/177)
Yes	94% (32/34)

\* One subject without a confirmed subtype for genotype 1 infection and one subject with genotype 4 infection were excluded from this subgroup analysis.

† Subjects with missing cirrhosis status were excluded from this subgroup analysis.

#### Previously-Treated Adults with or without Cirrhosis – ION-2 (Study 0109)

ION-2 was a randomized, open-label trial that evaluated 12 and 24 weeks of treatment with ledipasvir and sofosbuvir tablets (90 mg/400 mg) with or without ribavirin in genotype 1 HCV-infected subjects with or without cirrhosis who failed prior therapy with an interferon-based regimen, including regimens containing an HCV protease inhibitor. Subjects were randomized in a 1:1:1:1 ratio to receive ledipasvir and sofosbuvir for 12 weeks, ledipasvir and sofosbuvir + ribavirin for 12 weeks, ledipasvir and sofosbuvir for 24 weeks, or ledipasvir and sofosbuvir + ribavirin for 24 weeks. Randomization was stratified by the presence or absence of cirrhosis, HCV genotype (1a vs 1b) and response to prior HCV therapy (relapse/breakthrough vs nonresponse).

Demographics and baseline characteristics were balanced across the treatment groups. Of the 440 treated subjects, the median age was 57 years (range: 24 to 75); 65% of the subjects were male; 81% were White; 18% were Black; 9% were Hispanic or Latino; mean body mass index was 28 kg/m<sup>2</sup> (range: 19 to 50 kg/m<sup>2</sup>); 89% had baseline HCV RNA levels greater than or equal to 800,000 IU per mL; 79% had genotype 1a HCV infection; 88% had non-C/C IL28B alleles (CT or TT); and 20% had cirrhosis. Forty-seven percent (47%) of the subjects failed a prior therapy of pegylated interferon and ribavirin. Among these subjects, 49% were relapse/breakthrough and 51% were non-responder. Fifty-three percent (53%) of the subjects failed a prior therapy of pegylated interferon and ribavirin with an HCV protease inhibitor. Among these subjects, 62% were relapse/breakthrough and 38% were non-responder.

Table 15 presents the SVR12 for the ledipasvir and sofosbuvir treatment groups in the ION-2 trial. Ribavirin was not shown to increase SVR12 observed with ledipasvir and sofosbuvir. Therefore, the ledipasvir and sofosbuvir + ribavirin arms are not presented in Table 15.

**Table 15 Study ION-2: SVR12 after 12 and 24 Weeks of Treatment in Subjects with Genotype 1 HCV with or without Cirrhosis Who Failed Prior Therapy**

	<b>Ledipasvir and Sofosbuvir 12 Weeks (N=109)</b>	<b>Ledipasvir and Sofosbuvir 24 Weeks (N=109)</b>
SVR12	94% (102/109)	99% (108/109)
Outcome for Subjects without SVR		
On-Treatment Virologic Failure	0/109	0/109
Relapse*	6% (7/108)	0/109
Other†	0/109	1% (1/109)

\* The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

† Other includes subjects who did not achieve SVR12 and did not meet virologic failure criteria (e.g., lost to follow-up).

Among the subjects with available SVR12 and SVR24 data (206/218), all subjects who achieved SVR12 in the ION-2 study also achieved SVR24.

SVR12 and relapse rates for selected subgroups are presented in Tables 16 and 17.

**Table 16 Study ION-2: SVR12 for Selected Subgroups after 12 and 24 Weeks of Treatment in Subjects with Genotype 1 HCV Who Failed Prior Therapy**

	<b>Ledipasvir and Sofosbuvir 12 Weeks (N=109)</b>	<b>Ledipasvir and Sofosbuvir 24 Weeks (N=109)</b>
Genotype		
Genotype 1a	95% (82/86)	99% (84/85)
Genotype 1b	87% (20/23)	100% (24/24)
Cirrhosis*		
No	95% (83/87)	99% (85/86)
Yes	86% (19/22)	100% (22/22)
Prior HCV Therapy		
Peg-IFN + RBV	93% (40/43)	100% (58/58)
HCV protease inhibitor + Peg-IFN + RBV	94% (62/66)	98% (49/50)
Response to Prior HCV Therapy		
Relapse/Breakthrough	95% (57/60)	100% (60/60)
Non-responder	92% (45/49)	98% (48/49)

RBV = ribavirin.

\* Subjects with missing cirrhosis status were excluded from this subgroup analysis.

**Table 17 Study ION-2: Relapse Rates for Selected Subgroups after 12 and 24 Weeks of Treatment in Subjects with Genotype 1 HCV Who Failed Prior Therapy**

	<b>Ledipasvir and Sofosbuvir 12 Weeks (N=109)</b>	<b>Ledipasvir and Sofosbuvir 24 Weeks (N=109)</b>
Number of Responders at End of Treatment	108	109
Cirrhosis*		
No	5% (4/86) <sup>†</sup>	0% (0/86)
Yes	14% (3/22)	0% (0/22)
Presence of Baseline NS5A Resistance-Associated Polymorphisms <sup>‡</sup>		
No	2% (2/85)	0% (0/90)
Yes	22% (5/23)	0% (0/19)
IL28B Status		
C/C	0% (0/10)	0% (0/16)
Non-C/C	7% (7/98)	0% (0/93)

\* Subjects with missing cirrhosis status were excluded from this subgroup analysis.

† These 4 non-cirrhotic relapsers all had baseline NS5A resistance-associated polymorphisms.

‡ NS5A resistance-associated polymorphisms include any change at NS5A positions 24, 28, 30, 31, 58, 92, or 93.

#### Previously-Treated Adults with Cirrhosis – SIRIUS (Study 0121)

SIRIUS was a randomized, double-blind and placebo-controlled trial that evaluated the efficacy of ledipasvir and sofosbuvir tablets (90 mg/400 mg) + ribavirin for 12 weeks or ledipasvir and sofosbuvir tablets (90 mg/400 mg) without ribavirin for 24 weeks in genotype 1 HCV-infected subjects with compensated cirrhosis who failed prior therapy with a Peg-IFN + ribavirin regimen followed by a subsequent Peg-IFN + ribavirin + an HCV protease inhibitor regimen. Subjects were randomized in a 1:1 ratio to receive placebo for 12 weeks followed by ledipasvir and sofosbuvir + ribavirin for 12 weeks or ledipasvir and sofosbuvir for 24 weeks. Randomization was stratified by HCV genotype (1a vs 1b) and response to prior HCV therapy (never achieved HCV RNA less than LLOQ vs achieved HCV RNA less than LLOQ).

Demographics and baseline characteristics were balanced across the treatment groups. Of the 155 randomized subjects, the median age was 56 years (range: 23 to 77); 74% of the subjects were male; 97% were White; mean body mass index was 27 kg/m<sup>2</sup> (range: 19 to 47 kg/m<sup>2</sup>); 63% had genotype 1a HCV infection; 94% had non-C/C IL28B alleles (CT or TT). One subject discontinued therapy while on placebo, and was not included in the efficacy analysis.

The SVR12 was 96% (74/77) and 97% (75/77) in subjects treated with ledipasvir and sofosbuvir + ribavirin for 12 weeks and ledipasvir and sofosbuvir for 24 weeks without ribavirin, respectively. All 5 subjects who did not achieve SVR12 relapsed.

### **14.3 Clinical Trials in Subjects with Genotype 4, 5, or 6 HCV**

Below are trial descriptions, SVR12 and relapse data in the genotype 4, 5, and 6 HCV populations. Trial results in the genotype 4, 5, and 6 HCV populations are based upon limited number of subjects in some subgroups, particularly in subjects who have been previously treated and subjects with cirrhosis.

#### Genotype 4

In two open-label studies (Study 1119 and ION-4), ledipasvir and sofosbuvir tablets (90 mg/400 mg) were administered for 12 weeks to treatment-naïve and previously-treated adult subjects with genotype 4 HCV infection. Study 1119 enrolled 44 treatment-naïve or previously-treated subjects with genotype 4 HCV, with or without cirrhosis. ION-4 enrolled 4 treatment-naïve and 4 previously-treated subjects with genotype 4 HCV infection who were coinfecting with HIV-1, none of whom had cirrhosis.

In Study 1119, the overall SVR12 rate was 93% (41/44). SVR12 was similar based upon prior HCV treatment history and cirrhosis status. In ION-4, all 8 subjects achieved SVR12.

#### Genotype 5

In the open-label 1119 trial, ledipasvir and sofosbuvir tablets (90 mg/400 mg) were administered for 12 weeks to 41 treatment-naïve or previously treated adult subjects with genotype 5 HCV infection, with or without cirrhosis. The overall SVR12 was 93% (38/41). SVR12 was similar based upon prior HCV treatment history and cirrhosis status.

#### Genotype 6

In the open-label ELECTRON-2 trial, ledipasvir and sofosbuvir tablets (90 mg/400 mg) were administered for 12 weeks to 25 treatment-naïve or previously treated adult subjects with genotype 6 HCV infection, with or without cirrhosis. The overall SVR12 was 96% (24/25). SVR12 was similar based upon prior HCV treatment history and cirrhosis status. The single subject who relapsed discontinued study treatment early (at approximately Week 8).

### **14.4 Clinical Trials in Subjects Coinfected with HCV and HIV-1**

ION-4 was an open-label clinical trial that evaluated the safety and efficacy of 12 weeks of treatment with ledipasvir and sofosbuvir tablets (90 mg/400 mg) without ribavirin in HCV treatment-naïve and previously treated adult subjects with genotype 1 or 4 HCV infection who were coinfecting with HIV-1. Treatment-experienced subjects had failed prior treatment with Peg-IFN + ribavirin, Peg-IFN + ribavirin + an HCV protease inhibitor, or sofosbuvir + ribavirin. Subjects were on a stable HIV-1 antiretroviral therapy that included emtricitabine + tenofovir disoproxil fumarate, administered with efavirenz, rilpivirine, or raltegravir.

Of the 335 treated subjects, the median age was 52 years (range: 26 to 72); 82% of the subjects were male; 61% were White; 34% were Black; mean body mass index was 27 kg/m<sup>2</sup> (range: 18 to 66 kg/m<sup>2</sup>); 75% had genotype 1a HCV infection; 2% had genotype 4 infection; 76% had non-C/C IL28B alleles (CT or TT); and 20% had compensated

cirrhosis. Fifty-five percent (55%) of the subjects were treatment-experienced.

Table 18 presents the SVR12 in the ION-4 trial after 12 weeks of ledipasvir and sofosbuvir treatment.

**Table 18 Study ION-4: SVR12 in Subjects with Genotype 1 or 4 HCV Coinfected with HIV-1**

	<b>Ledipasvir and Sofosbuvir 12 Weeks (N=335)</b>
SVR12	96% (321/335)
Outcome for Subjects without SVR	
On-Treatment Virologic Failure	<1% (2/335)
Relapse*	3% (10/333)
Other†	<1% (2/335)

\* The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

† Other includes subjects who did not achieve SVR12 and did not meet virologic failure criteria (e.g., lost to follow-up).

SVR12 rates were 94% (63/67) in subjects with cirrhosis and 98% (46/47) in subjects who were previously treated and had cirrhosis. The relapse rate in the ION-4 trial in Black subjects was 9% (10/115), all of whom were IL28B non-CC genotype, and none in non-Black subjects (0/220). In the ION-1, ION-2, and ION-3 HCV mono-infection studies, relapse rates were 3% (10/305) in Black subjects and 2% (26/1637) in non-Black subjects.

No subject had HIV-1 rebound during the study. The percentage of CD4+ cells did not change during treatment. Median CD4+ cell count increase of 29 cells/mm<sup>3</sup> was observed at the end of treatment with ledipasvir and sofosbuvir tablets for 12 weeks.

#### **14.5 Clinical Trials in Liver Transplant Recipients and/or Subjects with Decompensated Cirrhosis**

SOLAR-1 and SOLAR-2 were two open-label trials that evaluated 12 and 24 weeks of treatment with ledipasvir and sofosbuvir tablets (90 mg/400 mg) in combination with ribavirin in HCV treatment-naïve and previously treated adult subjects with genotype 1 and 4 infection who had undergone liver transplantation and/or who had decompensated liver disease. The two trials were identical in study design. Subjects were enrolled in one of the seven groups in the trials based on liver transplantation status and severity of hepatic impairment (see Table 19). Subjects with a CPT score greater than 12 were excluded. Within each group, subjects were randomized in a 1:1 ratio to receive ledipasvir and sofosbuvir + ribavirin for 12 weeks or ledipasvir and sofosbuvir + ribavirin for 24 weeks. For subjects with decompensated cirrhosis in SOLAR-1 and SOLAR-2 studies, the starting ribavirin dosage was 600 mg per day regardless of transplantation status. Ribavirin dose adjustments were performed according to the ribavirin labeling [see *Clinical Studies (14.1)*].

Demographics and baseline characteristics were balanced across the treatment groups. Of the 670 treated subjects, the median age was 59 years (range: 21 to 81); 77% of the

subjects were male; 91% were White; mean body mass index was 28 kg/m<sup>2</sup> (range: 18 to 49 kg/m<sup>2</sup>); 94% and 6% had genotype 1 and 4 HCV infection, respectively; 78% of the subjects failed a prior HCV therapy.

Table 19 presents the pooled SVR12 rates for SOLAR-1 and SOLAR-2 in subjects with genotype 1 HCV treated with ledipasvir and sofosbuvir + ribavirin for 12 weeks. The SVR12 rates observed with 24 weeks of ledipasvir and sofosbuvir + ribavirin were similar to the SVR12 rates observed with 12 weeks of treatment. Therefore, the results for the ledipasvir and sofosbuvir + ribavirin 24 weeks arm are not presented in Table 19.

**Table 19 Studies SOLAR-1 and SOLAR-2: SVR12 and Relapse Rates After 12 Weeks of Treatment with Ledipasvir and Sofosbuvir and Ribavirin in Subjects with Genotype 1 HCV Who Were Post Liver Transplant and/or Who Had Decompensated Liver Disease**

	<b>Ledipasvir and Sofosbuvir + RBV 12 weeks (N=307)</b>	
	<b>SVR12 (N=300)*,†</b>	<b>Relapse (N=288)*,†,‡</b>
Pre-transplant		
CPT B	87% (45/52)	12% (6/51)
CPT C	88% (35/40)	5% (2/37)
Post-transplant		
Metavir score F0–F3	95% (94/99)	3% (3/97)
CPT A	98% (55/56)	0% (0/55)
CPT B	89% (41/46)	2% (1/42)
CPT C	57% (4/7)	33% (2/6)

\* Five subjects transplanted prior to post-treatment Week 12 with HCV RNA < LLOQ at last measurement prior to transplant were excluded.

† Two subjects were excluded due to failure to meet the inclusion criteria for any of the treatment groups (i.e., did not have decompensated cirrhosis and had also not received a liver transplant).

‡ Twelve subjects were excluded from relapse analysis because they died (N=11) or withdrew consent (N=1) prior to reaching the 12-week post-treatment follow-up visit.

There were 7 subjects with fibrosing cholestatic hepatitis in the 12-week treatment arm, and all subjects achieved SVR12.

In genotype 4 HCV post-transplant subjects without cirrhosis or with compensated cirrhosis treated with ledipasvir and sofosbuvir + ribavirin for 12 weeks (N=12), the SVR12 rate was similar to rates reported with genotype 1; no subjects relapsed. Available data in subjects with genotype 4 HCV who had decompensated cirrhosis (pre- and post-liver transplantation) were insufficient for dosing recommendations; therefore, these results are not presented.

#### **14.6 Clinical Trials in Adults with Severe Renal Impairment, Including those Requiring Dialysis**

Trial 0154 was an open-label clinical trial that evaluated 12 weeks of treatment with

ledipasvir and sofosbuvir tablets (90 mg/400 mg) in 18 treatment-naïve and treatment-experienced (subjects with prior exposure to an HCV NS5B polymerase inhibitor were excluded) genotype 1 HCV-infected adults with severe renal impairment not requiring dialysis. At baseline, two subjects (11%) had cirrhosis and the mean eGFR was 24.9 mL/min (range: 9.0 to 39.6). The SVR rate was 100% (18/18).

As shown in the table below, Trial 4063 was an open-label three-arm clinical trial that evaluated 8, 12, and 24 weeks of treatment with ledipasvir and sofosbuvir in a total of 63 adults with chronic HCV infection and ESRD requiring dialysis. Of the 63 subjects, 10% had cirrhosis, 24% were treatment-experienced, 95% were on hemodialysis, and 5% were on peritoneal dialysis; mean duration on dialysis was 12 years (range: 0.2 to 43 years). The SVR rates for the 8, 12, and 24 week ledipasvir and sofosbuvir treatment groups are shown in Table 20.

**Table 20 Trial 4063: SVR12 after 8, 12, and 24 Weeks of Treatment in Adults with HCV with or without Cirrhosis and with Severe Renal Impairment Requiring Dialysis**

	<b>Ledipasvir and sofosbuvir 8 Weeks (N=45)</b>	<b>Ledipasvir and sofosbuvir 12 Weeks (N=12)</b>	<b>Ledipasvir and sofosbuvir 24 Weeks (N=6)</b>
Population	Treatment-naïve, GT 1 HCV Non-cirrhotic	Treatment-naïve and treatment-experienced* GT 1, 5, 6† HCV Non-cirrhotic	Treatment-experienced, GT 1 HCV with compensated cirrhosis
SVR12	93% (42/45)	100% (12/12)	83% (5/6)
Outcome for Subjects without SVR			
On-Treatment Virologic Failure	0/45	0/12	0/6
Relapse	0/44	0/12	0/6
Other‡	7% (3/45)	0/12	17% (1/6)

\* Subjects with prior exposure to any HCV NS5A inhibitor were excluded.

† One subject had an indeterminate HCV GT.

‡ "Other" outcomes includes subjects who did not achieve SVR and did not meet virologic failure criteria. All subjects who failed without virologic relapse or on-treatment virologic failure died prior to follow-up Week 12. None of these deaths were assessed as treatment-related.

#### 14.7 Clinical Trial in Pediatric Subjects

The efficacy of ledipasvir and sofosbuvir was evaluated in an open-label trial (Study 1116) in 224 HCV treatment-naïve (N=186) and treatment-experienced (N=38) pediatric subjects 3 years of age or older. This study evaluated 12 weeks of treatment with ledipasvir and sofosbuvir once daily in genotype 1 (N=183) or genotype 4 (N=3) treatment-naïve subjects without cirrhosis or with compensated cirrhosis; genotype 1

treatment-experienced subjects without cirrhosis (N=37); and evaluated 24 weeks of treatment with ledipasvir and sofosbuvir once daily in one genotype 1 subject who was both treatment-experienced and cirrhotic.

*Subjects 12 Years to <18 Years of Age:* Ledipasvir and sofosbuvir was evaluated in 100 subjects 12 years to <18 years of age with HCV genotype 1 infection. Demographics and baseline characteristics were balanced across treatment-naïve and treatment-experienced subjects (patients had failed an interferon based regimen with or without ribavirin). The median age was 15 years (range: 12 to 17); 63% of the subjects were female; 91% were White, 7% were Black, and 2% were Asian; 13% were Hispanic/Latino; mean body mass index was 23 kg/m<sup>2</sup> (range: 13.1 to 36.6 kg/m<sup>2</sup>); mean weight was 61 kg (range 33 to 126 kg); 55% had baseline HCV RNA levels greater than or equal to 800,000 IU/mL; 81% had genotype 1a HCV infection. One subject (treatment-naïve) had known compensated cirrhosis. The majority of subjects (84%) had been infected through vertical transmission.

The SVR12 rate was 98% overall (98% [78/80] in treatment-naïve subjects and 100% [20/20] in treatment-experienced subjects). No subject experienced on-treatment virologic failure or relapse. Two subjects were lost to follow-up.

*Subjects 6 Years to <12 Years of Age:* Ledipasvir and sofosbuvir was evaluated in 90 subjects 6 years to <12 years of age with HCV genotype 1 or 4 infection. Among these subjects, 72 (80%) were treatment-naïve and 18 (20%) were treatment-experienced. Eighty-nine of the subjects (87 with genotype 1 HCV infection and 2 with genotype 4 HCV infection) were treated with ledipasvir and sofosbuvir for 12 weeks, 1 subject with genotype 1 HCV infection was treated with ledipasvir and sofosbuvir for 24 weeks. The median age was 9 years (range: 6 to 11); 59% of the subjects were male; 79% were White, 8% were Black, and 6% were Asian; 10% were Hispanic/Latino; mean body mass index was 18 kg/m<sup>2</sup> (range: 13 to 31kg/m<sup>2</sup>); mean weight was 33 kg (range 18 to 76 kg); 59% had baseline HCV RNA levels greater than or equal to 800,000 IU/mL; 86% had genotype 1a HCV infection; 2 subjects (1 treatment-naïve, 1 treatment-experienced) had known compensated cirrhosis. The majority of subjects (97%) had been infected through vertical transmission.

The SVR12 rate was 99% (86/87) in subjects with genotype 1 HCV infection, and 100% (2/2) in subjects with genotype 4 HCV infection. The one genotype 1 subject treated with ledipasvir and sofosbuvir for 24 weeks also achieved SVR12. The one subject (genotype 1) who did not achieve SVR12 and relapsed had been treated with ledipasvir and sofosbuvir for 12 weeks.

*Subjects 3 Years to <6 Years of Age:* Ledipasvir and sofosbuvir was evaluated in 34 subjects 3 years to <6 years of age with HCV genotype 1 (N = 33) or genotype 4 (N = 1) infection. All of the subjects were treatment-naïve and treated with ledipasvir and sofosbuvir for 12 weeks. The median age was 5 years (range: 3 to 5); 71% of the subjects were female; 79% were White, 3% were Black, and 6% were Asian; 18% were Hispanic/Latino; mean body mass index was 17 kg/m<sup>2</sup> (range: 13 to 25 kg/m<sup>2</sup>); mean weight was 19 kg (range 11 to 34 kg); 56% had baseline HCV RNA levels greater than or equal to 800,000 IU/mL; 82% had genotype 1a HCV infection; no subjects had known cirrhosis. All subjects (100%) had been infected through vertical transmission.

The SVR12 rate was 97% (32/33) in subjects with genotype 1 HCV infection, and the one subject with genotype 4 HCV infection also achieved SVR12. One subject prematurely discontinued study treatment due to an adverse event.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Ledipasvir and sofosbuvir tablets (90 mg/400 mg) are orange, diamond-shaped, film-coated, debossed with "ASE" on one side and "9875" on the other side of the tablet. Each carton contains 28 tablets (2 blister cards each containing 14 tablets) (NDC 72626-2601-1).

Store at room temperature below 30 °C (86 °F).

## **17 PATIENT COUNSELING INFORMATION**

Advise the patient to read the FDA-approved patient labeling (Patient Information and Instructions for Use).

### Risk of Hepatitis B Virus Reactivation in Patients Coinfected with HCV and HBV

Inform patients that HBV reactivation can occur in patients coinfecting with HBV during or after treatment of HCV infection. Advise patients to tell their healthcare provider if they have a history of HBV infection [see *Warnings and Precautions (5.1)*].

### Serious Symptomatic Bradycardia When Coadministered with Amiodarone

Advise patients to seek medical evaluation immediately for symptoms of bradycardia such as near-fainting or fainting, dizziness or lightheadedness, malaise, weakness, excessive tiredness, shortness of breath, chest pain, confusion, or memory problems [see *Warnings and Precautions (5.2)*, *Adverse Reactions (6.2)*, and *Drug Interactions (7.2)*].

### Drug Interactions

Inform patients that ledipasvir and sofosbuvir may interact with other drugs. Advise patients to report to their healthcare provider the use of any other prescription or nonprescription medication or herbal products including St. John's wort [see *Warnings and Precautions (5.2, 5.3)* and *Drug Interactions (7)*].

### Pregnancy

Advise patients to avoid pregnancy during combination treatment with ledipasvir and sofosbuvir and ribavirin and for 6 months after completion of treatment. Inform patients to notify their healthcare provider immediately in the event of a pregnancy [see *Use in Specific Populations (8.1)*].

### Hepatitis C Virus Transmission

Inform patients that the effect of treatment of hepatitis C infection on transmission is not known, and that appropriate precautions to prevent transmission of the hepatitis C virus during treatment or in the event of treatment failure should be taken.

### Administration

Advise patients to take ledipasvir and sofosbuvir every day at the regularly scheduled time with or without food. Inform patients that it is important not to miss or skip doses and to take ledipasvir and sofosbuvir for the duration that is recommended by the physician.

For ledipasvir and sofosbuvir (HARVONI) oral pellets, advise patients or caregivers to read and follow the Instructions for Use for preparing the correct dose.

Manufactured for:  
Asegua Therapeutics LLC  
An affiliate of Gilead Sciences, Inc.  
Foster City, CA 94404

Asegua is a trademark of Asegua Therapeutics LLC. All other trademarks referenced herein are the property of their respective owners.

205834-AG-012

**Patient Information**  
**Ledipasvir and Sofosbuvir**  
**(led" i pas' vir and soe fos' bue vir)**  
**tablets (90 mg/400 mg)**  
**Authorized generic of HARVONI®**

**Important: If you take ledipasvir and sofosbuvir with ribavirin, you should also read the Medication Guide for ribavirin.**

**What is the most important information I should know about ledipasvir and sofosbuvir?**

Ledipasvir and sofosbuvir can cause serious side effects, including, **Hepatitis B virus reactivation:** Before starting treatment with ledipasvir and sofosbuvir, your healthcare provider will do blood tests to check for hepatitis B virus infection. If you have ever had hepatitis B virus infection, the hepatitis B virus could become active again during or after treatment of hepatitis C virus with ledipasvir and sofosbuvir. Hepatitis B virus becoming active again (called reactivation) may cause serious liver problems including liver failure and death. Your healthcare provider will monitor you if you are at risk for hepatitis B virus reactivation during treatment and after you stop taking ledipasvir and sofosbuvir.

**For more information about side effects, see the section "What are the possible side effects of ledipasvir and sofosbuvir?"**

**What is ledipasvir and sofosbuvir?**

Ledipasvir and sofosbuvir is a prescription medicine used to treat adults and children 3 years of age and older with chronic (lasting a long time) hepatitis C virus (HCV):

- genotype 1, 4, 5, or 6 infection without cirrhosis or with compensated cirrhosis
- genotype 1 infection with advanced cirrhosis (decompensated) in combination with ribavirin
- genotype 1 or 4 infection without cirrhosis or with compensated cirrhosis who have had a liver transplant, in combination with ribavirin

It is not known if ledipasvir and sofosbuvir is safe and effective in children with HCV under 3 years of age.

**Before taking ledipasvir and sofosbuvir, tell your healthcare provider about all of your medical conditions, including if you:**

- have ever had hepatitis B virus infection
- have liver problems other than hepatitis C infection
- have had a liver transplant
- have kidney problems or you are on dialysis
- have HIV infection
- are pregnant or plan to become pregnant. It is not known if ledipasvir and sofosbuvir tablets will harm your unborn baby.

- **Males and females who take ledipasvir and sofosbuvir in combination with ribavirin should also read the ribavirin Medication Guide for important pregnancy, contraception, and infertility information.**
- are breastfeeding or plan to breastfeed. It is not known if ledipasvir and sofosbuvir pass into your breast milk.
  - Talk to your healthcare provider about the best way to feed your baby during treatment with ledipasvir and sofosbuvir.

**Tell your healthcare provider about all of the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Ledipasvir and sofosbuvir and other medicines may affect each other. This can cause you to have too much or not enough ledipasvir and sofosbuvir or other medicines in your body. This may affect the way ledipasvir and sofosbuvir or your other medicines work, or may cause side effects. **Keep a list of your medicines to show your healthcare provider and pharmacist.**

- You can ask your healthcare provider or pharmacist for a list of medicines that interact with ledipasvir and sofosbuvir.

**Do not start taking a new medicine without telling your healthcare provider.** Your healthcare provider can tell you if it is safe to take ledipasvir and sofosbuvir with other medicines.

#### **How should I take ledipasvir and sofosbuvir?**

- Take ledipasvir and sofosbuvir exactly as your healthcare provider tells you to take it. Do not change your dose unless your healthcare provider tells you to.
- Do not stop taking ledipasvir and sofosbuvir without first talking with your healthcare provider.
- Take ledipasvir and sofosbuvir by mouth, with or without food.
- It is important that you do not miss or skip doses of ledipasvir and sofosbuvir during treatment.
- For adults the usual dose of ledipasvir and sofosbuvir is one 90 mg/400 mg tablet each day.
- For children 3 years of age and older your healthcare provider will prescribe the right dose of ledipasvir and sofosbuvir based on your child's body weight.
  - Tell your healthcare provider if your child has problems with swallowing tablets.
  - If your healthcare provider prescribes ledipasvir and sofosbuvir (HARVONI) oral pellets for your child, see "How should I give ledipasvir and sofosbuvir (HARVONI) oral pellets to my child."
- Do not miss a dose of ledipasvir and sofosbuvir. Missing a dose lowers the amount of medicine in your blood. Refill your ledipasvir and sofosbuvir prescription before you run out of medicine.

If you take too much ledipasvir and sofosbuvir, call your healthcare provider or go to the nearest hospital emergency room right away.

#### **How should I give ledipasvir and sofosbuvir (HARVONI) oral pellets to my child?**

**See the detailed Instructions for Use for information about how to give or take a dose of ledipasvir and sofosbuvir (HARVONI) oral pellets.**

- Administer ledipasvir and sofosbuvir (HARVONI) oral pellets exactly as instructed by your healthcare provider.
- Do not open the packet until ready to use.

- Hold the ledipasvir and sofosbuvir (HARVONI) oral pellets packet with the cut line on top.
- Shake the ledipasvir and sofosbuvir (HARVONI) oral pellets packet gently to settle the pellets.
- Tear or cut the ledipasvir and sofosbuvir (HARVONI) oral pellets packet along the cut line.
- Ledipasvir and sofosbuvir (HARVONI) oral pellets can be taken right in the mouth without chewing, or with food.
- If ledipasvir and sofosbuvir (HARVONI) oral pellets are taken with food, sprinkle the pellets on one or more spoonfuls of non-acidic soft food at or below room temperature. Examples of non-acidic foods include pudding, chocolate syrup, mashed potato, and ice cream. Take ledipasvir and sofosbuvir (HARVONI) oral pellets within 30 minutes of gently mixing with food and swallow the entire contents without chewing to avoid a bitter taste.
- Do not store any leftover ledipasvir and sofosbuvir (HARVONI) mixture (oral pellets mixed with food) for use at a later time. Throw away any unused portion.

### **What are the possible side effects of ledipasvir and sofosbuvir?**

#### **Ledipasvir and sofosbuvir can cause serious side effects, including:**

- **Hepatitis B virus reactivation.** See "What is the most important information I should know about ledipasvir and sofosbuvir?"
- **Slow heart rate (bradycardia).** Ledipasvir and sofosbuvir treatment may result in slowing of the heart rate along with other symptoms when taken with amiodarone (Cordarone<sup>®</sup>, Nexterone<sup>®</sup>, Pacerone<sup>®</sup>), a medicine used to treat certain heart problems. In some cases bradycardia has led to death or the need for a heart pacemaker when amiodarone is taken with ledipasvir and sofosbuvir tablets. Get medical help right away if you take amiodarone with ledipasvir and sofosbuvir tablets and get any of the following symptoms:
  - fainting or near-fainting
  - dizziness or lightheadedness
  - not feeling well
  - weakness
  - extreme tiredness
  - shortness of breath
  - chest pains
  - confusion
  - memory problems

The most common side effects of ledipasvir and sofosbuvir include:

- tiredness
- headache
- weakness

These are not all the possible side effects of ledipasvir and sofosbuvir. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store ledipasvir and sofosbuvir tablets?**

- Store ledipasvir and sofosbuvir tablets at room temperature below 86°F (30°C).

### **Keep ledipasvir and sofosbuvir tablets and all medicines out of the reach of children.**

### **General information about the safe and effective use of ledipasvir and sofosbuvir**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use ledipasvir and sofosbuvir for a condition for which it was

not prescribed. Do not give ledipasvir and sofosbuvir to other people, even if they have the same symptoms you have. It may harm them.  
You can ask your healthcare provider or pharmacist for information about ledipasvir and sofosbuvir that is written for health professionals.

**What are the ingredients in ledipasvir and sofosbuvir tablets (90 mg/400 mg)?**

**Active ingredients:** ledipasvir and sofosbuvir

**Inactive ingredients:** colloidal silicon dioxide, copovidone, croscarmellose sodium, lactose monohydrate, magnesium stearate, and microcrystalline cellulose.

The tablet film-coat contains: FD&C yellow #6/sunset yellow FCF aluminum lake, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide.

Manufactured for:

Asegua Therapeutics LLC, an affiliate of Gilead Sciences, Inc., Foster City, CA 94404

For more information, call 1-800-445-3235 or go to [www.asegua.com](http://www.asegua.com).

Asegua is a trademark of Asegua Therapeutics LLC. All other trademarks referenced herein are the property of their respective owners.

205834-AG-012

This Patient Information has been approved by the U.S. Food and Drug Administration

Revised: 03/2020

**PRINCIPAL DISPLAY PANEL - 14 Tablet Blister Pack Card Carton**

NDC 72626-2601-1

ASEGUA™  
THERAPEUTICS

Ledipasvir and Sofosbuvir  
90 mg / 400 mg  
tablets

Take 1 tablet once daily

Rx only

Note to pharmacist:  
Do not cover ALERT box with pharmacy label.

ALERT: Find out about medicines that should NOT be taken with Ledipasvir and Sofosbuvir

Contents:  
Two blister cards  
Each blister card contains 14 tablets.

Authorized generic of Harvoni®

NDC 72626-2601-1

**ASEGUA™**  
THERAPEUTICS

# Ledipasvir and Sofosbuvir 90 mg / 400 mg tablets

Take 1 tablet once daily

**Rx only**

Note to pharmacist:  
Do not cover ALERT box with pharmacy label.

**ALERT: Find out about medicines that should  
NOT be taken with Ledipasvir and Sofosbuvir**

Contents:  
Two blister cards  
Each blister card contains 14 tablets.  
Authorized generic of Harvoni®

## LEDIPASVIR AND SOFOSBUVIR

ledipasvir and sofosbuvir tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:72626-2601
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LEDIPASVIR</b> (UNII: 013TE6E4WW) (LEDIPASVIR - UNII:013TE6E4WW)	LEDIPASVIR	90 mg
<b>SOFOSBUVIR</b> (UNII: WJ6CA3ZU8B) (SOFOSBUVIR - UNII:WJ6CA3ZU8B)	SOFOSBUVIR	400 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>COPOVIDONE K25-31</b> (UNII: D9C330MD8B)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	

**Product Characteristics**

<b>Color</b>	ORANGE	<b>Score</b>	no score
<b>Shape</b>	DIAMOND	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	ASE;9875
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72626-2601-1	2 in 1 CARTON	01/01/2019	
1		14 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC	NDA205834	01/01/2019	

**Labeler** - Asegu Therapeutics LLC (116670954)

Revised: 1/2025

Asegu Therapeutics LLC