

**GREEN GUARD SUPER STAT BLOOD CLOTTING FIRST AID- benzethonium chloride and lidocaine aerosol, spray**  
**Unifirst First Aid Corporation**

-----  
**Green Guard Super Stat Blood Clotter Spray**

***Active Ingredients***

Benzethonium Chloride 0.2% w/w

Lidocaine 4% w/w

***Purpose***

Topical Antiseptic

Topical Anesthetic

***Uses***

- for help on control of superficial bleeding
- for temporary relief of pain and to help protect against infection in minor cuts, scrapes and minor burns
- minor burns

***Warnings***

**For external use only.**

**Flammable**

- keep away from flame
- contents under pressure
- do not puncture or incinerate container

**Do not use**

- in or near eyes or other mucous membranes
- in case of serious burns
- in case of deep or puncture wounds
- for a prolonged period of time
- on large portions of the body
- on raw surfaces
- on blistered areas

**Stop use and ask a doctor if**

- condition worsens or symptoms persist for more than 7 days

- condition clears up and recurs within a few days
- redness, swelling, or irritation occurs

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

- clean the infected area
- shake can well before using
- hold 6-8 inches from moist injured area
- spray until area is covered and superficial bleeding has stopped
- scab-like cover will gradually disappear or may be removed with soap and water
- for adult institutional use only
- not intended for use on children under 12 years of age

***Other Information***

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating and inhaling may be harmful or fatal
- store at room temperature 59-86°F (15-30°C)
- do not expose to temperatures above 120°F

***Inactive Ingredients***

isobutane, isopropyl myristate, karaya, n-butane, propane, silica, sorbitan monooleate, talc, tragacanth

**Green Guard Super Stat Blood Clotter Spray**

Super Stat

Blood Clotter Spray

- Aids in control of superficial bleeding
- Temporarily relieves pain of minor cuts, burns and scrapes
- Helps prevent infection
- Contains no fluorocarbons

Green Guard®

Aerosol

WARNING: FLAMMABLE

Contents under Pressure, Read Back Panel Carefully

Net Wt. 3 oz. (85g)

### Drug Facts (continued)

#### Directions

- clean the affected area
- shake can well before using
- hold 6-8 inches from surface
- spray until area is covered and superficial bleeding is stopped
- scab-like cover will gradually disappear or may be removed with soap and water
- for adult institutional use only
- not intended for use on children under 12 years of age

#### Other information

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal
- store at room temperature 59-86°F (15-30°C)
- do not expose to temperatures above 120° F

#### Inactive ingredients

isobutane, isopropyl myristate, karaya, n-butane, propane, silica, sorbitan monooleate, talc, tragacanth.

Questions or comments? 1-800-869-6970

Distributed by:  
GREEN GUARD®  
St. Louis, MO 63045  
Reorder #3003

Rev. 7/25/18



470341

# Super Stat

## Blood Clotter Spray

- Aids in control of superficial bleeding
- Temporarily relieves pain of minor cuts, burns and scrapes
- Helps prevent infection
- Contains no fluorocarbons



Aerosol

WARNING: FLAMMABLE  
CONTENTS UNDER PRESSURE, READ BACK PANEL CAREFULLY  
Net Wt. 3 oz. (85 g)

### Drug Facts

Active ingredients	Purposes
Benzethonium Chloride 0.2% w/w.....	Topical antiseptic
Lidocaine 4% w/w.....	Topical anesthetic

#### Uses

- for help in control of superficial bleeding
- for temporary relief of pain and to help protect against infection in minor cuts, scrapes and minor burns

#### Warnings

For external use only.

#### Flammable

- keep away from fire or flame
- contents under pressure
- do not puncture or incinerate container

#### Do not use

- in or near eyes or other mucous membranes
- in case of serious burns
- in case of deep or puncture wounds
- for a prolonged period of time
- on a large portion of the body
- on raw surfaces
- on blistered areas

#### Stop use and ask a doctor if

- condition worsens or symptoms persist for more than 7 days
- condition clears up and recurs within a few days
- redness, swelling, or irritation occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

## GREEN GUARD SUPER STAT BLOOD CLOTTING FIRST AID

benzethonium chloride and lidocaine aerosol, spray

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-300
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	3400 mg in 85 g
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	170 mg in 85 g

### Inactive Ingredients

Ingredient Name	Strength
ISOBUTANE (UNII: BXR49TP611)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
KARAYA GUM (UNII: 73W9IQY50Q)	
BUTANE (UNII: 6LV4FOR43R)	
PROPANE (UNII: T75W9911L6)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
TALC (UNII: 7SEV7J4R1U)	

TRAGACANTH (UNII: 2944357020)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-300-17	85 g in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	04/05/2013	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	04/05/2013	

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 9/2025

Unifirst First Aid Corporation