

LLORENS CARE TUSSIN DM- dextromethorphan hbr, guaifenesin liquid
Llorens Pharmaceutical International Division, Inc.

LC Tussin DM 833

In each 5 mL

Dextromethorphan - 10 mg

Guaifenesin - 100 mg

Purpose

Cough Suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus)
- helps thin bronchial secretions to make coughs more productive

Warnings

Do not use if you are now taking a prescription monamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Do not use if you have ever had an allergic reaction to any of the ingredients in this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or head that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions:

- take every 4 hours as needed, or as directed by a doctor
- do not take more than 6 doses in 24 hours
- do not exceed recommended dose

Age	Dose
-----	------

Adults and children over 12 years and over	2 teaspoonfuls (10 mL) every 4 hours
Children under 12 years	ask a doctor

Inactive ingredients artificial and natural cherry flavor, citric acid, FD&C #40, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose, sucrose.

Questions or comments? 1-866-595-5598

Llorens Pharmaceutical

Llorens Care

Compare to Robitussin® DM Active Ingredients*

NDC: 54859-833-04

Drug Facts

Active Ingredients (in each 5 mL teaspoonful) **Purposes**
Dextromethorphan HBr, USP 10 mg..... Cough Suppressant
Guaifenesin, USP 100 mg Expectorant

USES • temporarily relieves cough due to minor throat and bronchial irritation • helps loosen phlegm (mucus) • helps thin bronchial secretions to make coughs more productive

WARNINGS
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Do not use if you have ever had an allergic reaction to any of the ingredients in this product.

Ask a doctor before use if you have • a cough that occurs with too much phlegm (mucus) • a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Stop use and ask a doctor if cough lasts for more than 7 days, comes back or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

adult

tussin DM

Dextromethorphan HBr - Cough Suppressant
Guaifenesin - Expectorant

Cough & Chest Congestion DM
Controls Cough
Relieves Chest Congestion
Thins & Loosens Mucus

Cherry Flavor

4 FL OZ (118 mL)

Drug Facts (Continued)

Directions • Take every 4 hours as needed, or as directed by a doctor
• Do not take more than 6 doses in a 24-hour period
• Do not exceed recommended dose

adults and children 12 years and over	2 teaspoonfuls (10 mL) every 4 hours
children under 12 years	ask a doctor

Other Information
• **TAMPER-EVIDENT:** Do not use if foil seal over bottle opening is torn, broken, or missing • store at room temperature 15°-30°C (59°-86°F) • protect from freezing • do not refrigerate • protect from light • each teaspoonful (5 mL) contains: sodium 2 mg

Inactive Ingredients: Artificial and natural cherry flavor, citric acid, FD&C red #40, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose, sucrose

Questions or Comments? 1-866-595-5598

*This product is not manufactured or distributed by the owner of the registered trademark Robitussin® DM.

Manufactured by:
Llorens Pharmaceutical International Division, Inc.
7080 NW 37th Ct, Miami, FL 33147

Code #: L-190
REV: 06/22

www.LlorensPharm.com

Lot #:

Exp Date:

3 54859 83304 4

LLORENS CARE TUSSIN DM

dextromethorphan hbr, guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54859-833
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

MENTHOL (UNII: L7T10EIP3A)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54859-833-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2025	

Labeler - Llorens Pharmaceutical International Division, Inc. (037342305)