

**GERI CARE TUSSIN CF MAX- acetaminophen, dextromethorphan hbr,
guaifenesin liquid
GERI-CARE PHARMACEUTICAL CORP**

GeriCare TussinCFMax 596

Active Ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Purpose

Pain reliever/fever reducer

Cough Suppressant

Expectorant

Uses

- temporarily relieves these symptoms occurring with a cold or flu:
- cough due to minor throat and bronchial irritation
- minor aches and pains
- sore throat
- headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in any 24-hour period, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking any other pain reliever/fever reducer

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- mL = milliliter
- do not take more than 6 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage
- measure only with dosing cup provided
- keep dosing cup with product
- **adults and children 12 years and over: 20 mL every 4 hours**
- **children under 12 years of age: do not use**

Inactive ingredients: artificial and natural flavor, citric acid, FD&C #40, hydroxyethyl cellulose, menthol, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium citrate, sucralose

Questions or comments: 1-800-540-3765

TAMPER-EVIDENT: Do not use if foil seal over bottle opening is torn, broken or missing.

Drug Facts

Active ingredients (in each 20 mL) **Purposes**
Acetaminophen 650 mg.....Pain reliever/Fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant

Uses • temporarily relieves these symptoms occurring with a cold or flu:
• cough due to minor throat and bronchial irritation
• minor aches and pains • sore throat • headache
• temporarily reduces fever • helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 doses in any 24-hour period, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have • liver disease • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema.

Ask a doctor or pharmacist before use if you are • taking the blood thinning drug warfarin • taking any other pain reliever/fever reducer ►

GERI CARE®

NDC 57896-596-08

TUSSIN CF MAX

Adult - For Ages 12 & Older

MAXIMUM STRENGTH

ACETAMINOPHEN - Pain Reliever/Fever Reducer
DEXTRMETHORPHAN HBr - Cough Suppressant
GUAIFENESIN - Expectorant

MULTI-SYMPOTOM RELIEF

- Cough, Sore Throat
- Head & Body Aches, Fever
- Chest Congestion
- Thins & Loosens Mucus

Compare to the active ingredients in Robitussin CF®
MAX SEVERE MULTI-SYMPOTOM Cough, Cold + Flu*

8 FL OZ (237 mL)

Drug Facts (continued)

Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days
• redness or swelling is present • new symptoms occur
• cough comes back or occurs with rash or headache that lasts
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions • mL = milliliter
• do not take more than 6 doses in any 24-hour period
• do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
• measure only with dosing cup provided • keep dosing cup with product
• **adults and children 12 years and over:** 20 mL every 4 hours
• **children under 12 years of age:** do not use

Other information • each 20 mL contains: sodium 9 mg
• store at 20°C-25°C (68°F-77°F); excursions permitted between 15°C-30°C (59°F-86°F)

Inactive ingredients: artificial and natural flavor, citric acid, FD&C Red #40, hydroxyethyl cellulose, menthol, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium citrate, sucralose.

Questions or comments: 1-800-540-3765

*This product is not manufactured or distributed by the owner of the registered trademark: Robitussin CF® Max, Severe Multi-Symptom Cough, Cold + Flu. Maximum Strength claim based on maximum levels of active ingredients per 4 hour dose.

Code # L-260 REV 11/25

Dist. By: Geri-Care
Pharmaceuticals Corp.
1295 Towbin Avenue
Lakewood, NJ 08701

Lot#
Exp. Date:

3 57896 59608

GERI CARE TUSSIN CF MAX

acetaminophen, dextromethorphan hbr, guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-596
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KOOR)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

MENTHOL (UNII: L7T10EIP3A)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-596-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2025	

Labeler - GERI-CARE PHARMACEUTICAL CORP (611196254)

Revised: 12/2025

GERI-CARE PHARMACEUTICAL CORP