

CARBOPLATIN- carboplatin injection, solution
Hospira, Inc.

Carboplatin Injection

R_x only

WARNING

Carboplatin Injection should be administered under the supervision of a qualified physician experienced in the use of cancer chemotherapeutic agents. Appropriate management of therapy and complications is possible only when adequate treatment facilities are readily available.


Bone marrow suppression is dose related and may be severe, resulting in infection and/or bleeding. Anemia may be cumulative and may require transfusion support. Vomiting is another frequent drug-related side effect.

Anaphylactic-like reactions to Carboplatin Injection have been reported and may occur within minutes of Carboplatin Injection administration. Epinephrine, corticosteroids, and antihistamines have been employed to alleviate symptoms.

DESCRIPTION

Carboplatin Injection is supplied as a sterile, pyrogen-free, aqueous solution available in 50 mg/5 mL, 150 mg/15 mL, 450 mg/45 mL or 600 mg/60 mL multiple-dose vials containing 10 mg/mL of carboplatin for administration by intravenous infusion. Each mL contains 10 mg carboplatin and Water for Injection, USP.

Carboplatin is a platinum coordination compound. The chemical name for carboplatin is platinum, diammine [1,1-cyclobutane-dicarboxylato(2-)-0,0']-, (SP-4-2), and carboplatin has the following structural formula:



Carboplatin is a crystalline powder with the molecular formula of C₆H₁₂N₂O₄Pt and a molecular weight of 371.25. It is soluble in water at a rate of approximately 14 mg/mL,

and the pH of a 1% solution is 5 to 7. It is virtually insoluble in ethanol, acetone, and dimethylacetamide.

CLINICAL PHARMACOLOGY

Carboplatin, like cisplatin, produces predominantly interstrand DNA cross-links rather than DNA-protein cross-links. This effect is apparently cell-cycle nonspecific. The aquation of carboplatin, which is thought to produce the active species, occurs at a slower rate than in the case of cisplatin. Despite this difference, it appears that both carboplatin and cisplatin induce equal numbers of drug-DNA cross-links, causing equivalent lesions and biological effects. The differences in potencies for carboplatin and cisplatin appear to be directly related to the difference in aquation rates.

In patients with creatinine clearances of about 60 mL/min or greater, plasma levels of intact carboplatin decay in a biphasic manner after a 30-minute intravenous infusion of 300 mg/m² to 500 mg/m² of carboplatin. The initial plasma half-life (alpha) was found to be 1.1 to 2 hours (n=6), and the postdistribution plasma half-life (beta) was found to be 2.6 to 5.9 hours (n=6). The total body clearance, apparent volume of distribution and mean residence time for carboplatin are 4.4 L/hour, 16 L and 3.5 hours, respectively. The C_{max} values and areas under the plasma concentration versus time curves from 0 to infinity (AUC inf) increase linearly with dose, although the increase was slightly more than dose proportional. Carboplatin, therefore, exhibits linear pharmacokinetics over the dosing range studied (300 mg/m² to 500 mg/m²).

Carboplatin is not bound to plasma proteins. No significant quantities of protein-free, ultrafilterable platinum-containing species other than carboplatin are present in plasma. However, platinum from carboplatin becomes irreversibly bound to plasma proteins and is slowly eliminated with a minimum half-life of 5 days.

The major route of elimination of carboplatin is renal excretion. Patients with creatinine clearances of approximately 60 mL/min or greater excrete 65% of the dose in the urine within 12 hours and 71% of the dose within 24 hours. All of the platinum in the 24-hour urine is present as carboplatin. Only 3% to 5% of the administered platinum is excreted in the urine between 24 and 96 hours. There are insufficient data to determine whether biliary excretion occurs.

In patients with creatinine clearances below 60 mL/min, the total body and renal clearances of carboplatin decrease as the creatinine clearance decreases. Carboplatin Injection dosages should therefore be reduced in these patients (see **DOSAGE AND ADMINISTRATION**).

The primary determinant of Carboplatin Injection clearance is glomerular filtration rate (GFR) and this parameter of renal function is often decreased in elderly patients. Dosing formulas incorporating estimates of GFR (see **DOSAGE AND ADMINISTRATION**) to provide predictable Carboplatin Injection plasma AUCs should be used in elderly patients to minimize the risk of toxicity.

CLINICAL STUDIES

Use with Cyclophosphamide for Initial Treatment of Ovarian Cancer

In two prospectively randomized, controlled studies conducted by the National Cancer

Institute of Canada, Clinical Trials Group (NCIC), and the Southwest Oncology Group (SWOG), 789 chemotherapy naive patients with advanced ovarian cancer were treated with carboplatin or cisplatin, both in combination with cyclophosphamide every 28 days for six courses before surgical reevaluation. The following results were obtained from both studies:

Comparative Efficacy

Overview of Pivotal Trials

| | NCIC | SWOG |
|---|-----------------------|-----------------------|
| Number of patients randomized | 447 | 342 |
| Median age (years) | 60 | 62 |
| Dose of cisplatin | 75 mg/m ² | 100 mg/m ² |
| Dose of carboplatin | 300 mg/m ² | 300 mg/m ² |
| Dose of cyclophosphamide | 600 mg/m ² | 600 mg/m ² |
| Residual tumor <2 cm (number of patients) | 39% (174/447) | 14% (49/342) |

Clinical Response in Measurable Disease Patients

| | NCIC | SWOG |
|--|-----------------|----------------|
| Carboplatin (number of patients) | 60% (48/80) | 58% (48/83) |
| Cisplatin (number of patients) | 58% (49/85) | 43% (33/76) |
| 95% C.I. of difference (Carboplatin - Cisplatin) | (-13.9%, 18.6%) | (-2.3%, 31.1%) |

Pathologic Complete Response*

| | NCIC | SWOG |
|--|----------------|---------------|
| Carboplatin (number of patients) | 11% (24/224) | 10% (17/171) |
| Cisplatin (number of patients) | 15% (33/223) | 10% (17/171) |
| 95% C.I. of difference (Carboplatin - Cisplatin) | (-10.7%, 2.5%) | (-6.9%, 6.9%) |

* 114 Carboplatin and 109 Cisplatin patients did not undergo second look surgery in NCIC study. 90 Carboplatin and 106 Cisplatin patients did not undergo second look surgery in SWOG study.

Progression-Free Survival (PFS)

| | NCIC | SWOG |
|--|-------------|-------------|
| Median | | |
| Carboplatin | 59 weeks | 49 weeks |
| Cisplatin | 61 weeks | 47 weeks |
| 2-year PFS* | | |
| Carboplatin | 31% | 21% |
| Cisplatin | 31% | 21% |
| 95% C.I. of difference (Carboplatin - Cisplatin) | (-9.3, 8.7) | (-9.0, 9.4) |
| 3-year PFS* | | |

| | | |
|--|--------------|--------------|
| Carboplatin | 19% | 8% |
| Cisplatin | 23% | 14% |
| 95% C.I. of difference (Carboplatin - Cisplatin) | (-11.5, 4.5) | (-14.1, 0.3) |
| Hazard Ratio[†] | 1.10 | 1.02 |
| 95% C.I. (Carboplatin - Cisplatin) | (0.89, 1.35) | (0.81, 1.29) |

* Kaplan-Meier Estimates

Unrelated deaths occurring in the absence of progression were counted as events (progression) in this analysis.

† Analysis adjusted for factors found to be of prognostic significance were consistent with unadjusted analysis.

Survival

| | NCIC | SWOG |
|--|--------------|--------------|
| Median | | |
| Carboplatin | 110 weeks | 86 weeks |
| Cisplatin | 99 weeks | 79 weeks |
| 2-year Survival* | | |
| Carboplatin | 51.9% | 40.2% |
| Cisplatin | 48.4% | 39.0% |
| 95% C.I. of difference (Carboplatin - Cisplatin) | (-6.2, 13.2) | (-9.8, 12.2) |
| 3-year Survival* | | |
| Carboplatin | 34.6% | 18.3% |
| Cisplatin | 33.1% | 24.9% |
| 95% C.I. of difference (Carboplatin - Cisplatin) | (-7.7, 10.7) | (-15.9, 2.7) |
| Hazard Ratio[†] | 0.98 | 1.01 |
| 95% C.I. (Carboplatin - Cisplatin) | (0.78, 1.23) | (0.78, 1.30) |

* Kaplan-Meier Estimates

† Analysis adjusted for factors found to be of prognostic significance were consistent with unadjusted analysis.

Comparative Toxicity

The pattern of toxicity exerted by the carboplatin containing regimen was significantly different from that of the cisplatin-containing combinations. Differences between the two studies may be explained by different cisplatin dosages and by different supportive care.

The carboplatin-containing regimen induced significantly more thrombocytopenia and, in one study, significantly more leukopenia and more need for transfusional support. The cisplatin-containing regimen produced significantly more anemia in one study. However, no significant differences occurred in incidences of infections and hemorrhagic episodes.

Non-hematologic toxicities (emesis, neurotoxicity, ototoxicity, renal toxicity, hypomagnesemia, and alopecia) were significantly more frequent in the cisplatin-containing arms.

ADVERSE EXPERIENCES IN PATIENTS WITH OVARIAN CANCER NCIC STUDY

| | Carboplatin Arm Percent* | Cisplatin Arm Percent* | P-Values† |
|---|-----------------------------|---------------------------|-----------|
| Bone Marrow | | | |
| Thrombocytopenia <100,000/mm ³ | 70 | 29 | <0.001 |
| <50,000/mm ³ | 41 | 6 | <0.001 |
| Neutropenia <2,000 cells/mm ³ | 97 | 96 | ns |
| <1,000 cells/mm ³ | 81 | 79 | ns |
| Leukopenia <4,000 cells/mm ³ | 98 | 97 | ns |
| <2,000 cells/mm ³ | 68 | 52 | 0.001 |
| Anemia <11 g/dL | 91 | 91 | ns |
| <8 g/dL | 18 | 12 | ns |
| Infections | 14 | 12 | ns |
| Bleeding | 10 | 4 | ns |
| Transfusions | 42 | 31 | 0.018 |
| Gastrointestinal | | | |
| Nausea and vomiting | 93 | 98 | 0.010 |
| Vomiting | 84 | 97 | <0.001 |
| Other GI side effects | 50 | 62 | 0.013 |
| Neurologic | | | |
| Peripheral neuropathies | 16 | 42 | <0.001 |
| Ototoxicity | 13 | 33 | <0.001 |
| Other sensory side effects | 6 | 10 | ns |
| Central neurotoxicity | 28 | 40 | 0.009 |
| Renal | | | |
| Serum creatinine elevations | 5 | 13 | 0.006 |
| Blood urea elevations | 17 | 31 | <0.001 |
| Hepatic | | | |
| Bilirubin elevations | 5 | 3 | ns |
| SGOT elevations | 17 | 13 | ns |
| Alkaline phosphatase elevations | - | - | - |
| Electrolytes loss | | | |
| Sodium | 10 | 20 | 0.005 |
| Potassium | 16 | 22 | ns |
| Calcium | 16 | 19 | ns |
| Magnesium | 63 | 88 | <0.001 |
| Other side effects | | | |
| Pain | 36 | 37 | ns |
| Asthenia | 40 | 33 | ns |
| Cardiovascular | 15 | 19 | ns |
| Respiratory | 8 | 9 | ns |

| | | | |
|---------------|----|----|-------|
| Allergic | 12 | 9 | ns |
| Genitourinary | 10 | 10 | ns |
| Alopecia ‡ | 50 | 62 | 0.017 |
| Mucositis | 10 | 9 | ns |

* Values are in percent of evaluable patients

† ns = not significant, p>0.05

‡ May have been affected by cyclophosphamide dosage delivered

ADVERSE EXPERIENCES IN PATIENTS WITH OVARIAN CANCER SWOG STUDY

| | Carboplatin Arm Percent* | Cisplatin Arm Percent* | P-Values† |
|---|-----------------------------|---------------------------|-----------|
| Bone Marrow | | | |
| Thrombocytopenia <100,000/mm ³ | 59 | 35 | <0.001 |
| <50,000/mm ³ | 22 | 11 | 0.006 |
| Neutropenia <2,000 cells/mm ³ | 95 | 97 | ns |
| <1,000 cells/mm ³ | 84 | 78 | ns |
| Leukopenia <4,000 cells/mm ³ | 97 | 97 | ns |
| <2,000 cells/mm ³ | 76 | 67 | ns |
| Anemia <11 g/dL | 88 | 87 | ns |
| <8 g/dL | 8 | 24 | <0.001 |
| Infections | 18 | 21 | ns |
| Bleeding | 6 | 4 | ns |
| Transfusions | 25 | 33 | ns |
| Gastrointestinal | | | |
| Nausea and vomiting | 94 | 96 | ns |
| Vomiting | 82 | 91 | 0.007 |
| Other GI side effects | 40 | 48 | ns |
| Neurologic | | | |
| Peripheral neuropathies | 13 | 28 | 0.001 |
| Ototoxicity | 12 | 30 | <0.001 |
| Other sensory side effects | 4 | 6 | ns |
| Central neurotoxicity | 23 | 29 | ns |
| Renal | | | |
| Serum creatinine elevations | 7 | 38 | <0.001 |
| Blood urea elevations | - | - | - |
| Hepatic | | | |
| Bilirubin elevations | 5 | 3 | ns |
| SGOT elevations | 23 | 16 | ns |
| Alkaline phosphatase elevations | 29 | 20 | ns |
| Electrolytes loss | | | |
| Sodium | - | - | - |
| Potassium | - | - | - |

| | | | |
|---------------------------|----|----|--------|
| Calcium | - | - | - |
| Magnesium | 58 | 77 | <0.001 |
| Other side effects | | | |
| Pain | 54 | 52 | ns |
| Asthenia | 43 | 46 | ns |
| Cardiovascular | 23 | 30 | ns |
| Respiratory | 12 | 11 | ns |
| Allergic | 10 | 11 | ns |
| Genitourinary | 11 | 13 | ns |
| Alopecia † | 43 | 57 | 0.009 |
| Mucositis | 6 | 11 | ns |

* Values are in percent of evaluable patients

† ns = not significant, $p > 0.05$

‡ May have been affected by cyclophosphamide dosage delivered

Use as a Single Agent for Secondary Treatment of Advanced Ovarian Cancer

In two prospective, randomized controlled studies in patients with advanced ovarian cancer previously treated with chemotherapy, carboplatin achieved six clinical complete responses in 47 patients. The duration of these responses ranged from 45 to 71 + weeks.

INDICATIONS

Initial Treatment of Advanced Ovarian Carcinoma

Carboplatin Injection is indicated for the initial treatment of advanced ovarian carcinoma in established combination with other approved chemotherapeutic agents. One established combination regimen consists of Carboplatin Injection and cyclophosphamide. Two randomized controlled studies conducted by the NCIC and SWOG with carboplatin versus cisplatin, both in combination with cyclophosphamide, have demonstrated equivalent overall survival between the two groups (see **CLINICAL STUDIES**).

There is limited statistical power to demonstrate equivalence in overall pathologic complete response rates and long-term survival (≥ 3 years) because of the small number of patients with these outcomes: the small number of patients with residual tumor < 2 cm after initial surgery also limits the statistical power to demonstrate equivalence in this subgroup.

Secondary Treatment of Advanced Ovarian Carcinoma

Carboplatin Injection is indicated for the palliative treatment of patients with ovarian carcinoma recurrent after prior chemotherapy, including patients who have been previously treated with cisplatin.

Within the group of patients previously treated with cisplatin, those who have developed progressive disease while receiving cisplatin therapy may have a decreased response rate.

CONTRAINDICATIONS

Carboplatin Injection is contraindicated in patients with a history of severe allergic reactions to cisplatin or other platinum-containing compounds.

Carboplatin Injection should not be employed in patients with severe bone marrow depression or significant bleeding.

WARNINGS

Bone marrow suppression (leukopenia, neutropenia, and thrombocytopenia) is dose-dependent and is also the dose-limiting toxicity. Peripheral blood counts should be frequently monitored during carboplatin treatment and, when appropriate, until recovery is achieved. Median nadir occurs at day 21 in patients receiving single agent carboplatin. In general, single intermittent courses of carboplatin should not be repeated until leukocyte, neutrophil, and platelet counts have recovered.

Since anemia is cumulative, transfusions may be needed during treatment with carboplatin, particularly in patients receiving prolonged therapy.

Bone marrow suppression is increased in patients who have received prior therapy, especially regimens including cisplatin. Marrow suppression is also increased in patients with impaired kidney function. Initial carboplatin dosages in these patients should be appropriately reduced (see **DOSAGE AND ADMINISTRATION**) and blood counts should be carefully monitored between courses. The use of carboplatin in combination with other bone marrow suppressing therapies must be carefully managed with respect to dosage and timing in order to minimize additive effects.

Hemolytic anemia with the presence of serologic drug-induced antibodies has been reported in patients treated with carboplatin. This event can be fatal.

Hemolytic-uremic syndrome is a potentially life-threatening side effect. Carboplatin should be discontinued at the first sign of microangiopathic hemolytic anemia, such as rapidly falling hemoglobin with concomitant thrombocytopenia or elevation of serum bilirubin, serum creatinine, blood urea nitrogen, or lactate dehydrogenase (LDH). Renal failure may not be reversible with discontinuation of therapy and dialysis may be required (see **ADVERSE REACTIONS**).

Carboplatin has limited nephrotoxic potential, but concomitant treatment with aminoglycosides has resulted in increased renal and/or audiologic toxicity, and caution must be exercised when a patient receives both drugs. Clinically significant hearing loss has been reported to occur in pediatric patients when carboplatin was administered at higher than recommended doses in combination with other ototoxic agents. Delayed onset hearing loss has been reported in pediatric patients. Long-term audiometric follow-up in this population is recommended.

Carboplatin can induce emesis, which can be more severe in patients previously receiving emetogenic therapy. The incidence and intensity of emesis have been reduced by using premedication with antiemetics. Although no conclusive efficacy data exist with the following schedules of carboplatin, lengthening the duration of single intravenous administration to 24 hours or dividing the total dose over five consecutive daily pulse doses has resulted in reduced emesis.

Although peripheral neurotoxicity is infrequent, its incidence is increased in patients

older than 65 years and in patients previously treated with cisplatin. Pre-existing cisplatin-induced neurotoxicity does not worsen in about 70% of the patients receiving carboplatin as secondary treatment.

Loss of vision, which can be complete for light and colors, has been reported after the use of carboplatin with doses higher than those recommended in the package insert. Vision appears to recover totally or to a significant extent within weeks of stopping these high doses.

As in the case of other platinum-coordination compounds, allergic reactions to carboplatin have been reported. These may occur within minutes of administration and should be managed with appropriate supportive therapy. There is increased risk of allergic reactions including anaphylaxis in patients previously exposed to platinum therapy (see **CONTRAINDICATIONS** and **ADVERSE REACTIONS: Allergic Reactions**).

Hypersensitivity reactions which progressed to Kounis syndrome have also been reported (see **ADVERSE REACTIONS: Allergic Reactions**).

Patients at high risk of Tumor Lysis Syndrome (TLS), such as those with high tumor burden, high sensitivity to cytotoxic agents, or tumors that have high proliferative rate, should be monitored closely and appropriate precaution taken.

High dosages of carboplatin (more than four times the recommended dose) have resulted in severe abnormalities of liver function tests. Cases of hepatic veno-occlusive disease (sinusoidal obstructive syndrome) have been reported. Some of them were fatal.

Carboplatin may cause fetal harm when administered to a pregnant woman. Carboplatin has been shown to be embryotoxic and teratogenic in rats. There are no adequate and well-controlled studies in pregnant women. If this drug is used during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant.

PRECAUTIONS

General

Needles or intravenous administration sets containing aluminum parts that may come in contact with Carboplatin Injection should not be used for the preparation or administration of the drug. Aluminum can react with carboplatin causing precipitate formation and loss of potency.

Drug Interactions

The renal effects of nephrotoxic compounds may be potentiated by carboplatin. A decrease in phenytoin serum levels has been observed with concurrent administration of carboplatin and phenytoin/fosphenytoin. This may lead to exacerbation of seizures.

Carcinogenesis, Mutagenesis, Impairment of Fertility

The carcinogenic potential of carboplatin has not been studied, but compounds with

similar mechanisms of action and mutagenicity profiles have been reported to be carcinogenic. Carboplatin has been shown to be mutagenic both *in vitro* and *in vivo*. It has also been shown to be embryotoxic and teratogenic in rats receiving the drug during organogenesis. Secondary malignancies have been reported in association with multi-drug therapy. Acute promyelocytic leukemia (APL) and myelodysplastic syndrome (MDS)/acute myeloid leukemia (AML) have been reported years after therapy with carboplatin and other antineoplastic treatments.

Pregnancy

(see **WARNINGS**).

Nursing Mothers

It is not known whether carboplatin is excreted in human milk. Because there is a possibility of toxicity in nursing infants secondary to carboplatin treatment of the mother, it is recommended that breast-feeding be discontinued if the mother is treated with carboplatin.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established (see **WARNINGS; "Audiologic Toxicity"**).

Geriatric Use

Of the 789 patients in initial treatment combination therapy studies (NCIC and SWOG), 395 patients were treated with carboplatin in combination with cyclophosphamide. Of these, 141 were over 65 years of age and 22 were 75 years or older. In these trials, age was not a prognostic factor for survival. In terms of safety, elderly patients treated with carboplatin were more likely to develop severe thrombocytopenia than younger patients. In a combined database of 1,942 patients (414 were ≥ 65 years of age) that received single agent carboplatin for different tumor types, a similar incidence of adverse events was seen in patients 65 years and older and in patients less than 65. Other reported clinical experience has not identified differences in responses between elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. Because renal function is often decreased in the elderly, renal function should be considered in the selection of carboplatin dosage (see **DOSAGE AND ADMINISTRATION**).

ADVERSE REACTIONS

For a comparison of toxicities when carboplatin or cisplatin was given in combination with cyclophosphamide, see **CLINICAL STUDIES: Use with Cyclophosphamide for Initial Treatment of Ovarian Cancer: Comparative Toxicity**.

ADVERSE EXPERIENCES IN PATIENTS WITH OVARIAN CANCER

| | First Line Combination Therapy* Percent | Second Line Single Agent Therapy† Percent |
|--|---|---|
|--|---|---|

| | | | |
|---------------------------------|------------------------------|----|----|
| Bone Marrow | | | |
| Thrombocytopenia | <100,000/mm ³ | 66 | 62 |
| | <50,000/mm ³ | 33 | 35 |
| Neutropenia | <2,000 cells/mm ³ | 96 | 67 |
| | <1,000 cells/mm ³ | 82 | 21 |
| Leukopenia | <4,000 cells/mm ³ | 97 | 85 |
| | <2,000 cells/mm ³ | 71 | 26 |
| Anemia | <11 g/dL | 90 | 90 |
| | <8 g/dL | 14 | 21 |
| Infections | | 16 | 5 |
| Bleeding | | 8 | 5 |
| Transfusions | | 35 | 44 |
| Gastrointestinal | | | |
| Nausea and vomiting | | 93 | 92 |
| Vomiting | | 83 | 81 |
| Other GI side effects | | 46 | 21 |
| Neurologic | | | |
| Peripheral neuropathies | | 15 | 6 |
| Ototoxicity | | 12 | 1 |
| Other sensory side effects | | 5 | 1 |
| Central neurotoxicity | | 26 | 5 |
| Renal | | | |
| Serum creatinine elevations | | 6 | 10 |
| Blood urea elevations | | 17 | 22 |
| Hepatic | | | |
| Bilirubin elevations | | 5 | 5 |
| SGOT elevations | | 20 | 19 |
| Alkaline phosphatase elevations | | 29 | 37 |
| Electrolytes loss | | | |
| Sodium | | 10 | 47 |
| Potassium | | 16 | 28 |
| Calcium | | 16 | 31 |
| Magnesium | | 61 | 43 |
| Other side effects | | | |
| Pain | | 44 | 23 |
| Asthenia | | 41 | 11 |
| Cardiovascular | | 19 | 6 |
| Respiratory | | 10 | 6 |
| Allergic | | 11 | 2 |
| Genitourinary | | 10 | 2 |
| Alopecia | | 49 | 2 |
| Mucositis | | 8 | 1 |

* **Use with Cyclophosphamide for Initial Treatment of Ovarian Cancer:** Data are based on the experience of 393 patients with ovarian cancer (regardless of baseline status) who received initial combination therapy with carboplatin and cyclophosphamide in two randomized controlled studies conducted by SWOG and NCIC (see **CLINICAL STUDIES**).

Combination with cyclophosphamide as well as duration of treatment may be responsible for the differences that can be noted in the adverse experience table.

† **Single Agent Use for the Secondary Treatment of Ovarian Cancer:** Data are based on the experience of 553 patients with previously treated ovarian carcinoma (regardless of baseline status) who received single agent carboplatin.

In the narrative section that follows, the incidences of adverse events are based on data from 1,893 patients with various types of tumors who received carboplatin as single agent therapy.

Hematologic Toxicity

Bone marrow suppression is the dose-limiting toxicity of carboplatin. Thrombocytopenia with platelet counts below 50,000/mm³ occurs in 25% of the patients (35% of pretreated ovarian cancer patients); neutropenia with granulocyte counts below 1,000/mm³ occurs in 16% of the patients (21% of pretreated ovarian cancer patients); leukopenia with WBC counts below 2,000/mm³ occurs in 15% of the patients (26% of pretreated ovarian cancer patients). The nadir usually occurs about day 21 in patients receiving single agent therapy. By day 28, 90% of patients have platelet counts above 100,000/mm³; 74% have neutrophil counts above 2,000/mm³; 67% have leukocyte counts above 4,000/mm³.

Marrow suppression is usually more severe in patients with impaired kidney function. Patients with poor performance status have also experienced a higher incidence of severe leukopenia and thrombocytopenia.

The hematologic effects, although usually reversible, have resulted in infectious or hemorrhagic complications in 5% of the patients treated with carboplatin, with drug-related death occurring in less than 1% of the patients. Fever has also been reported in patients with neutropenia.

Anemia with hemoglobin less than 11 g/dL has been observed in 71% of the patients who started therapy with a baseline above that value. The incidence of anemia increases with increasing exposure to carboplatin. Transfusions have been administered to 26% of the patients treated with carboplatin (44% of previously treated ovarian cancer patients).

Bone marrow depression may be more severe when carboplatin is combined with other bone marrow suppressing drugs or with radiotherapy.

Gastrointestinal Toxicity

Vomiting occurs in 65% of the patients (81% of previously treated ovarian cancer patients) and in about one-third of these patients it is severe. Carboplatin, as a single agent or in combination, is significantly less emetogenic than cisplatin; however, patients previously treated with emetogenic agents, especially cisplatin, appear to be more prone to vomiting. Nausea alone occurs in an additional 10% to 15% of patients. Both nausea and vomiting usually cease within 24 hours of treatment and are often responsive to antiemetic measures. Although no conclusive efficacy data exist with the following schedules, prolonged administration of carboplatin, either by continuous 24-hour infusion or by daily pulse doses given for five consecutive days, was associated with less severe vomiting than the single-dose intermittent schedule. Emesis was increased when carboplatin was used in combination with other emetogenic compounds. Other gastrointestinal effects observed frequently were pain, in 17% of the patients; diarrhea,

in 6%; and constipation, also in 6%.

Neurologic Toxicity

Peripheral neuropathies have been observed in 4% of the patients receiving carboplatin (6% of pretreated ovarian cancer patients) with mild paresthesias occurring most frequently. Carboplatin therapy produces significantly fewer and less severe neurologic side effects than does therapy with cisplatin. However, patients older than 65 years and/or previously treated with cisplatin appear to have an increased risk (10%) for peripheral neuropathies. In 70% of the patients with pre-existing cisplatin-induced peripheral neurotoxicity, there was no worsening of symptoms during therapy with carboplatin. Clinical ototoxicity and other sensory abnormalities such as visual disturbances and change in taste have been reported in only 1% of the patients. Central nervous system symptoms have been reported in 5% of the patients and appear to be most often related to the use of antiemetics.

Although the overall incidence of peripheral neurologic side effects induced by carboplatin is low, prolonged treatment, particularly in cisplatin pretreated patients, may result in cumulative neurotoxicity.

Nephrotoxicity

Development of abnormal renal function test results is uncommon, despite the fact that carboplatin, unlike cisplatin, has usually been administered without high-volume fluid hydration and/or forced diuresis. The incidences of abnormal renal function tests reported are 6% for serum creatinine and 14% for blood urea nitrogen (10% and 22%, respectively, in pretreated ovarian cancer patients). Most of these reported abnormalities have been mild and about one-half of them were reversible.

Creatinine clearance has proven to be the most sensitive measure of kidney function in patients receiving carboplatin, and it appears to be the most useful test for correlating drug clearance and bone marrow suppression. Twenty-seven percent of the patients who had a baseline value of 60 mL/min or more demonstrated a reduction below this value during carboplatin therapy.

Hepatic Toxicity

The incidences of abnormal liver function tests in patients with normal baseline values were reported as follows: total bilirubin, 5%; SGOT, 15%; and alkaline phosphatase, 24%; (5%, 19%, and 37%, respectively, in pretreated ovarian cancer patients). These abnormalities have generally been mild and reversible in about one-half of the cases, although the role of metastatic tumor in the liver may complicate the assessment in many patients. In a limited series of patients receiving very high dosages of carboplatin and autologous bone marrow transplantation, severe abnormalities of liver function tests were reported.

Electrolyte Changes

The incidences of abnormally decreased serum electrolyte values reported were as follows: sodium, 29%; potassium, 20%; calcium, 22%; and magnesium, 29%; (47%, 28%, 31%, and 43%, respectively, in pretreated ovarian cancer patients). Electrolyte supplementation was not routinely administered concomitantly with carboplatin, and these electrolyte abnormalities were rarely associated with symptoms.

Allergic Reactions

Hypersensitivity to carboplatin has been reported in 2% of the patients. These allergic reactions have been similar in nature and severity to those reported with other platinum-containing compounds, i.e., rash, urticaria, erythema, pruritus, and rarely bronchospasm and hypotension. Anaphylactic reactions have been reported as part of post-marketing surveillance (see **WARNINGS**). These reactions have been successfully managed with standard epinephrine, corticosteroid, and antihistamine therapy.

There have been reports of hypersensitivity reactions which progressed to Kounis syndrome (acute allergic coronary arteriospasm that can result in myocardial infarction). Kounis syndrome can develop in patients with and without cardiac risk factors, and may present with cardiac and allergic symptoms. Coronary vasospasm may be eliminated with steroids, antihistamines, and spasmolytics treatment.

Injection Site Reactions

Injection site reactions, including redness, swelling, and pain, have been reported during post-marketing surveillance. Necrosis associated with extravasation has also been reported.

Other Events

Pain and asthenia were the most frequently reported miscellaneous adverse effects; their relationship to the tumor and to anemia was likely. Alopecia was reported (3%). Cardiovascular, respiratory, genitourinary, and mucosal side effects have occurred in 6% or less of the patients. Cardiovascular events (cardiac failure, embolism, cerebrovascular accidents) were fatal in less than 1% of the patients and did not appear to be related to chemotherapy. Cancer-associated hemolytic-uremic syndrome has been reported rarely. Hemolytic anemia (sometimes fatal) has also been reported (see **WARNINGS**).

Malaise, anorexia, hypertension, dehydration, and stomatitis have been reported as part of post-marketing surveillance.

OVERDOSAGE

There is no known antidote for Carboplatin Injection overdose. The anticipated complications of overdose would be secondary to bone marrow suppression and/or hepatic toxicity.

DOSAGE AND ADMINISTRATION

NOTE: Aluminum reacts with carboplatin causing precipitate formation and loss of potency, therefore, needles or intravenous sets containing aluminum parts that may come in contact with the drug must not be used for the preparation or administration of Carboplatin Injection.

Single Agent Therapy

Carboplatin Injection, as a single agent, has been shown to be effective in patients with

recurrent ovarian carcinoma at a dosage of 360 mg/m² IV on day 1 every 4 weeks (alternatively see **Formula Dosing**). In general, however, single intermittent courses of Carboplatin Injection should not be repeated until the neutrophil count is at least 2,000 and the platelet count is at least 100,000.

Combination Therapy with Cyclophosphamide

In the chemotherapy of advanced ovarian cancer, an effective combination for previously untreated patients consists of:

Carboplatin Injection - 300 mg/m² IV on day 1 every four weeks for six cycles (alternatively see **Formula Dosing**).

Cyclophosphamide - 600 mg/m² IV on day 1 every four weeks for six cycles. For directions regarding the use and administration of cyclophosphamide please refer to its package insert. (see **CLINICAL STUDIES**).

Intermittent courses of Carboplatin Injection in combination with cyclophosphamide should not be repeated until the neutrophil count is at least 2,000 and the platelet count is at least 100,000.

Dose Adjustment Recommendations

Pretreatment platelet count and performance status are important prognostic factors for severity of myelosuppression in previously treated patients.

The suggested dose adjustments for single agent or combination therapy shown in the table below are modified from controlled trials in previously treated and untreated patients with ovarian carcinoma. Blood counts were done weekly, and the recommendations are based on the lowest post-treatment platelet or neutrophil value.

| Platelets | Neutrophils | Adjusted Dose* (From Prior Course) |
|------------------|--------------------|---|
| >100,000 | >2,000 | 125% |
| 50-100,000 | 500-2,000 | No Adjustment |
| <50,000 | <500 | 75% |

* Percentages apply to Carboplatin Injection as a single agent or to both Carboplatin Injection and cyclophosphamide in combination. In the controlled studies, dosages were also adjusted at a lower level (50% to 60%) for severe myelosuppression. Escalations above 125% were not recommended for these studies.

Carboplatin Injection is usually administered by an infusion lasting 15 minutes or longer. No pre- or post-treatment hydration or forced diuresis is required.

Patients with Impaired Kidney Function

Patients with creatinine clearance values below 60 mL/min are at increased risk of severe bone marrow suppression. In renally-impaired patients who received single agent Carboplatin Injection therapy, the incidence of severe leukopenia, neutropenia, or thrombocytopenia has been about 25% when the dosage modifications in the table below have been used.

| Baseline Creatinine Clearance | Recommended Dose on Day 1 |
|--------------------------------------|----------------------------------|
| 41 - 59 mL/min | 250 mg/m ² |
| 16 - 40 mL/min | 200 mg/m ² |

The data available for patients with severely impaired kidney function (creatinine clearance below 15 mL/min) are too limited to permit a recommendation for treatment.

These dosing recommendations apply to the initial course of treatment. Subsequent dosages should be adjusted according to the patient's tolerance based on the degree of bone marrow suppression.

Formula Dosing

Another approach for determining the initial dose of Carboplatin Injection is the use of mathematical formulae, which are based on a patient's pre-existing renal function or renal function and desired platelet nadir. Renal excretion is the major route of elimination for carboplatin. (see **CLINICAL PHARMACOLOGY**). The use of dosing formulae, as compared to empirical dose calculation based on body surface area, allows compensation for patient variations in pretreatment renal function that might otherwise result in either underdosing (in patients with above average renal function) or overdosing (in patients with impaired renal function).

A simple formula for calculating dosage, based upon a patient's glomerular filtration rate (GFR in mL/min) and Carboplatin Injection target area under the concentration versus time curve (AUC in mg/mL•min), has been proposed by Calvert. In these studies, GFR was measured by ⁵¹Cr-EDTA clearance.

| CALVERT FORMULA FOR CARBOPLATIN DOSING |
|--|
| Total Dose (mg) = (target AUC) × (GFR + 25) |
| Note: With the Calvert formula, the total dose of Carboplatin Injection is calculated in mg, <u>not</u> mg/m². |

The target AUC of 4 mg/mL•min to 6 mg/mL•min using single agent Carboplatin Injection appears to provide the most appropriate dose range in previously treated patients. This study also showed a trend between the AUC of single agent Carboplatin Injection administered to previously treated patients and the likelihood of developing toxicity.

| % Actual Toxicity in Previously Treated Patients | | |
|--|----------------------------------|----------------------------|
| AUC (mg/mL•min) | Gr 3 or Gr 4 Thrombocytopenia | Gr 3 or Gr 4 Leukopenia |
| 4 to 5 | 16% | 13% |
| 6 to 7 | 33% | 34% |

Geriatric Dosing

Because renal function is often decreased in elderly patients, formula dosing of Carboplatin Injection based on estimates of GFR should be used in elderly patients to

provide predictable plasma Carboplatin Injection AUCs and thereby minimize the risk of toxicity.

PREPARATION OF INTRAVENOUS SOLUTIONS

Carboplatin Injection 10 mg/mL is supplied as a Ready To Use (RTU) sterile solution in 5 mL, 15 mL, 45 mL or 60 mL vials. Total content of carboplatin per vial is described in following table:

| Vial Strength | Diluent Volume |
|---------------|----------------|
| 50 mg | 5 mL |
| 150 mg | 15 mL |
| 450 mg | 45 mL |
| 600 mg | 60 mL |

Carboplatin Injection aqueous solution can be further diluted to concentrations as low as 0.5 mg/mL with 5% Dextrose in Water (D₅W) or 0.9% Sodium Chloride Injection, USP.

When prepared as directed, Carboplatin Injection aqueous solutions are stable for 8 hours at room temperature (25°C). Since no antibacterial preservative is contained in the formulation, it is recommended that Carboplatin Injection aqueous solutions be discarded 8 hours after dilution.

HOW SUPPLIED

Carboplatin Injection 10 mg/mL sterile solution is available in the following presentations:

| Unit of Sale | Concentration |
|--|----------------------------|
| NDC 61703-360-18 Carton containing 1 multiple-dose vial (Blue flip-off seals) | 50 mg/5 mL (10 mg/mL) |
| NDC 61703-150-05 Carton containing 1 multiple-dose vial (Blue flip-off seals) | 150 mg/15 mL (10 mg/mL) |
| NDC 61703-262-05 Carton containing 1 multiple-dose vial (Blue flip-off seals) | 450 mg/45 mL (10 mg/mL) |
| NDC 61703-600-05 Carton containing 1 multiple-dose vial (Blue flip-off seals) | 600 mg/60 mL (10 mg/mL) |

STORAGE

Unopened vials of Carboplatin Injection are stable to the date indicated on the package when stored at 25°C (77°F); excursions permitted from 15°-30°C (59°-86°F) [see USP Controlled Room Temperature]. Protect from light.

Carboplatin Injection multiple-dose vials maintain microbial, chemical, and physical

stability for up to 14 days at 25°C following multiple needle entries.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration. Solutions for infusion should be discarded 8 hours after preparation.

HANDLING AND DISPOSAL

Caution should be exercised in handling and preparing Carboplatin Injection. Several guidelines on this subject have been published.¹⁻⁴

To minimize the risk of dermal exposure, always wear impervious gloves when handling vials containing Carboplatin Injection. If Carboplatin Injection contacts the skin, immediately wash the skin thoroughly with soap and water. If Carboplatin Injection contacts mucous membranes, the membranes should be flushed immediately and thoroughly with water. More information is available in the references listed below.

REFERENCES

1. NIOSH Alert: Preventing occupational exposures to antineoplastic and other hazardous drugs in healthcare settings. 2004. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 2004- 165.
2. OSHA Technical Manual, TED 1-0.15A, Section VI: Chapter 2. Controlling occupational exposure to hazardous drugs. OSHA, 1999. http://www.osha.gov/dts/osta/otm/otm_vi/otm_vi_2.html
3. American Society of Health-System Pharmacists. ASHP guidelines on handling hazardous drugs. *Am J Health-Syst Pharm.* 2006; 63:1172-1193.
4. Polovich M, White JM, Kelleher LO, eds. 2005. Chemotherapy and biotherapy guidelines and recommendations for practice. 2nd ed. Pittsburgh, PA: Oncology Nursing Society.

Distributed by Hospira, Inc., Lake Forest, IL 60045

novaplus[™] 

Novaplus is a registered trademark of Vizient, Inc.

LAB-1505-2.0

Revised: 6/2024

PATIENT INFORMATION

R_x only

Read this entire leaflet carefully. Keep it for future reference.

Carboplatin Injection

This information will help you learn more about Carboplatin Injection. It cannot, however, cover all the possible warnings or side effects relating to Carboplatin Injection, and it does not list all of the benefits and risks of Carboplatin Injection. Your doctor should always be your first choice for detailed information about your medical condition and your treatment. Be sure to ask your doctor about any questions you may have.

What is cancer?

Under normal conditions, the cells in your body divide and grow in an orderly, controlled fashion. Cell division and growth are necessary for the human body to perform its functions and to repair itself. Cancer cells are different from normal cells because they are not able to control their own growth. The reasons for this abnormal growth are not yet fully understood.

A tumor is a mass of unhealthy cells that are dividing and growing fast and in an uncontrolled way. When a tumor invades surrounding healthy body tissue it is known as a malignant tumor. A malignant tumor can spread (metastasize) from its original location to other parts of the body.

What is Carboplatin Injection?

Carboplatin Injection is a medicine that is used to treat cancer of the ovaries. It acts by interfering with the division of rapidly multiplying cells, particularly cancer cells.

Who should not take Carboplatin Injection?

Treatment with Carboplatin Injection is not recommended if you:

- are allergic to Carboplatin Injection or other platinum-containing products;
- have a weakened blood-forming system (bone marrow depression) or significant bleeding;
- are pregnant, intend to become pregnant, or are breast-feeding a baby.

How is Carboplatin Injection used?

Only a professional experienced in the use of cancer drugs should give you this medication. Carboplatin Injection is given by dripping the medicine slowly and directly into a vein (intravenous infusion) for 15 minutes or longer. Your doctor will determine the dose of Carboplatin Injection for you based on your weight, height, and kidney function. Carboplatin Injection may be given alone or with other drugs. Treatment is usually repeated every four weeks for a number of cycles.

Before and after Carboplatin Injection treatment, your doctor may give you medication to lessen the nausea and vomiting associated with this cancer treatment.

What should you tell your doctor before starting treatment with Carboplatin Injection?

Discuss the benefits and risks of Carboplatin Injection with your doctor before beginning treatment.

Be sure to inform your doctor:

- If you are allergic to Carboplatin Injection or other platinum containing products;
- If you are or intend to become pregnant, since Carboplatin Injection may harm the developing fetus. It is important to use effective birth control while you are being treated with Carboplatin Injection;
- If you are breast-feeding, since nursing infants may be exposed to Carboplatin Injection in this way;
- If you are taking other medicines, including all prescription and non prescription (over-the-counter) drugs, since Carboplatin Injection may affect the action of other medicines;
- If you have any other medical problems, especially chicken pox (including recent exposure to adults or children with chicken pox), shingles, hearing problems, infection, or kidney disease, since treatment with Carboplatin Injection increases the risk and severity of these conditions.

What should I avoid while taking Carboplatin Injection?

If you are pregnant or think you might be pregnant, or if you are breast feeding, let your doctor know right away. Carboplatin Injection may harm your developing fetus or breast-feeding baby. If you are a woman of childbearing age, you should use birth control to avoid getting pregnant while you are taking Carboplatin Injection.

You should avoid contact with adults and children who have infections, and tell your doctor right away if you show signs of infection such as cough, fever, and/or chills.

Also, while you are being treated with Carboplatin Injection or after you stop treatment, first check with your doctor **before** getting any immunizations (vaccinations). Avoid contact with adults or children who have received oral polio vaccine since they can pass the polio virus to you.

What are the possible side effects of Carboplatin Injection?

Carboplatin Injection may cause unwanted effects, particularly because Carboplatin Injection interferes with the growth of normal cells as well as cancer cells. For example, the occurrence of another cancer (secondary malignancy) has been reported in patients receiving cancer chemotherapy with multiple drugs. It is not always possible to tell whether such effects are caused by Carboplatin Injection, another drug you may be taking, or your illness. Because some of these effects may be serious, you will need close medical supervision during treatment with Carboplatin Injection.

The most serious side effects of Carboplatin Injection are:

- **bleeding and reduced blood cells, including reduced red blood cells (anemia) and platelets (needed for proper blood clotting)**, which may be severe enough to require blood transfusion. You should tell your doctor right away if you notice any unusual bruising or bleeding, including black tarry stools or blood in the urine.
- **infection** – Carboplatin Injection can temporarily lower the number of white blood cells in your blood, increasing the risk of infection;
- **life-threatening allergic reaction** – during and after treatment the doctor or nurse will observe you carefully for signs of allergic reaction;
- **kidney and liver problems;**
- **loss of hearing or ringing in the ears;**

Contact your doctor right away if you experience any of these effects, or notice effects that worry you or are troublesome.

Of the less serious side effects associated with Carboplatin Injection treatment, the most common are nausea, vomiting, diarrhea, loss of appetite, hair loss and numbness, tingling, burning, or pain in the hands or feet.

This medicine was prescribed for your particular condition. It must be given under close medical supervision by a doctor trained in the use of drugs for the treatment of cancer. This summary does not include everything there is to know about Carboplatin Injection. Medicines are sometimes prescribed for purposes other than those listed in patient leaflets. If you have questions or concerns, or want more information about Carboplatin Injection, your physician and pharmacist have the complete prescribing information upon which this information is based. You may want to read it and discuss it with your doctor. Remember, no written summary can replace careful discussion with your doctor.

Distributed by Hospira, Inc., Lake Forest, IL 60045

novaplus[™] 

Novaplus is a registered trademark of Vizient, Inc.
LAB-1506-1.0
Revised: 3/2022

PRINCIPAL DISPLAY PANEL - 5 mL Vial Label

Sterile

NDC 61703-360-18

5 mL Multiple-dose Vial

Rx only

CARBOplatin

Injection

50 mg/5 mL

(10 mg/mL)

For Intravenous Use

Cytotoxic Agent

Sterile NDC 61703-360-18
5 mL Multiple-dose Vial Rx only

CARBOplatin Injection

50 mg/5 mL
(10 mg/mL)

For Intravenous Use

Cytotoxic Agent

Each vial contains:
50 mg Carboplatin and
Water for Injection, USP q.s.
CONTAINS NO PRESERVATIVES.

**READ ACCOMPANYING
CIRCULAR FOR COMPLETE
DIRECTIONS FOR USE.**

**Store at 25°C (77°F); excursions
permitted to 15°-30°C (59°-86°F)
(see USP Controlled Room
Temperature).**

Protect from light.

Retain in carton until time of use.
Distributed by Hospira, Inc.
Lake Forest, IL 60045

novaplus⁺



(01)10361703360189

PAA182748



LOT S014321AA
EXP YYYYMMDD

PRINCIPAL DISPLAY PANEL - 5 mL Vial Carton

VIAL

1 x 5 mL Vial
NDC 61703-360-18

Sterile
Rx only

CARBOplatin
Injection

50 mg/5 mL
(10 mg/mL)

For Intravenous Use

Multiple-dose Vial

Cytotoxic Agent

Cytotoxic Agent

(01)00361703360182



VIAL



VIAL

Sterile

CARBOplatin Injection

50 mg/5 mL (10 mg/mL)



LOT S014321AA
EXP YYYYMMDD
SN XXXXXXXX
GTIN 00361703360182

*These manufacturing details,
along with 2D datamatrix,
will be printed at time of packing.*



VIAL



VIAL

1 x 5 mL Vial

NDC 61703-360-18

Sterile

Rx only

CARBOplatin
Injection

50 mg/5 mL
(10 mg/mL)

For Intravenous Use



VIAL

Each vial contains:

50 mg Carboplatin and Water for Injection,
USP q.s.

CONTAINS NO PRESERVATIVES.

**READ ACCOMPANYING CIRCULAR FOR
COMPLETE DIRECTIONS FOR USE.**

Store at 25°C (77°F); excursions
permitted to 15°-30°C (59°-86°F)
(see USP Controlled Room Temperature).

Protect from light.

Retain in carton until time of use.



MADE IN GERMANY

Distributed by Hospira, Inc.
Lake Forest, IL 60045



Multiple-dose Vial

Cytotoxic Agent

Cytotoxic Agent



PRINCIPAL DISPLAY PANEL - 15 mL Vial Label

Sterile
 NDC 61703-150-05
 15 mL Multiple-dose Vial
 Rx only
 CARBOplatin
 Injection
 150 mg/15 mL
 (10 mg/mL)
 For Intravenous Use
 Cytotoxic Agent

Sterile NDC 61703-150-05
 15 mL Multiple-dose Vial Rx only

**CARBOplatin
 Injection**

**150 mg/15 mL
 (10 mg/mL)**

For Intravenous Use
Cytotoxic Agent

Each vial contains:
 150 mg Carboplatin and
 Water for Injection, USP
 q.s.

CONTAINS NO
 PRESERVATIVES.

READ
 ACCOMPANYING
 CIRCULAR FOR
 COMPLETE
 DIRECTIONS FOR USE.

Store at 25°C (77°F);
 excursions permitted
 to 15°-30°C (59°-86°F)
 (see USP Controlled
 Room Temperature).
Protect from light.
 Retain in carton until
 time of use.

Distributed by
 Hospira, Inc.
 Lake Forest, IL 60045

novaplus+

(01)10361703150056
 PAA182750



LOT S014321AA
 EXP YYYYYMMMDD

PRINCIPAL DISPLAY PANEL - 15 mL Vial Carton

VIAL

1 x 15 mL Vial
NDC 61703-150-05

Sterile
Rx only

CARBOplatin
Injection

150 mg/15 mL
(10 mg/mL)

For Intravenous Use

Multiple-dose Vial

Cytotoxic Agent

Cytotoxic Agent

(01)00361703150059



VIAL

VIAL



Sterile

CARBOplatin Injection

150 mg/15 mL (10 mg/mL)

LOT S014321AA
EXP YYYYMMDD
SN XXXXXXXX
GTIN 00361703150059

*These manufacturing details,
along with 2D datamatrix,
will be printed at time of packing.*

VIAL



VIAL



1 x 15 mL Vial

NDC 61703-150-05

Sterile

Rx only

CARBOplatin
Injection

150 mg/15 mL
(10 mg/mL)

For Intravenous Use

VIAL



Each vial contains:

150 mg Carboplatin and Water for Injection, USP q.s.

CONTAINS NO PRESERVATIVES.

**READ ACCOMPANYING CIRCULAR
FOR COMPLETE DIRECTIONS FOR USE.**

Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F) (see USP Controlled Room Temperature).

Protect from light.

Retain in carton until time of use.

MADE IN GERMANY

Distributed by Hospira, Inc.
Lake Forest, IL 60045

novaplus⁺


Novaplus is a registered trademark of

Multiple-dose Vial

Cytotoxic Agent

Cytotoxic Agent





PRINCIPAL DISPLAY PANEL - 45 mL Vial Label


Sterile
NDC 61703-262-05
45 mL Multiple-dose Vial
Rx only
CARBOplatin
Injection
450 mg/45 mL
(10 mg/mL)
For Intravenous Use
Cytotoxic Agent

Sterile NDC 61703-262-05
45 mL Multiple-dose Vial Rx only

**CARBOplatin
Injection**
450 mg/45 mL
(10 mg/mL)
For Intravenous Use
Cytotoxic Agent

Each vial contains:
450 mg Carboplatin and
Water for Injection, USP q.s.
CONTAINS NO
PRESERVATIVES.
**READ ACCOMPANYING
CIRCULAR FOR
COMPLETE
DIRECTIONS FOR USE.**
Store at 25°C (77°F);
excursions permitted to
15°-30°C (59°-86°F)
(see USP Controlled Room
Temperature).
Protect from light.
Retain in carton until time
of use.
Distributed by Hospira, Inc.
Lake Forest, IL 60045

novaplus
Novaplus is a registered
trademark of Vizient, Inc.



(01) 10361703262056
PAA162764



↑

| | |
|-----|-----------|
| LOT | S014321AA |
| EXP | YYYYMMDD |

PRINCIPAL DISPLAY PANEL - 45 mL Vial Carton

VIAL

1 x 45 mL Vial
NDC 61703-262-05

Sterile
Rx only

CARBOplatin
Injection



450 mg/45 mL
(10 mg/mL)

For Intravenous Use

Multiple-dose Vial

Cytotoxic Agent

LOT S014321AA
EXP YYYYMMDD
SN XXXXXXXX
GTIN 00361703262059
*These manufacturing details,
along with 2D datamatrix,
will be printed at time of packing.*



PRINCIPAL DISPLAY PANEL - 60 mL Vial Label

Sterile
NDC 61703- 600-05
60 mL Multiple-dose Vial
Rx only
CARBOplatin
Injection

600 mg/60 mL
(10 mg/mL)

For Intravenous Use

Cytotoxic Agent

Sterile NDC 61703- 600-05
60 mL Multiple-dose Vial Rx only

**CARBOplatin
Injection**

**600 mg/60 mL
(10 mg/mL)**

For Intravenous Use

Cytotoxic Agent

Each vial contains:
600 mg Carboplatin and
Water for Injection, USP q.s.
CONTAINS NO
PRESERVATIVES.

**READ ACCOMPANYING
CIRCULAR FOR COMPLETE
DIRECTIONS FOR USE.**

**Store at 25°C (77°F);
excursions permitted to
15°-30°C (59°-86°F) (see USP
Controlled Room
Temperature).**

Protect from light.
Retain in carton until time of
use.


Distributed by Hospira, Inc.
Lake Forest, IL 60045

novaplus⁺

Novaplus is a registered
trademark of Vizient, Inc.


(01)10361703600056

PA1182767



PRINCIPAL DISPLAY PANEL - 60 mL Vial Carton

VIAL

1 x 60 mL Vial
NDC 61703-600-05

Sterile
Rx only


CARBOplatin
Injection

600 mg/60 mL
(10 mg/mL)

For Intravenous Use

Multiple-dose Vial

Cytotoxic Agent



CARBOPLATIN

carboplatin injection, solution

Product Information

| | | | |
|--------------------------------|-------------------------|---------------------------|---------------|
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:61703-360 |
| Route of Administration | INTRAVENOUS | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|-----------------|-------------------|----------|
|-----------------|-------------------|----------|

| | | | | |
|---|---|---|-----------------------------|---------------------------|
| CARBOPLATIN (UNII: BG3F62OND5) (CARBOPLATIN - UNII:BG3F62OND5) | | CARBOPLATIN | 10 mg in 1 mL | |
| Inactive Ingredients | | | | |
| Ingredient Name | | | Strength | |
| WATER (UNII: 059QF0KO0R) | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:61703-360-18 | 1 in 1 CARTON | 05/23/2022 | |
| 1 | | 5 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | | Marketing Start Date | Marketing End Date |
| ANDA | ANDA076517 | | 05/23/2022 | |

| | | | | |
|--|-------------------------|--|-----------------------------|---------------------------|
| CARBOPLATIN | | | | |
| carboplatin injection, solution | | | | |
| Product Information | | | | |
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:61703-150 | |
| Route of Administration | INTRAVENOUS | | | |
| Active Ingredient/Active Moiety | | | | |
| Ingredient Name | | | Basis of Strength | Strength |
| CARBOPLATIN (UNII: BG3F62OND5) (CARBOPLATIN - UNII:BG3F62OND5) | | | CARBOPLATIN | 10 mg in 1 mL |
| Inactive Ingredients | | | | |
| Ingredient Name | | | Strength | |
| WATER (UNII: 059QF0KO0R) | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:61703-150-05 | 1 in 1 CARTON | 05/23/2022 | |
| 1 | | 15 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a | | |

| Combination Product | | | |
|------------------------------|--|----------------------|--------------------|
| Marketing Information | | | |
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| ANDA | ANDA076517 | 05/23/2022 | |

| CARBOPLATIN | | | | |
|--|--|--|----------------------|--------------------|
| carboplatin injection, solution | | | | |
| Product Information | | | | |
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:61703-262 | |
| Route of Administration | INTRAVENOUS | | | |
| Active Ingredient/Active Moiety | | | | |
| Ingredient Name | | Basis of Strength | Strength | |
| CARBOPLATIN (UNII: BG3F62OND5) (CARBOPLATIN - UNII:BG3F62OND5) | | CARBOPLATIN | 10 mg in 1 mL | |
| Inactive Ingredients | | | | |
| Ingredient Name | | Strength | | |
| WATER (UNII: 059QF0KO0R) | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:61703-262-05 | 1 in 1 CARTON | 05/23/2022 | |
| 1 | | 45 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| ANDA | ANDA076517 | 05/23/2022 | | |

| | | | |
|---------------------------------|--|--|--|
| CARBOPLATIN | | | |
| carboplatin injection, solution | | | |
| Product Information | | | |

| | | | |
|--------------------------------|-------------------------|---------------------------|---------------|
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:61703-600 |
| Route of Administration | INTRAVENOUS | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|---------------|
| CARBOPLATIN (UNII: BG3F62OND5) (CARBOPLATIN - UNII:BG3F62OND5) | CARBOPLATIN | 10 mg in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:61703-600-05 | 1 in 1 CARTON | 05/23/2022 | |
| 1 | | 60 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA077059 | 05/23/2022 | |

Labeler - Hospira, Inc. (141588017)

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------------------|---------|-----------|---|
| Hospira Australia Pty Ltd | | 758967652 | ANALYSIS(61703-150, 61703-262, 61703-360, 61703-600) , MANUFACTURE(61703-150, 61703-262, 61703-360, 61703-600) , PACK(61703-150, 61703-262, 61703-360, 61703-600) , LABEL(61703-150, 61703-262, 61703-360, 61703-600) |

Revised: 8/2024

Hospira, Inc.