LIDOCAINE- lidocaine cream Nivagen Pharmaceuticals, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Lidocaine 5%

Drug Facts

Active Ingredient

Lidocaine 5% w/w

Purpose

Local anesthetic

Uses

Helps relieve the pain, itching, and burning associated with hemorrhoids and other anorectal disorders.

Warnings

For external use only.

When using this product

- avoid contact with eyes
- do not exceed recommended dosage unless directed by a doctor
- do not put this product into the rectum by using fingers or any mechanical device or applicator.

Stop use and ask a doctor if

- rectal bleeding occurs
- condition worsens or does not improve within 7 days
- allergic reaction occurs to ingredients in this product
- symptom being treated does not subside or if redness, irritation, swelling, pain or other symptoms develop or increase
- symptoms clear up and return within a few days

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or soft cloth before applying.
- Adults and children 12 years and older: apply externally to the affected area up to 6 times a day.
- Children under 12 years of age: consult a doctor.
- To use finger cots: Roll one finger cot over finger. Gently squeeze cream onto finger cot. Smooth a layer of the cream over affected area.

Other Information

Store at 25°C (77°C); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature].

Inactive Ingredients

benzyl alcohol,carbomer hompolymer type C, C10-18 triglycerides, cholesterol, lecithin, isopropyl myristate, polysorbate 80, propylene glycol, tocopherol, trolamine, water

Questions?

1-877-977-0687

PRINCIPAL DISPLAY PANEL - 30 g Tube Carton

NDC 75834-113-30

Lidocaine 5% Anorectal Cream

Relieves:

- Pain
- Itching
- Burning

NIVAGEN PHARMACEUTICALS

net wt. 30 g (1 oz)

NDC 75834-113-30



Relief of Pain for Hemorrhoids & Other Anorectal Disorders



Lidocaine 5%

Anorectal Cream

FINGER COTS INCLUDED

> Caution: The Finger Cots Contain Natural Rubber Latex Which May Cause Allergic Reactions

Relieves Pain Quickly

Lidocaine 5% Anorectal Cream

NDC 75834-113-30

Lidocaine 5%

Anorectal Cream



Relieves:

✓ Pain

✓ Itching

Burning

net wt. 30 g (1 oz)

Lidocaine 5%

Anorectal Cream





See end flap for Lot No. and exp. date Rev 8/16

7830-776-778-1 Nivagen Pharmaceuticals, Inc Sacramento, CA 95827 USA



Push down and furn counterclockwise. To open child-resistant cap:

the tube is damaged. Do not use if tube seal is not intact or if Child-Resistant & Tamper Evident; Anorectal Cream

%g əuisəobi**7**

T880-TT8-TT8-1 SRIOITS94D

80, propylene glycol, tocopherol, trolamine, water cholesterol, lecithin, isopropyl myristate, polysorbate hompolymer type C, C10-18 triglycerides, Inactive Ingredients benzylalcohol,carbomer

[see USP Controlled Room Temperature]. excursions permitted to 15°-30°C (59°-86°F) Other Information Store at 25°C (77°C);

layer of the cream over affected area. Gently squeeze cream onto finger cot. Smooth a ■ To use finger cots: Roll one finger cot over finger Children under 12 years of age: consult a doctor. externally to the affected area up to 6 times a day. Yearls and children 12 years and older: apply

Drug Facts (continued)

ussne or son cloth betore applying. thoroughly. Gently dry by patting or blotting with toilet stea with mild soap and warm water and rinse Directions - When practical, cleanse the affected

help or contact a Poison Control Center right away. Keep out of reach of children. If swallowed, get medical

- ■eymptoms clear up and return within a few days gevelop or increase
- redness, irritation, swelling, pain or other symptoms symptom being treated does not subside or if
- silergic reaction occurs to ingredients in this product ■ condition worsens or does not improve within 7 days
 - rectal bleeding occurs

Stop use and ask a doctor it

Drug Facts (continued)

fingers or any mechanical device or applicator.

- do not put this product into the rectum by using directed by a doctor
 - do not exceed recommended dosage unless
 - svojq contact with eyes

Myeu najud tuja broduct


For external use only. **Marnings**

anorectal disorders.

burningassociated with hemorrhoids and other USes Helps relieve the pain, itching, and

LIDOCRING 5% W/W. " Focal anesthetic Active Ingredient.

Side Hacis



LIDOCAINE

lidocaine cream

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:75834-113	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Lidocaine (UNII: 98PI200987) (Lidocaine - UNII:98PI200987)	Lidocaine	50 mg in 1 g	

Inactive Ingredients		
Ingredient Name	Strength	
BENZYL ALCOHOL (UNII: LKG8494WBH)		
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)		
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)		
CHOLESTEROL (UNII: 97C5T2UQ7J)		
LECITHIN, SO YBEAN (UNII: 1DI56 QDM62)		
ISOPROPYL MYRISTATE (UNII: 0 RE8 K4LNJS)		
POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
TOCOPHEROL (UNII: R0ZB2556P8)		
TROLAMINE (UNII: 9O3K93S3TK)		
WATER (UNII: 059QF0KO0R)		

Product Characteristics			
Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging				
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:75834-113-30	30 g in 1 TUBE; Type 0: Not a Combination Product	12/15/20 16	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH FINAL	part346	12/15/20 16		

Labeler - Nivagen Pharmaceuticals, Inc. (052032418)

Revised: 1/2020 Nivagen Pharmaceuticals, Inc.