

SKIN METAL GREEN STEALTH- zinc oxide lotion
Skin Metal

Skin Metal Green Stealth

Skin Metal Green Stealth

Drug Facts

Active ingredient
Zinc Oxide (15%)

Purpose
Sunscreen

Uses

- Helps prevent sunburn.
- It is used in conjunction with other sun protection measures like: Avoidance, decreasing the use of sun cancer and early skin aging (DAILY METAL).

Warnings

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if you become itchy or red.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Wash well before use.

- Apply liberally. 15 minutes before sun exposure.

- Reapply after 80 minutes of swimming or sweating.

- Immediately after towel drying.

- Use every 2 hours.

- Children under 6 months: Ask a doctor.

- **Sun Protection Measures** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures.

- **Use** in the sun, especially from 10 a.m. - 2 p.m.

- Wear long-sleeved shirts, pants, hats, and sunglasses.

Inactive Ingredients

Water, Dimethicone, Capryl/Capric Triglyceride, Lauryl PEG-9 Dimethicone, Trisnopal Salicylate.

Drug Facts (Continued)

Active ingredients

Octinoxate

Octisalate

Octocrylene

Avobenzone

Homosalate

Butyl Methoxy

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

Succinate

Diethylmalonate

Diethylhexyl

DAILY METAL
Green Smooth SPF 51

**125
skinnmetal**

BROAD SPECTRUM SPF 51 SUNSCREEN

WATER RESISTANT (80 MINUTES)

NET WT. 3.8 FL. OZ. 55 mL



Skin Metal Green Stealth

Drug Facts

Active Ingredient	Purpose
Zinc Oxide 12.5%	Sunscreen

Uses

- Helps prevent sunburn.
- Helps prevent dry skin.
- Helps prevent skin cancer when used with other sun protection measures (see **Directions**, **Warnings** for use, and **Directions** for use of sunscreen and other sun protection measures).

Warnings

For external use only.

- Do not use on damaged or broken skin.
- When using this product keep out of eyes. Rinse with water to remove.
- Stop use and ask a doctor if rash occurs.
- Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake well before use.
- Apply liberally 15 minutes before sun exposure.
- Reapply:
- After 80 minutes of swimming or sweating.
- Immediately after towel drying.
- At least every 2 hours.
- Cotton swab or sponge. Ask a doctor.

Sun Protection Measure Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 30 or higher and other sun protection measures including:

- Limiting time in the sun, especially from 10 a.m. to 2 p.m.
- Wear long-sleeved shirts, pants, hats, and sunglasses.

Inactive Ingredients
Base: Dimethicone, Caprylic/Capric Triglyceride, Butyl Stearate, Dimethicone Trisiloxane, **125/32**

Drug Facts (Continued)

Active Ingredients (continued)

DAILY METAL
Green Stealth SPF 51

125/32
skinneral

Question or Comments?
CALL 1-800-4-A-MEDICINE

OTT, SKIN METAL, INC.
www.skinmetal.com

**USA: 1-800-4-A-MEDICINE
CANADA: 1-800-268-3873**

**BROAD SPECTRUM SPF 51 SUNSCREEN
ANTHRACENYL 180 MINUTE**
NET WT. 3.8 FL. OZ. 55 mL

8 60014 52731

Skin Metal Green Stealth

Drug Facts

Active Ingredient	Purpose
Zinc Oxide (8%)	Sunscreen

Uses

- Helps prevent sunburn.
- When used correctly with other sun protection measures (see Directions), decreasing the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

- Do not use on damaged or broken skin.
- When using this product keep out of eyes. Rinse with water to remove.
- Stop use and ask a doctor if redness occurs.
- Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake well before use.
- Apply liberally 15 minutes before sun exposure.
- Reapply:
 - After 80 minutes of swimming or sweating.
 - Immediately after towel drying.
 - At least every 2 hours.
- Children under 6 months: Ask a doctor.
- Sun Protection Measure:** Spending limited time in the sun reduces your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - Avoiding the sun, especially from 10 a.m. to 4 p.m.
 - Wearing long-sleeved shirts, pants, hats, and sunglasses.

Inactive Ingredients

Water, Dimethicone, Caprylic/Capric Triglyceride, Isopropyl Alcohol, Dimethicone, Triethyl Salicylate.

Drug Facts (Continued)

Active ingredients (continued)

Other Ingredients

Octyl Methacrylate, Dimethylsiloxane, Polybutyl Acrylate, Caprylic/Capric Triglyceride, Hydroxyethyl Acrylate, Ethylhexyl Methacrylate, Isododecane, Polyethylene Glycol-10, Polyethylene Glycol-15, Polyethylene Glycol-20, Polyethylene Glycol-30, Polyethylene Glycol-40, Polyethylene Glycol-60, Polyethylene Glycol-80, Polyethylene Glycol-100, Polyethylene Glycol-120, Polyethylene Glycol-150, Polyethylene Glycol-180, Polyethylene Glycol-200, Polyethylene Glycol-250, Polyethylene Glycol-300, Polyethylene Glycol-350, Polyethylene Glycol-400, Polyethylene Glycol-450, Polyethylene Glycol-500, Polyethylene Glycol-600, Polyethylene Glycol-750, Polyethylene Glycol-900, Polyethylene Glycol-1000, Polyethylene Glycol-1200, Polyethylene Glycol-1500, Polyethylene Glycol-1800, Polyethylene Glycol-2000, Polyethylene Glycol-2500, Polyethylene Glycol-3000, Polyethylene Glycol-3500, Polyethylene Glycol-4000, Polyethylene Glycol-4500, Polyethylene Glycol-5000, Polyethylene Glycol-6000, Polyethylene Glycol-7500, Polyethylene Glycol-9000, Polyethylene Glycol-10000.

DAILY METAL

Green Stealth SPF 51

125/32

125/32 skinmetal

QUESTIONS or COMMENTS?
Email: info@skinmetal.com

SKIN METAL, INC.
1000 N. W. 10TH AVE.
MIAMI, FL 33136

**MADE IN THE USA OF
COMBINE A FOREIGN
PRODUCT**

BROAD SPECTRUM SPF 51 SUNSCREEN
WATER RESISTANT (80 MINUTES)
NET WT. 3.3 FL. OZ. 85 mL

Skin Metal Green Stealth

Drug Facts

Active Ingredient	Purpose
Zinc Oxide (15%)	Sunscreen

Uses

- Helps prevent sunburn.
- It used as directed with other sun protection measures like clothing, decreasing the risk of skin cancer and early skin aging.

Warnings

For external use only.

- Do not use on damaged or broken skin.
- When using this product keep out of eyes. Rinse with water to remove.
- Stop use and ask a doctor if rash occurs.
- Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake well before use.
- Apply liberally 15 minutes before sun exposure.
- Reapply:
 - After 80 minutes of swimming or sweating.
 - Immediately after towel drying.
 - at least every 2 hours.
- Children under 6 months: Ask a doctor.
- Sun Protection Measure:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - Limiting time in the sun, especially from 10 a.m. to 2 p.m.
 - Wearing long-sleeved shirts, pants, hats, and sunglasses.

Inactive Ingredients

Water, Dimethicone, Capryl/Capric Triglyceride, Lauryl PEG-9 Dimethicone, Trisopal Silylate

Drug Facts (Continued)

Active Ingredients (Continued)

Dimethicone, C14-20 Alkyl Dimethicones, Bisphenol Acrylate, Capryl Methacrylate, Hexyl Methacrylate, Hydroxyethyl Methacrylate, Glyceryl Triacrylate, Hydroxyethyl Methacrylate, Propylene Glycol, Stearyl Dimethyl Silylate, Diphenylhexatriene, Octyl Methacrylate, Crocodylimer, Sodium Chloride, Octylalcohol, Hydroxyacetone, Chromium Oxide Green, Phosphorylcholine, Glycerol, C12-15 Alkyl Lactate, Polybutyl Methacrylate, Iron Oxides, C12-15 Alkyl Benzoate, Capryl Glycol, Aluminum Dimethacrylate, Hydroxyethyl Methacrylate, Hexylene Glycol, Tetrahydrofuran, Chloroform, Triethylacrylate, Polymethyl Methacrylate, Polybutyl Methacrylate, Hydroxyethyl Methacrylate, Hydroxyethyl Methacrylate, Sodium Hydroxide.

Contains or Contains Derivatives of: DRUG INGREDIENTS

SEE: SKIN METAL, INC.
www.skinmetal.com

MADE IN THE USA OF DOMESTIC AND FOREIGN MATERIALS.

DAILY METAL
Green Stealth SPF 51

125 skinmetal

BROAD SPECTRUM SPF 51 SUNSCREEN
WATER RESISTANT (80 MINUTES)
NET WT. 3.3 FL. OZ. 95 mL

8 60014 52731 9

SKIN METAL GREEN STEALTH

zinc oxide lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:87098-270
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	132 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BAKUCHIOL (UNII: OT12HJU3AR)	
CI 77288 (UNII: X5Z09SU859)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
C13-15 ALKANE (UNII: 114P5I43UJ)	
GLYCERYL STEARATE (UNII: 230OU9XXE4)	
LAURYL PEG-8 DIMETHICONE (300 CPS) (UNII: ELL2U7K8T8)	
OCTYLDODECYL NEOPENTANOATE (UNII: X8725R883T)	
CAPRYLYL METHICONE (UNII: Q95M2P1KJL)	
HYDROGENATED SOYBEAN LECITHIN (UNII: H1109Z9J4N)	
POLYGLYCERYL-4 DIISOSTEARATE/POLYHYDROXYSTEARATE/SEBACATE (UNII: 687U3PEB2Y)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
CAPRYLYL GLYCOL (UNII: 00YIU5438U)	
ALLANTOIN (UNII: 344S277G0Z)	
.ALPHA.-BISABOLOL, (+)- (UNII: 105S6I733Z)	
BUTYLOCTYL SALICYLATE (UNII: 2EH13UN8D3)	
NIACINAMIDE (UNII: 25X51I8RD4)	
WATER (UNII: 059QF0K00R)	
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
HEXYLENE GLYCOL (UNII: KEH0A3F75J)	
PROPANEDIOL (UNII: 5965N8W85T)	
TETRASODIUM GLUTAMATE DIACETATE (UNII: 5EHL50I4MY)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
ALKYL (C12-15) BENZOATE (UNII: A9EJ3J61HQ)	
CAPRYLIC/CAPRIC TRIGLYCERIDE (UNII: C9H2L21V7U)	
DIMETHICONE CROSSPOLYMER (UNII: UF7620L1W6)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CI 77492 (UNII: EX438O2MRT)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
TRIDECYL SALICYLATE (UNII: AZQ08K38Z1)	

Product Characteristics

Color	green	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87098-270-02	1 in 1 CARTON	11/03/2025	
1	NDC:87098-270-01	55 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	11/03/2025	

Labeler - Skin Metal (119577227)

Revised: 1/2026

Skin Metal