

PINWORM TREATMENT- pyrantel pamoate suspension
AMERISOURCE BERGEN

pinworm treatment-ABC

Active ingredient (in each mL)

Pyrantel Pamoate 144mg/mL

(the equivalent of 50mg pyrantel base)

Purpose

Anthelmintic

Use

for the treatment of pinworms

Warnings

Do not exceed recommended dosage

Ask a doctor before use if

■ you are pregnant ■ have liver disease ■ nursing a baby

When using this product

abdominal cramps, nausea, vomiting, diarrhea, headache, or dizziness sometimes occur after taking this drug. If any of these symptoms persist, consult a doctor.

Keep out of reach of children.

In case of overdose get medical help or contact a poison control center right away.

Directions

■ **shake well before use**

■ **read bottle label and package insert carefully before taking this medication**

■ treat the entire household unless otherwise advised. See Warnings

■ do not repeat treatment unless directed by a doctor

■ this product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during or after medication

■ if signs of pinworms persist after treatment, consult a doctor

| Weight | Dosage (taken as a single dose) |
|-------------------------------------|---|
| Less than 25 lb. or under 2 yrs.old | Do not use unless directed by a doctor |
| 25-37 lb. | 1/2 teaspoonful |
| 38-62 lb. | 1 teaspoonful |
| 63-87 lb. | 1 1/2 teaspoonfuls |
| 88-112 lb. | 2 teaspoonfuls |
| 113-137 lb. | 2 1/2 teaspoonfuls |
| 138-162 lb. | 3 teaspoonfuls |
| 163-187 lb. | 3 1/2 teaspoonfuls |
| 188 lb. and over | 4 teaspoonfuls |

Other information

■ Store between 15°C-30°C (59°- 86°F)

Inactive ingredients

banana flavor, citric acid, glycerin, lecithin, magnesium aluminum silicate, methylcellulose, methylparaben, povidone, propylene glycol, propylparaben, purified water, simethicone, sodium benzoate, sodium citrate, sodium saccharin and sorbitol.

Questions or comments?

call toll-free **1-888-235-2466**

CAUTION: DO NOT USE IF IMPRINTED SEAL ON BOTTLE IS BROKEN OR MISSING

Distributed by:

AmerisourceBergen

1 West First Avenue

Conshohocken, PA 19428

Questions or Concerns?

www.mygnp.com

MADE IN INDIA

Code: TN/Drugs/TN00002222/2006

949129349

L0001159

R0925

READ THIS INSERT CAREFULLY BEFORE TAKING MEDICATION

PINWORM MEDICINE

Pyrantel Pamoate 144 mg/mL
(the equivalent of 50 mg Pyrantel base per mL)

FOR THE TREATMENT OF PINWORMS

DIRECTIONS

Adults, children 12 years of age and over, and children 2 years to under 12 years of age: oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight not to exceed 1 gram.

Take only according to directions and do not exceed the recommended dosage unless directed by a doctor. Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See warnings. If any worms other than pinworms are present before or after treatment, consult a doctor. If any symptoms or pinworms are still present after treatment, consult a doctor.

Shake well before using.

| Weight | Dosage (taken as a single dose) |
|--|--|
| Less than 25 pounds or under 2 years old | consult a doctor |
| 25 to 37 pounds | ½ teaspoonful |
| 38 to 62 pounds | 1 teaspoonful |
| 63 to 87 pounds | 1½ teaspoonfuls |
| 88 to 112 pounds | 2 teaspoonfuls |
| 113 to 137 pounds | 2½ teaspoonfuls |
| 138 to 162 pounds | 3 teaspoonfuls |
| 163 to 187 pounds | 3½ teaspoonfuls |
| 188 pounds and over | 4 teaspoonfuls |

This product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during, or after medication.

WARNINGS

Abdominal cramps, nausea, vomiting, diarrhea, headaches, or dizziness, sometimes occur after taking this drug, if any of these conditions persist consult a doctor.

If you are pregnant, nursing or have liver disease, do not take this product unless directed by your doctor.

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

OTHER INFORMATION

This brochure will tell you how to identify and treat pinworm infections effectively with pinworm medicine. Many who have pinworms do not have symptoms, however, many will have itching in and around the rectal opening. The itching may be very annoying and

continued scratching can cause an irritation in this area. The itching may be prominent at night when sleeping. That is why restless sleep is sometimes a sign of pinworms, especially in children. Other symptoms include insomnia, gastrointestinal distress, irritability, enuresis (bed wetting) and secondary infection due to localized scratching. You should treat with pinworm medicine only after you have seen and identified the pinworms.

Pinworms look like tiny white threads. The female pinworm is about 1/2 inch long (the male pinworm is shorter); they live in the bowel. Usually, at night, the female pinworm travels to the rectal opening and lays eggs within the skin folds around the opening. This usually takes place within 1-2 hours after the child has been put to sleep for the night. The sticky gelatin like substance in which the eggs (they are too small to see) are deposited and the movement of the female pinworm may cause annoying itching and possibly restless sleep. By checking the rectal opening at this time, you can attempt to see the pinworm as described.

Scratching will cause pinworm eggs to stick to the fingers. Reinfection will result if the fingers are placed in the mouth. The eggs, which are too small to see, contaminate whatever they come in contact with: bed clothes, underwear, toys, hands, and food touched by contaminated hands. Even eggs floating in the air can be swallowed and cause infection. Eggs deposited can survive for as long as 3 weeks. Pinworms are highly contagious; therefore, after treatment with pinworm medication you should wash hands and fingernails with soap often during the day in order to prevent spreading to others and reinfection. Wear tight underpants both day and night. For several days after treatment clean the bedroom floor by vacuuming or damp mopping. After treatment, wash bed linens and night clothes (don't shake them). Keep toilet seats clean.

Distributed by:
AmerisourceBergen

1 West First Avenue

Conshohocken, PA 19428

Questions or Concerns?www.mygnp.com

MADE IN INDIA

948026898

L0001176

R1025

Code: TN/Drugs/TN00002222/2006

carton label 1oz

GOOD

NEIGHBOR

PHARMACY

NDC 46122-825-29

Pinworm

Treatment

pyrantel pamoate suspension

- Single dose effectiveness
- Measuring cup included
- Doctor Recommended

FOR THE ENTIRE FAMILY (AGES 2 & UP)

1 fl. oz. (30 ml)



PINWORM TREATMENT

pyrantel pamoate suspension

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:46122-825 |
| Route of Administration | ORAL | | |

| Active Ingredient/Active Moiety | | | | |
|--|--------------------------------|--|----------------------|--------------------|
| Ingredient Name | | | Basis of Strength | Strength |
| PYRANTEL PAMOATE (UNII: 81BK194Z5M) (PYRANTEL - UNII:4QIH0N49E7) | | | PYRANTEL | 144 mg in 1 mL |
| | | | | |
| Inactive Ingredients | | | | |
| Ingredient Name | | | | Strength |
| CITRIC ACID (UNII: 2968PHW8QP) | | | | |
| GLYCERIN (UNII: PDC6A3C0OX) | | | | |
| MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC) | | | | |
| METHYLCELLULOSE (UNII: Z944H5SN0H) | | | | |
| POVIDONE (UNII: FZ989GH94E) | | | | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | | | | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | | | | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | | | | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | | | | |
| SORBITOL (UNII: 506T60A25R) | | | | |
| WATER (UNII: 059QF0KO0R) | | | | |
| PROPYLPARABEN (UNII: Z8IX2SC1OH) | | | | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | | | | |
| METHYLPARABEN (UNII: A2I8C7HI9T) | | | | |
| | | | | |
| Product Characteristics | | | | |
| Color | yellow (pale yellow to yellow) | | Score | |
| Shape | | | Size | |
| Flavor | BANANA | | Imprint Code | |
| Contains | | | | |
| | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:46122-825-29 | 1 in 1 CARTON | 01/21/2026 | |
| 1 | | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |
| | | | | |
| Marketing Information | | | | |
| Marketing Category | | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug | | M024 | 01/21/2026 | |

Labeler - AMERISOURCE BERGEN (007914906)

Registrant - BIONPHARMA INC. (079637826)

Establishment

| Name | Address | ID/FEI | Business Operations |
|--|---------|-----------|------------------------|
| OrBion Pharmaceuticals Private Limited | | 854403569 | manufacture(46122-825) |

Revised: 1/2026

AMERISOURCE BERGEN