

IBUPROFEN AND PSEUDOEPHEDRINE HYDROCHLORIDE- ibuprofen and pseudoephedrine hydrochloride tablet, coated
Rite-Aid

IBUPROFEN & PSEUDOEPHEDRINE HCl TABLETS, USP

Drug Facts

Active ingredients (in each caplet)	Purposes
Ibuprofen, USP 200 mg (NSAID)*	Pain reliever/fever reducer
Pseudoephedrine HCl, USP 30 mg	Nasal decongestant

* nonsteroidal anti-inflammatory drug

Uses

temporarily relieves these symptoms associated with the common cold or flu:

- headache
- fever
- sinus pressure
- nasal congestion
- minor body aches and pains

Warnings

Allergy alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other product that contains pseudoephedrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain

- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years of age and over:
 - take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used.
 - do not use more than 6 caplets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use

Other information

- store at 20 - 25° C (68 - 77° F). Avoid excessive heat above 40° C (104° F).
- read all warnings and directions before use. Keep carton.

Inactive ingredients

acacia, calcium carbonate, carnauba wax, confectioner's sugar, corn starch, croscarmellose sodium, crospovidone, FD&C Blue no. 2 Aluminum Lake, FD&C Red no. 40 Aluminum Lake, FD&C Yellow no. 6 Aluminum Lake, gelatin, guar gum, hydrogenated vegetable oil, hydroxypropyl cellulose, iron oxide black, kaolin, polyethylene glycol, powdered cellulose, povidone, pregelatinized starch, propylene glycol, shellac, silicon dioxide, sodium benzoate, sucrose, talc, titanium dioxide, white wax

Questions?

call **1-800-406-7984**

**DISTRIBUTED BY: RITE AID
30 HUNTER LANE, CAMP HILL, PA 17011**

PRINCIPAL DISPLAY PANEL - 200 mg/30 mg Caplet Blister Pack Carton

NDC 11822-0084-6

†Compare to the active ingredients in Advil® Cold & Sinus

NON-DROWSY

COLD & SINUS RELIEF

IBUPROFEN, USP 200 mg &
PSEUDOEPHEDRINE HCl, USP 30 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)*
NASAL DECONGESTANT

Relieves sinus pressure,
Nasal congestion &
Fever

*nonsteroidal
anti-inflammatory drug

Actual Size

40
COATED CAPLETS††
(††OVAL-SHAPED TABLETS)

5224961

5224961

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, can increase the risk of heart attack, heart failure, and stroke. These risks are higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if:

- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic
- you are taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- do not take more than directed
- ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition
- under a doctor's care for any serious condition
- take any other product that contains pseudoephedrine or any other nasal decongestant
- decrease this benefit of aspirin
- taking any other drug

Drug Facts (continued)

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - stomach pain that does not get better
 - feel faint
 - vomit blood
 - have bloody or black stools
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

Pregnant or breast-feeding: ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- ask a doctor if you are under a doctor's care for any serious condition
- do not use more than 6 caplets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use

Drug Facts

Active Ingredients (in each caplet) Purposes

Ibuprofen, USP 200 mg (NSAID)*: Pain reliever/fever reducer
 Pseudoephedrine HCl, USP 30 mg: Nasal decongestant
 Nonsteroidal anti-inflammatory drug

Uses temporarily relieves these symptoms associated with the common cold or flu:

- nasal congestion
- fever
- headache
- sinus pressure
- minor body aches and pains

Warnings

Warning alert: Ibuprofen may cause a severe allergic reaction, which may be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

aspirin in people allergic to aspirin. Symptoms may include: hives, skin redness, rash, facial swelling, asthma (wheezing), shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed

Drug Facts (continued)

Directions

- do not use more than directed
- ask a doctor if you are under a doctor's care for any serious condition
- do not use more than 6 caplets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use

Compare to the active ingredients in Advil® Cold & Sinus

NON-DROWSY COLD & SINUS RELIEF

IBUPROFEN, USP 200 mg & PSEUDOEPHEDRINE HCl, USP 30 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)* NASAL DECONGESTANT

Relieves sinus pressure, Nasal congestion & Fever

Actual Size

40 COATED CAPLETS††
(††OVAL-SHAPED TABLETS)



*nonsteroidal anti-inflammatory drug

Expiration Date: _____
Batch No. _____

Non Varnish Area

Drug Facts (continued)

Other Information

- store at 20° - 25° C (68° - 77° F). Avoid excessive heat above 40° C (104° F).
- read all warnings and directions before use. Keep carton.

Inactive ingredients

acacia, calcium carbonate, carnuba wax, confectioner's sugar, corn starch, croscarmellose sodium, crospovidone, FD&C Blue no. 2, Aluminum Lake, FD&C Red no. 40, Aluminum Lake, FD&C Yellow no. 6, Aluminum Lake, gelatin, guar gum, hydrogenated vegetable oil, hydroxypropyl cellulose, iron oxide black, kaolin, poly(vinylpyrrolidone), powdered cellulose, povidone, polygalactulose starch, propylene glycol, she'llac, silicon dioxide, sodium benzoate, sucrose, talc, titanium dioxide, white wax

Questions? call 1-800-405-7964

TAMPER EVIDENT: DO NOT USE IF BUBBLER UNITS ARE LOOSE, BROKEN OR SHOW ANY SIGNS OF TAMPERING.

DISTRIBUTED BY: RTE AID
 30 HUNTER LANE,
 CAMP HILL, PA 17011
 www.rteaid.com

SCITRACKION SUBANTER
 It's not just a name. It's a commitment to help you feel better.

† All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Advil® Cold & Sinus.

0350487 R1021



5224961

IBUPROFEN AND PSEUDOEPHEDRINE HYDROCHLORIDE

ibuprofen and pseudoephedrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0084
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (120 .MU.M) (UNII: 68401960MK)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GUAR GUM (UNII: E89I1637KE)	
GLYCERYL TRISTEARATE (UNII: P6OCJ2551R)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
KAOLIN (UNII: 24H4NWX5CO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	

Product Characteristics

Color	brown	Score	no score
Shape	OVAL	Size	14mm

Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0084-6	1 in 1 CARTON	02/01/2018	
1		40 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-0084-5	1 in 1 CARTON	02/01/2018	
2		20 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA074567	02/01/2018		

Labeler - Rite-Aid (014578892)

Registrant - OHM LABORATORIES INC. (184769029)

Establishment			
Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(11822-0084)