

**POLYETHYLENE GLYCOL 3350- polyethylene glycol 3350 powder, for solution**  
**YYBA Corp**

-----

**Camber PEG 3350**

**ACTIVE INGREDIENT (IN EACH DOSE)**

**(Packet Only)**

Polyethylene Glycol 3350, 17 g

**PURPOSE**

Osmotic Laxative

**USE**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

**WARNINGS**

**Allergy alert: Do not use if you are allergic to polyethylene glycol**

**DO NOT USE**

if you have kidney disease, except under the advice and supervision of a doctor

**ASK A DOCTOR BEFORE USE IF YOU HAVE**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE**

taking a prescription drug

**WHEN USING THIS PRODUCT**

you may have loose, watery, more frequent stools

**STOP USE AND ASK A DOCTOR IF**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.

- you get diarrhea
- you need to use a laxative for longer than 1 week

## **IF PREGNANT OR BREAST-FEEDING**

ask a health professional before use.

## **KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a Poison Control Center right away.

## **DIRECTIONS**

### **(Bottle Only)**

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - use once a day
  - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficult swallowing
- ensure that the powder is fully dissolved before drink:iog
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

### **(Packet Only)**

- **do not take more than directed unless advised by your doctor**
- adults and children 17 years of age and older:
  - use once a day
  - stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficult swallowing
- ensure that the powder is fully dissolved before drink:iog
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

## **OTHER INFORMATION**

- store at 20° to 25°C (68° to 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

## **INACTIVE INGREDIENTS**

none

## **QUESTIONS OR COMMENTS**

call

1-866-933-6337

## **PRINCIPAL DISPLAY PANEL**

Polyethylene glycol 3350 powder for oral solution carton label

Cut here to open



NDC 73581-000-80

100%  
SATISFACTION  
GUARANTEED

# LAXATIVE

**OSMOTIC POWDER  
TO SOLUTION**

**Polyethylene  
Glycol 3350**

UNFLAVORED  
GRIT  
FREE



- ✓ Relieves Occasional Constipation & Irregularity
- ✓ 40 single serve doses (individually wrapped)
- ✓ Softens Stool and Increases Frequencies of Bowel Movements

NET WT 0.5 OZ (17 g)

This package is NOT child-resistant



## Drug Facts

**Active ingredient (in each dose).....Purpose**  
Polyethylene Glycol 3350, 17 g.....Osmotic Laxative

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use if you have kidney disease, except under the advice and supervision of a doctor**

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are taking a prescription drug**

**When using this product you may have loose, watery, more frequent stools**

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

## Drug Facts (continued)

### Directions

- do not take more than directed unless advised by your doctor
- adults and children 17 years of age and older:
  - use once a day
  - stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20° to 25°C (68° to 77°F)
- tamper-evident: do not use if foil is open or broken

**Inactive ingredients** none

**Questions or comments?** call 1-888-933-6337

## Not For Individual Sale

- Unflavored
- Sugar Free
- Dissolves in Any Beverage

Mfg. Lic. No.:

Distributed by: Wellspring

Airmont, NY 10952 USA.



Polyethylene glycol 3350 powder for oral solution carton label



# POLYETHYLENE GLYCOL 3350

polyethylene glycol 3350 powder, for solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73581-913
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73581-913-40	40 in 1 CARTON	01/01/2026	
1		17 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214990	01/01/2026	

**Labeler** - YYBA Corp (006339772)

**Registrant** - YYBA Corp (006339772)

Revised: 1/2026

YYBA Corp