

**LANSOPRAZOLE- lansoprazole capsule, delayed release
LEADER/ Cardinal Health 110, Inc.**

Lansoprazole Delayed Release Capsules USP

Active ingredient (in each capsule)

Lansoprazole USP, 15 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- do not use if you are allergic to lansoprazole
- lansoprazole may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a healthcare professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (**1-800-222-1222**).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20 - 25°C (68° - 77° F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients

D&C Red 28, FD&C Blue No.1, FD&C Green 3, FD&C Red 40, gelatin, hydroxypropyl cellulose, iron oxide black, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol 6000, polysorbate 80, sodium lauryl sulphate, starch (corn), sucrose, sugar spheres, talc, titanium dioxide

Questions or comments?

call **1-888-375-3784**

Tips For Managing Heartburn

Tips For Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

PRINCIPAL DISPLAY PANEL- Carton Label

Lansoprazole Delayed- Release Capsules USP, 15 mg



24 Hour
Lansoprazole
Delayed-Release Capsules USP, 15 mg
Acid Reducer

1 BOTTLE INSIDE

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.



1 BOTTLE INSIDE
14 CAPSULES TOTAL

Actual Size

WARNING: Do not use if foil seal under cap is broken with "hatched" for your protection" or black band around center of capsule is broken or missing. SOURCE DATA: Read the directions and warnings before use. Keep the carton. It contains important information.

Drug Facts

Active ingredient (in each capsule)
Lansoprazole USP, 15 mg.

Purpose
Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- do not use if you are allergic to lansoprazole
- lansoprazole may cause severe skin reactions.

Symptoms may include:

- skin redness
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss

Drug Facts (continued)

- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are:

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- if you take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Drug Facts (continued)

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° - 25°C (68° - 77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients DSC Red 28, FD&C Blue No. 1, FD&C Green 3, FD&C Red 40, gelatin, hydroxypropyl cellulose, iron oxides black, five substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol 6000, polyborate 80, sodium lauryl sulfate, starch (corn), sucrose, sugar spheres, talc, titanium dioxide

Questions or comments?
call 1-888-375-3784

24 Hour
Lansoprazole
Delayed-Release Capsules USP, 15 mg
Acid Reducer

14 CAPSULES
One 14-Day Course of Treatment

Treats Frequent Heartburn
May Take 1 to 4 Days for Full Effect

COMPARE TO PREVACID® 24HR
active ingredient*

100% Money Back Guarantee



*Actual Size

LEADER?

100% Money Back Guarantee

©2013 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the LEADER LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

All LEADER™ Brand Products Have A 100% Money Back Guarantee

Return to place of purchase if not satisfied.

DECEMBER 2013
DISTRIBUTED BY CARDINAL HEALTH
KANSAS CITY, MO 64101
www.myleader.com 1-800-280-6335
Made in India

LOT
EXP

LOT & EXP

1 BOTTLE INSIDE
14 CAPSULES TOTAL

Actual Size

WARNING: Do not use if foil seal under cap is broken with "hatched" for your protection" or black band around center of capsule is broken or missing. SOURCE DATA: Read the directions and warnings before use. Keep the carton. It contains important information.



100% Money Back Guarantee

Return to place of purchase if not satisfied.

100% Money Back Guarantee

Return to place of purchase if not satisfied.

PRINCIPAL DISPLAY PANEL - Container Label

NDC 70000-0612-1
24 Hour
Lansoprazole
 Delayed-Release Capsules USP, 15 mg
 Acid Reducer

Treats Frequent Heartburn
 May Take 1 to 4 Days for Full Effect
14 CAPSULES
 One 14-Day Course of Treatment

LEADER

IMPORTANT: This label does not contain full product information. See carton for complete information. Read the directions and warnings before use. Retain carton for reference.

TAMPER EVIDENT: Do not use if foil seal under cap printed with "sealed for your protection" or black band around center of capsule is broken or missing.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

DIST. BY CAH DUBLIN OH, 43017
 www.myleader.com 1-800-200-6315
 Made in India ©2022 Cardinal Health

100% Money Back Guarantee
 Return to place of purchase if not satisfied.
CIN 5795448 REV. 9/22
150089713
 LOT/EXP

Unvarnished Area
 PEEL HERE

Reseal Area

Directions (continued)

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- You may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

CONTINUED

Hinge

Hinge

Directions (continued)

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information
- store at 20° - 25°C (68° - 77° F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use
- if pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments?
 call 1-888-375-3784

LANSOPRAZOLE			
lansoprazole capsule, delayed release			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0612(NDC:43598-109)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Lansoprazole (UNII: 0K5C5T2QPG) (Lansoprazole - UNII:0K5C5T2QPG)	Lansoprazole	15 mg	
Inactive Ingredients			
Ingredient Name	Strength		
D&C RED NO. 28 (UNII: 767IP0Y5NH)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			

FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 2165RE0K14)	
MAGNESIUM CARBONATE (UNII: 0E53J927NA)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BLUE (opaque cyan colored cap) , PINK (opaque pink colored body)	Score	no score
Shape	CAPSULE	Size	3mm
Flavor		Imprint Code	RDY;398
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0612-1	1 in 1 CARTON	10/26/2022	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:70000-0612-3	3 in 1 CARTON	10/26/2022	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:70000-0612-2	2 in 1 CARTON	10/26/2022	
3		14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202194	10/26/2022	

Labeler - LEADER/ Cardinal Health 110, Inc. (063997360)