

NAPROXEN SODIUM - naproxen sodium tablet, coated tablet
MARKSANS PHARMA LIMITED

NAPROXEN SODIUM- naproxen sodium tablet,coated

Active ingredient(s)

For Tablets:

Active ingredient (in each tablet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

For Caplets:

Active ingredient (in each caplet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/ fever reducer

Uses

temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold

temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

you have difficulty swallowing

it feels like the pill is stuck in your throat

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

For Tablets:

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none">• take 1 tablet every 8 to 12 hours while symptoms last• for the first dose you may take 2 tablets within the first hour
--	---

	<ul style="list-style-type: none"> • do not exceed 2 tablets in any 8- to 12-hour period • do not exceed 3 tablets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none"> • ask a doctor

For Caplets:

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"> • take 1 caplet every 8 to 12 hours while symptoms last • for the first dose you may take 2 caplets within the first hour • do not exceed 2 caplets in any 8- to 12-hour period • do not exceed 3 caplets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none"> • ask a doctor

Other information

For Tablets:

- **each tablet contains:** sodium 20 mg
- store at 20-25°C (68-77°F).
- avoid high humidity and excessive heat above 40°C (104°F).

For Caplets:

- **each caplet contains:** sodium 20 mg
- store at 20-25°C (68-77°F).
- avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD&C blue #2 Lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide

Questions or comments?

call toll free **1-877-376-4271**

Manufactured for:

Time-Cap Labs, Inc.

7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:

Marksans Pharma Ltd.

Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, Goa - 403722, India.

Mfg. Lic. No.: GO/DRUGS/515

Principal Display Panel

NDC 25000-141-43
Naproxen Sodium Tablets, USP 220 mg
10's count bottle label

NDC 25000-141-43

Naproxen Sodium Tablets, USP
220 mg (NSAID)

Pain reliever / fever reducer

STRENGTH TO LAST 12 HOURS

10 Tablets



Marksans

See Carton for Full information. **Warnings**
Allergy alert : Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. if an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning** : This

LOT :

EXP :

R504/22

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others) ■ have 3 or more alcoholic drinks every day while using this product. ■ take more or for a longer time than

directed. **Heart attack and stroke warning**: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use if**

you have ever had an allergic reaction to any other pain reliever/fever reducer. **Ask a doctor or pharmacist before use if you are** taking any other drug. **Stop use and ask a doctor** if you experience any of the following

signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or

side of body, slurred speech, leg swelling), or if pain or fever gets worse. **If pregnant or breast feeding**, ask a health professional before use. **Keep out of reach of children**. In case of overdose, get medical help or contact a Poison

Control Center right away (1-800-222-1222).

Directions Adults and children 12 years and older: take 1 tablet every 8 to 12 hours while symptoms last. For the first dose you may take 2

exceed 2 tablets in any 8- to 12-hour period. Do not exceed 3 tablets in a 24-hour period. Children under 12 years: ask a doctor

Questions or comments?
call toll free 1-877-376-4271

Do not use if seal under cap imprinted with "SEALED for your PROTECTION" is broken or missing

tablets within the first hour. Do not

Manufactured for:

Time-Cap Labs, Inc.

7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:

Marksans Pharma Ltd.


Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-141-43
Naproxen Sodium Tablets, USP 220 mg
10's count carton label



NDC 25000-141-02
 Naproxen Sodium Tablets, USP 220 mg
 24's count bottle label



product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others) ■ have 3 or more alcoholic drinks every day while using this product. ■ take more or for a longer time than

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use if**

you have ever had an allergic reaction to any other pain reliever/fever reducer. **Ask a doctor or pharmacist before use if you are** taking any other drug. **Stop use and ask a doctor** if you experience any of the following

signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or

side of body, slurred speech, leg swelling), or if pain or fever gets worse. **If pregnant or breast feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison

Control Center right away (1-800-222-1222).

Directions Adults and children 12 years and older: take 1 tablet every 8 to 12 hours while symptoms last. For the first dose you may take 2 tablets within the

first hour. Do not exceed 2 tablets in any 8- to 12-hour period. Do not exceed 3 tablets in a 24-hour period. Children under 12 years: ask a doctor.

Questions or comments?
call toll free 1-877-376-4271

Do not use if seal under cap imprinted with "SEALED for your PROTECTION" is broken or missing

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale

Manufactured by:
Marksans Pharma Ltd.
Plot No. 1-82, 1-83

7 Michael Avenue, Farmingdale,
NY 11735, USA

Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-141-02
Naproxen Sodium Tablets, USP 220 mg
24's count carton label

Drug Facts (Continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- taking a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

Drug Facts (Continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and Children 12 years and older:

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8-to-12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years:

- ask a doctor

Other information

- each tablet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide.

Questions or comments?
call toll free 1-877-376-4271

Do not use if carton is open or seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Manufactured by:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-141-02

**Naproxen Sodium Tablets, USP
220 mg (NSAID)**

Pain reliever / fever reducer

STRENGTH TO LAST 12 HOURS

24 Tablets

Marksans

Drug Facts

Active ingredient (in each tablet) Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*

Purposes Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

NDC 25000-141-14
 Naproxen Sodium Tablets, USP 220 mg
 1000's count bottle label

NDC 25000-141-14

Naproxen Sodium Tablets, USP 220 mg (NSAID)

Pain reliever / fever reducer

STRENGTH TO LAST 12 HOURS

1000 Tablets



Marksans

Drug Facts

Active ingredient (in each tablet)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis ■ muscular aches
 - backache ■ menstrual cramps ■ headache
 - toothache ■ the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include :

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

LOT :

EXP :

R1104/22



Drug Facts (Continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you :

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Drug Facts (Continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain ■ trouble breathing ■ weakness in one part or side of body
 - slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may

Drug Facts (Continued)

cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and Children
12 years and older:

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8-to 12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years:

- ask a doctor

Other information

- each tablet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, ▶

Drug Facts (Continued)

FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide.

Questions or comments?

call toll free 1-877-376-4271

Do not use if seal under cap imprinted with
"SEALED for YOUR PROTECTION"
is broken or missing.

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-144-14

Naproxen Sodium Tablets, USP 220 mg (Capsule-shaped tablet)
1000's count bottle label

NDC 25000-144-14

Naproxen Sodium Tablets, USP 220 mg (NSAID)

Pain reliever / fever reducer

STRENGTH TO LAST 12 HOURS

1000 Tablets (oval-shaped)



Marksans

Drug Facts

<i>Active ingredient (in each caplet)</i>	<i>Purposes</i>
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis ■ muscular aches
 - backache ■ menstrual cramps ■ headache
 - toothache ■ the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include :

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

LOT :

EXP :

R1104/22



Drug Facts (Continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you :

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Drug Facts (Continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain ■ trouble breathing ■ weakness in one part or side of body
 - slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may

Drug Facts (Continued)

Drug Facts (Continued)

cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and Children
12 years and older:

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8-to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years:

- ask a doctor

Other information

- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium,

FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide.

Questions or comments?

call toll free 1-877-376-4271

Do not use if seal under cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-144-43

Naproxen Sodium Tablets, USP 220 mg (Capsule-shaped tablet)

10's count bottle label


NDC 25000-144-43

**Naproxen Sodium Tablets, USP
220 mg (NSAID)**

Pain reliever / fever reducer

STRENGTH TO LAST 12 HOURS

10 Tablets (oval-shaped)



Marksans

See Carton for Full information. Warnings

Allergy alert : Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. if an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning :** This

LOT : _____

EXP : _____

R504/22

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others) ■ have 3 or more alcoholic drinks every day while using this product. ■ take more or for a longer time than

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use if**

you have ever had an allergic reaction to any other pain reliever/fever reducer. **Ask a doctor or pharmacist before use if you are** taking any other drug. **Stop use and ask a doctor** if you experience any of the following

signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or

side of body, slurred speech, leg swelling), or if pain or fever gets worse. **If pregnant or breast feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison

Control Center right away (1-800-222-1222).

Directions Adults and children 12 years and older: take 1 caplet every 8 to 12 hours while symptoms last. For the first dose you may take 2 caplets within the

first hour. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period. Children under 12 years: ask a doctor.

Questions or comments?
call toll free **1-877-376-4271**

Do not use if seal under cap imprinted with "SEALED for your PROTECTION" is broken or missing

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-144-43

Naproxen Sodium Tablets, USP 220 mg (Capsule-shaped tablet)

10's count carton label



NDC 25000-144-02
 Naproxen Sodium Tablets, USP 220 mg (Capsule-shaped tablet)
 24's count bottle label





people allergic to aspirin. Symptoms may include:

help fight away. **bleeding warning** : This

LOT :

EXP :

R1104/22

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others) ■ have 3 or more alcoholic drinks every day while using this product. ■ take more or for a longer time than

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use if**

you have ever had an allergic reaction to any other pain reliever/fever reducer. **Ask a doctor or pharmacist before use if you are** taking any other drug. **Stop use and ask a doctor** if you experience any of the following

signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or

side of body, slurred speech, leg swelling), or if pain or fever gets worse. **If pregnant or breast feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison

Control Center right away (1-800-222-1222).

Directions Adults and children 12 years and older: take 1 caplet every 8 to 12 hours while symptoms last. For the first dose you may take 2 caplets within the

first hour. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period. Children under 12 years: ask a doctor.

Questions or comments? call toll free **1-877-376-4271**

Do not use if seal under cap imprinted with "SEALED for your PROTECTION" is broken or missing

Manufactured for:
Time-Cap Labs, Inc.
 7 Michael Avenue, Farmingdale,
 NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
 Plot No. L-82, L-83,
 Verna Indl. Estate,
 Verna, GOA - 403 722, India.
 Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-144-02
 Naproxen Sodium Tablets, USP 220 mg (Capsule-shaped tablet)
 24's count carton label

: dX3
: LOT



NDC 25000-144-02

Naproxen Sodium Tablets, USP
220 mg (NSAID)

Pain reliever / fever reducer

STRENGTH TO LAST 12 HOURS

24 Tablets (oval-shaped)

Marksans



: dX3
: LOT

Drug Facts (Continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

Drug Facts (Continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

<p>Adults and Children 12 years and older:</p> <ul style="list-style-type: none"> ■ take 1 caplet every 8 to 12 hours while symptoms last ■ for the first dose you may take 2 caplets within the first hour ■ do not exceed 2 caplets in any 8-to 12-hour period ■ do not exceed 3 caplets in a 24-hour period 	<p>Children under 12 years:</p> <ul style="list-style-type: none"> ■ ask a doctor
--	--

Other information

- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)

Drug Facts

Active ingredient (in each caplet)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (Continued)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide.

Questions or comments?
 call toll free 1-877-376-4271

Do not use if carton is open or seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Manufactured for:
Time-Cap Labs, Inc.
 7 Michael Avenue, Farmingdale,
 NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
 Plot No. L-82, L-83,
 Verna Indl. Estate,
 Verna, GOA - 403 722, India.
 Mfg. Lic. No. : GO/DRUGS/515

NAPROXEN SODIUM

naproxen sodium tablet, coated tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:25000-144
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BLUE	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	144
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-144-14	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/04/2011	
2	NDC:25000-144-20	4 in 1 BOX	05/04/2011	
2		6500 in 1 BAG; Type 0: Not a Combination Product		
3	NDC:25000-144-30	6 in 1 BOX	05/04/2011	
3		6500 in 1 BAG; Type 0: Not a Combination Product		
4	NDC:25000-	100 in 1 BOX	05/04/2011	

4	144-02	100 in 1 BOX	05/04/2011	
4		1 in 1 CARTON		
4		24 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:25000-144-43	10 in 1 BOTTLE; Type 0: Not a Combination Product	05/04/2011	
6	NDC:25000-144-98	1 in 1 BOX	05/04/2011	
6		32165 in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090545	05/04/2011	

NAPROXEN SODIUM

naproxen sodium, coated tablets tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:25000-141
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	BLUE	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	141
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-141-14	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/28/2011	
2	NDC:25000-141-26	2 in 1 BOX	05/28/2011	
2		1600 in 1 BAG; Type 0: Not a Combination Product		
3	NDC:25000-141-30	6 in 1 BOX	05/28/2011	
3		6500 in 1 BAG; Type 0: Not a Combination Product		
4	NDC:25000-141-02	160 in 1 BOX	05/28/2011	
4		1 in 1 CARTON		
4		24 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:25000-141-43	10 in 1 BOTTLE; Type 0: Not a Combination Product	05/28/2011	
6	NDC:25000-141-98	1 in 1 BOX	05/28/2011	
6		32165 in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090545	05/28/2011	

Labeler - MARKSANS PHARMA LIMITED (925822975)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	MANUFACTURE(25000-141, 25000-144)

Revised: 6/2023

MARKSANS PHARMA LIMITED