

**ITRAFUNGOL- itraconazole injection, solution**  
**Elanco US Inc.**

-----  
**Itrafungol™**  
**(itraconazole oral solution)**

10 mg/ml

Antifungal for oral use in cats only

**Caution:**

Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

**Description:**

ITRAFUNGOL oral solution is a yellow to slightly amber, clear solution containing the active ingredient, itraconazole, at 10 mg/mL.

**Indication:**

ITRAFUNGOL oral solution is indicated for the treatment of dermatophytosis caused by *Microsporum canis* in cats.

**Dosage and Administration:**

The solution should be administered orally using the enclosed graduated dosing syringe.

The daily dosage is 5 mg/kg (0.5 mL/kg) body weight administered once daily on alternating weeks for 3 treatment cycles. Cats are treated during weeks 1, 3, and 5, and left untreated during weeks 2 and 4.

7 days	7 days	7 days	7 days	7 days
Daily treatment	No treatment	Daily treatment	No treatment	Daily treatment

Each line on the dosing syringe represents 0.05 mL of oral solution.

**Table 1: Dose Table for ITRAFUNGOL**

Weight of Cat	Volume of ITRAFUNGOL
0.5 lb	0.1 mL
1.0 lb	0.2 mL
1.5 lb	0.35 mL
2.0 lb	0.45 mL
2.5 lb	0.55 mL
3.0 lb	0.7 mL
3.5 lb	0.8 mL
4.0 lb	0.9 mL
4.5 lb	1.0 mL
5.0 lb	1.15 mL
6.0 lb	1.35 mL

7.0 lb	1.6 mL
8.0 lb	1.8 mL
9.0 lb	2.0 mL
10.0 lb	2.25 mL
11.0 lb	2.5 mL
12.0 lb	2.7 mL
13.0 lb	3.0 mL

The solution should be administered orally using the enclosed graduated dosing syringe. Keep the bottle upright and insert the dosing syringe through the opening of the top of the bottle (Figure 1). Do not invert the bottle (Figure 2). Fill the syringe by pulling the plunger until it reaches the graduation corresponding to the correct mL dose as indicated at the top of the syringe ring (Figure 3). Treat the cat by slowly and gently administering the liquid into the mouth, allowing the cat to swallow the product (Figure 4). For cats weighing more than 13.0 lbs, the total dose will need to be calculated and given over two doses as the dosing syringe only holds 3.0 mL of solution.

Fig. 1



Fig. 2



Fig. 3



Fig. 4



After dosing, the syringe should be removed from the bottle, rinsed and dried and the bottle cap should be screwed back on tightly.

#### **Contraindications:**

Do not administer to cats with hypersensitivity to itraconazole.

**Warnings:**

ITRAFUNGOL (itraconazole oral solution) has not been shown to be safe in pregnant cats (see Animal Safety section). ITRAFUNGOL should only be used in pregnant or lactating cats when the benefits outweigh the potential risks.

**User Safety Warnings:**

Not for use in humans. Keep this and all medications out of reach of children. Wash hands and exposed skin after use. In case of accidental contact with eyes, rinse thoroughly with water. In case of pain or irritation, seek medical advice. In case of accidental ingestion, rinse mouth with water and seek medical advice.

**Special precautions for person administering the veterinary product to the animal:**

*Microsporium canis* dermatophytosis is a zoonotic disease (a disease that can be transmitted from animals to humans); therefore consult a physician if a suspected lesion occurs on a human. Wear protective gloves when handling the animal during treatment or when cleaning the syringe. Wash hands and exposed skin after handling the animal.

ITRAFUNGOL has not been shown to be sporicidal; therefore in order to reduce zoonotic potential, environmental contamination, and to decrease course of the disease, topical and environmental treatment should also be utilized.

**Precautions:**

ITRAFUNGOL has been associated with renal changes found on histopathology that were not noted after an eight week recovery period (see Animal Safety). Use with caution in cats with renal dysfunction.

ITRAFUNGOL is metabolized by the liver (mainly CYP3A) and can cause elevated liver enzymes (see Animal Safety section). Use with caution in cats with impaired liver function. If clinical signs suggestive of liver dysfunction develop, treatment should be discontinued.

ITRAFUNGOL is a cytochrome p-450 inhibitor and may increase or prolong plasma concentrations of other drugs metabolized by this pathway, such as amitriptyline, amlodipine, benzodiazepines, buspirone, cisapride, corticosteroids, cyclosporine, ivermectin, and macrolide antibiotics.

Negative inotropic effects have been reported in literature when itraconazole was administered intravenously to dogs and healthy human volunteers. Cats suffering from heart disease should be carefully monitored during treatment.

**Adverse Reactions:**

In the laboratory effectiveness study, adverse reactions related to exposure to ITRAFUNGOL were primarily related to the gastrointestinal tract. Two ITRAFUNGOL-treated cats experienced transient hypersalivation during the dosing period. Vomiting was observed in 5 ITRAFUNGOL-treated cats (12.5%) during the dosing period compared to four cats (10%) in the control group. Diarrhea was observed in 9 ITRAFUNGOL-treated cats (22.5%) during the dosing period as compared to 7 cats (17.5%) in the control group. One ITRAFUNGOL-treated cat showed mild increases in alanine aminotransferase (ALT) and aspartate aminotransferase (AST) at the end of the dosing period. No related clinical signs were observed, and these values returned to normal by the end of the follow-up period. One cat in the ITRAFUNGOL-treated group was noted to have lip erythema and lip induration once during the study.

Field safety was evaluated in 266 cats randomized to receive itraconazole oral solution. Of the 266 cats that received at least one dose of itraconazole oral solution, adverse reactions included 35 cases (13%)

of one or more elevated hepatic enzymes and 8 cases (3%) of gastrointestinal upset, including decreased appetite, vomiting and/or diarrhea. Other infrequent adverse reactions included less than 3 cases each of somnolence, depression, and increased salivation.

For technical assistance or to report suspected adverse drug events, contact Elanco US Inc. at 1-888-545-5973.

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or on line at [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae)

### **Clinical Pharmacology:**

The mode of action of itraconazole is based on its highly selective binding ability to fungal cytochrome p-450 iso-enzymes. This inhibits the synthesis of ergosterol and affects membrane-bound enzyme function and membrane permeability. This effect is irreversible and causes structural degeneration of the fungal organism.

Itraconazole was absorbed rapidly following oral administration of ITRAFUNGOL (itraconazole oral solution) to laboratory cats. Compared to the fasted state, administration of ITRAFUNGOL with food results in slightly higher (1.3 fold) mean total itraconazole exposure (AUC), delay in median  $T_{max}$  (Fed 4 hours vs. Fasted 2 hours) and significant decrease (approximately 0.55 fold) in maximum plasma concentration ( $C_{max}$ ). ITRAFUNGOL can be administered with or without food. Itraconazole oral solution binds extensively to plasma proteins (> 99%), and distributes well to tissues. More than 30 metabolites are formed. Hydroxy-itraconazole is the parent metabolite and has antifungal activity. Excretion is rapid and primarily via the feces.

In cats, a single oral dose of 5 mg/kg results in a  $C_{max}$  of 0.525  $\mu$ /ml post dose at 2 hours ( $T_{max}$ ). The  $AUC_{0-24h}$  is 5.09  $\mu$ g.h/ml and the half-life in plasma is 12.1 hours. After repeated daily administration for seven days at 5 mg/kg/day, the  $C_{max}$  is doubled (1.05  $\mu$ g/ml), the  $AUC_{0-24h}$  is increased 3-fold (15.4  $\mu$ g.h/ml) and the plasma half-life is increased to 36 hours.

In the therapeutic treatment schedule, itraconazole is almost completely cleared from plasma after each wash-out period. The hydroxy-itraconazole remains near or below the quantification limit in feline plasma after a single dose of itraconazole oral solution at 5 mg/kg. However, after repeated daily doses of itraconazole oral solution at 5 mg/kg for one week, the hydroxy-itraconazole  $C_{max}$  of 0.059  $\mu$ g/ml was reached at 2 hours ( $T_{max}$ ). Itraconazole concentrations in cat's hair vary; an increase occurs during treatment to a median value of 3.0  $\mu$ g/g (mean 5.2  $\mu$ g/g) at the end of the third dosing week and concentrations drop slowly to 1.5  $\mu$ g/g (mean 1.9  $\mu$ g/g) at 14 days after final dosing. Concentrations of hydroxy-itraconazole in hair are insignificant.

### **Effectiveness:**

#### **Laboratory Study**

Effectiveness was demonstrated using ITRAFUNGOL (itraconazole oral solution) in a masked, placebo controlled laboratory study. Eighty cats were experimentally infected with *Microsporum canis* and treated with either ITRAFUNGOL or sterile water (control product) for the proposed therapeutic treatment schedule followed by a 4-week follow-up period. No topical therapy was used during this study. A statistical difference ( $P = 0.0003$ ) in mycological cure rate (defined as two consecutive negative mycological cultures) was demonstrated between cats treated with ITRAFUNGOL (24/40 or 60%) versus control (1/40 or 2.5%). Ninety percent of ITRAFUNGOL-treated cats (36/40) achieved at least one negative culture by the end of the study. Improvement was seen in inoculation site erythema and skin thickening by Day 7 and in crusts and scales by Day 14. By the end of the study, 98% of ITRAFUNGOL-treated cats had complete resolution of all clinical lesions, compared to 15% in the control group. Wood's lamp cure (defined as no fluorescence at the base and mid-shaft of the hair) in the ITRAFUNGOL-treated group (39/40 or 97.5%) was higher compared to the control group (6/40 or

15%). Itraconazole MICs indicative of susceptibility were obtained in *M. canis* isolates from the two cats unsuccessfully treated with ITRAFUNGOL.

## **Field Study**

A masked, positive-controlled, multi-site field study was conducted in client-owned cats in Europe. In this study, 514 cats diagnosed with dermatophytosis were randomly administered itraconazole oral solution or an active control. Cats received a daily dose of either itraconazole oral solution for three alternating weeks plus a placebo tablet once daily for 5 consecutive weeks, or a placebo solution for three alternating weeks plus the active control once daily for five weeks. Success was evaluated on clinical cure, which was noted with a complete resolution of all clinical lesions. Four weeks after the end of treatment, 175 (83%) out of 207 cats treated with itraconazole oral solution were clinically cured.

## **Animal Safety:**

### **Margin of Safety Study with Recovery**

In a margin of safety study, ITRAFUNGOL (itraconazole oral solution) was administered orally to 9-10 week old healthy kittens once daily at 0X (saline control), 1X (5 mg/kg), 3X (15 mg/kg), and 5X (25 mg/kg) the therapeutic dose for 17 alternating weeks (9 total weeks of dosing) followed by an 8 week recovery period. Hypersalivation during or immediately following dosing, vomiting, and loose stool were the most frequent abnormal clinical observations related to treatment with ITRAFUNGOL. Hypersalivation was limited to the 3X and 5X groups and was observed in every dosing cycle. Vomiting was noted at similar levels in the control, 1X and 3X groups; however, it occurred approximately twice as often in the 5X group. Mild gingival bleeding and perioral irritation (patchy alopecia and erythema) was noted in cats in the 3X and 5X groups. Food consumption was consistently higher throughout the study in the control group than the ITRAFUNGOL group. The control group gained more weight during the study than the groups administered ITRAFUNGOL. Mild elevations in ALT were sporadically noted in all groups; however, the number of affected cats increased with the higher doses (two cats in the control group, two cats in the 1X group, three cats in the 3X group, and four cats in the 5X group). In most cats, ALT values peaked just above the upper limit of the reference range and were continuing to trend upward or were elevated yet stable at the end of the study. One cat in the 5X group exhibited inappetence progressing to anorexia, dehydration and vomiting during the first dosing cycle. This cat had repeated episodes of inappetence during the second and third dosing cycles. This cat also had markedly elevated ALT and AST values on Day 36 (693 U/L and 283 U/L, respectively), was treated with minimal supportive care and recovered to complete the study.

### **Margin of Safety Study**

In a margin of safety study, ITRAFUNGOL (itraconazole oral solution) was administered orally to healthy adult cats once daily at 0X (saline control), 1X (5 mg/kg), 3X (15 mg/kg), and 5X (25 mg/kg) the therapeutic dose for 17 alternating weeks (9 total weeks of dosing) with no recovery period. Hypersalivation was the most frequent abnormal clinical observation related to treatment with ITRAFUNGOL and the incidence increased with the higher doses. One cat in group 4 (5X; Cat #26302) lost 22% of its body weight and had a number of episodes of vomiting, salivation, and anorexia during the treatment period. This cat also had renal lesions found on histopathology. Increasing trends were noted in ALT, AST, and creatinine values in some cats administered ITRAFUNGOL as compared to baseline values. Abnormal renal findings included proximal convoluted tubule acute degeneration in 3 cats in the 1X group and 3 cats in the 5X group; one 5X cat (cat #26302) also had proximal convoluted tubule marked pallor and focal mononuclear cell infiltration in the kidneys. In the lungs, one 3X group cat and five 5X cats had intra-alveolar foamy macrophages; five 5X group cats had intra-alveolar syncytial cells.

These histopathology findings are likely related to exposure to ITRAFUNGOL, specifically the

vehicle component hydroxypropyl- $\beta$ -cyclodextrin (HP $\beta$ CD). There were no corresponding adverse clinical effects noted on observation or on clinical pathology analysis. In addition, similar changes have been described in literature in other species exposed to HP $\beta$ CD and have been reported to be reversible.

### **Reproductive Safety**

In a study of 16 pregnant queens administered itraconazole oral solution at 5 mg/kg bodyweight for a total of 21 days (7 days on alternate weeks) during gestation or lactation, there was a high frequency of fetal resorption (partial and total), abnormal fetuses, and abnormal maternal behaviors. Confounding factors, such as infectious disease (*Chlamydia psittaci*) in some cats made it difficult to establish a definitive relationship between administration of itraconazole and the abnormal findings. However, the results of this study reveal potential reproductive safety risks and do not support the safe use of ITRAFUNGOL in pregnant queens.

### **Storage conditions:**

Store at 68-77°F (20-25°C). Excursions permitted between 59-86°F (15-30°C).

### **How supplied:**

ITRAFUNGOL (itraconazole oral solution) is available in a glass bottle containing 52 mL of oral solution, closed with a child resistant screw cap and packaged in a cardboard box that includes a package insert and a graduated dosing syringe.

Approved by FDA under NADA # 141-474

Manufactured for Elanco US Inc.

Greenfield, IN 46140

Version Date: September 2019

Itrafungol, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

**Elanco™**

### **Principal Display Panel - 52 mL Carton Label.**

NDC 58198-0796-4

**52 ml**

**Itrafungol™**

**(itraconazole  
oral solution)**

**Antifungal for oral  
use in cats only**

For the treatment of dermatophytosis  
caused by *Microsporum canis* in cats

Oral dosing syringe included

**10 mg/ml**

### **Caution:**

Federal (USA) law restricts this  
drug to use by or on the order  
of a licensed veterinarian.



**Principal Display Panel – 52 mL Bottle Label**

**52 ml**

**Itrafungol™**

**(itraconazole oral solution)**

## Antifungal for oral use in cats only

For the treatment of dermatophytosis caused by *Microsporum canis* in cats

## Oral dosing syringe included

### Caution:

Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

**10 mg/ml**

**Elanco**

**Itrafungol<sup>TM</sup>**  
**(itraconazole oral solution)**  
Antifungal for oral use in cats only

52 ml

For the treatment of dermatophytosis caused by *Microsporum canis* in cats

Oral dosing syringe included

Caution:  
Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

10 mg/ml

**Elanco**

Not for use in humans. Keep this and all medications out of the reach of children. For complete product information and warnings, see package insert.

Store at 68-77°F (20-25°C). Excursions permitted between 59-86°F (15-30°C).

Manufactured for Elanco US Inc. Greenfield, IN 46140

Approved by FDA under NADA # 141-474

Extra Print & Varnish Free Area If 3 x Variable Data is required

P16082488

Lot No.:

Exp.:

## ITRAFUNGOL

itraconazole injection, solution

### Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:58198-0796
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
itraconazole (UNII: 304NUG5GF4) (itraconazole - UNII:304NUG5GF4)	itraconazole	10 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
Hydroxypropyl Betadex (UNII: 1I96OHX6EK)	
Hydrochloric acid (UNII: QTT17582CB)	
Propylene glycol (UNII: 6DC9Q167V3)	
Sodium hydroxide (UNII: 55X04QC32I)	



<b>Saccharin sodium</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>Cherry</b> (UNII: BUC5I9595W)	
<b>Caramel</b> (UNII: T9D99G2B1R)	
<b>Water</b> (UNII: 059QF0K00R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58198-0796-4	1 in 1 BOX		
1		52 mL in 1 BOTTLE, GLASS		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141474	11/10/2016	

**Labeler** - Elanco US Inc. (966985624)

### Establishment

Name	Address	ID/FEI	Business Operations
SMS Pharmaceuticals Limited		677605406	API MANUFACTURE

### Establishment

Name	Address	ID/FEI	Business Operations
Lusomedicamenta Sociedade Tecnica Farmaceutica S.A		450630343	MANUFACTURE, PACK, LABEL

Revised: 4/2020

Elanco US Inc.