

STERILE WATER- water injection
Cardinal Health 107, LLC

Sterile Water for Injection, USP

DESCRIPTION

This preparation is designed solely for parenteral use only after addition to drugs that require dilution or must be dissolved in an aqueous vehicle prior to injection.

Sterile Water for Injection, USP is a sterile, nonpyrogenic preparation of water for injection which contains no bacteriostat, antimicrobial agent or added buffer and is supplied only in single dose containers to dilute or dissolve drugs for injection. For IV injection, add sufficient amount to a solute to make an approximately isotonic solution. pH 5.0 to 7.0.

Water for Injection, USP is chemically designated H₂O.

CLINICAL PHARMACOLOGY

Water is an essential constituent of all body tissues and accounts for approximately 70% of total body weight. Average normal adult daily requirement ranges from two to three liters (1 to 1.5 liters each for insensible water loss by perspiration and urine production).

Water balance is maintained by various regulatory mechanisms. Water distribution depends primarily on the concentration of electrolytes in the body compartments and sodium (Na⁺) plays a major role in maintaining physiologic equilibrium.

The small volume of fluid provided by Sterile Water for Injection, USP when used only as a pharmaceutical aid for diluting or dissolving drugs for parenteral injection, is unlikely to exert a significant effect on fluid balance except possibly in neonates or very small infants.

INDICATIONS AND USAGE

This parenteral preparation is indicated only for diluting or dissolving drugs for intravenous, intramuscular or subcutaneous injection, according to instructions of the manufacturer of the drug to be administered.

CONTRAINDICATIONS

Sterile Water for Injection must be made approximately isotonic prior to use.

WARNINGS

Intravenous administration of Sterile Water for Injection without a solute may result in hemolysis.

PRECAUTIONS

Do not use for intravenous injection unless the osmolar concentration of additives results in an approximate isotonic admixture.

Consult the manufacturer's instructions for choice of vehicle, appropriate dilution or volume for dissolving the drugs to be injected, including the route and rate of injection.

Inspect reconstituted (diluted or dissolved) drugs for clarity (if soluble) and freedom from unexpected precipitation or discoloration prior to administration.

Pregnancy

Animal reproduction studies have not been conducted with Sterile Water for Injection. It is also not known whether sterile water containing additives can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Sterile Water for Injection with additives should be given to a pregnant woman only if clearly needed.

Pediatric Use

Safety and effectiveness have been established in pediatric patients. However, in neonates or very small infants the volume of fluid may affect fluid and electrolyte balance.

Drug Interactions

Some drugs for injection may be incompatible in a given vehicle, or when combined in the same vehicle or in a vehicle containing benzyl alcohol. Consult with pharmacist, if available.

Use aseptic technique for single or multiple entry and withdrawal from all containers.

When diluting or dissolving drugs, mix thoroughly and use promptly. Do not store reconstituted solutions of drugs for injection unless otherwise directed by the manufacturer of the solute.

Do not use unless the solution is clear and seal intact. Do not reuse single-dose containers. Discard unused portion.

ADVERSE REACTIONS

Reactions which may occur because of this solution, added drugs or the technique of reconstitution or administration include febrile response, local tenderness, abscess, tissue necrosis or infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection and extravasation. If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate countermeasures, and if possible, retrieve and save the remainder of the unused vehicle for examination.

OVERDOSAGE

Use only as a diluent or solvent. This parenteral preparation is unlikely to pose a threat

of fluid overload except possibly in neonates or very small infants. In the event these should occur, re-evaluate the patient and institute appropriate corrective measures (see **WARNINGS, PRECAUTIONS** and **ADVERSE REACTIONS**).

DOSAGE AND ADMINISTRATION

The volume of the preparation to be used for diluting or dissolving any drug for injection is dependent on the vehicle concentration, dose and route of administration as recommended by the manufacturer.

This parenteral should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

How Supplied

Overbagged with 5 x 10 mL Single Dose Vials in each bag, NDC 55154-9588-5

Packaged in a plastic vial.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Single dose use. No preservative added.

Unused portion of vial should be discarded.

Use only if solution is clear and seal intact.



Lake Zurich, IL 60047

www.fresenius-kabi.com/us

Distributed By:

Cardinal Health

Dublin, OH 43017

L58749380424

45768H

Revised: April 2022

Package/Label Display Panel

STERILE WATER FOR INJECTION, USP

5 x 10 mL SINGLE-DOSE VIALS



F115

NDC 55154-9588-5

STERILE WATER FOR INJECTION, USP 5 x 10 mL SINGLE-DOSE VIALS

FOR DRUG DILUENT USE ONLY

Sterile, Nonpyrogenic
Preservative Free

Discard unused portion.

Do not administer intravenously unless rendered
nearly isotonic.

STORAGE: Store at 20° to 25° C (68° to 77° F) [See USP
Controlled Room Temperature].

RX ONLY

WARNING: This Unit Dose package is not child resistant
and is Intended for Institutional Use Only.
Keep this and all drugs out of the reach of children.

Fresenius Kabi
Lake Zurich, IL 60047
www.fresenius-kabi.com/us

Distributed by Cardinal Health
Dublin, OH 43017

L58749380424

STERILE WATER

water injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55154-9588(NDC:63323-185)
Route of Administration	INTRAMUSCULAR, SUBCUTANEOUS, INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WATER (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)	WATER	1 mL in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-9588-5	5 in 1 BAG	10/31/2024	
1		10 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA088400	10/31/2024	

Labeler - Cardinal Health 107, LLC (118546603)

Revised: 10/2024

Cardinal Health 107, LLC