

**TYLENOL COLD PLUS FLU MULTI-SYMPTOM LIQUID GELS DAY PLUS NIGHT-
acetaminophen, dextromethorphan hydrobromide and doxylamine succinate
Kenvue Brands LLC**

TYLENOL® Cold + Flu Multi-Symptom Liquid Gels Day + Night

Drug Facts

TYLENOL® Cold + Flu Multi-Symptom Liquid Gels Day

Active ingredient (in each capsule)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Pain reliever/fever reducer

Cough suppressant

Uses

- temporarily relieves these common cold/flu symptoms:
- minor aches and pains
- headache
- sore throat
- cough
- temporarily reduces fever

Warnings

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ **do not take more than directed (see overdose warning)**

| | |
|---------------------------------------|---|
| adults and children 12 years and over | ■ take 2 capsules every 4 hours while symptoms last ■ do not take more than 8 capsules in 24 hours |
| Children 4 to under 12 years | ask a doctor |
| Children under 4 years | do not use |

Other information

- store at 20° to 25°C (68° to 77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- protect from light
- **do not use if blister unit is torn or broken**

Inactive ingredients

FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, lecithin, mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Drug Facts

TYLENOL® Cold + Flu Multi-Symptom Liquid Gels Night

Active ingredient (in each capsule)

Active ingredient (in each capsule)

Acetaminophen 325 mg
Dextromethorphan HBr 15 mg
Doxylamine succinate 6.25 mg

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves these common cold/flu symptoms:
- minor aches and pains
- headache
- sore throat
- cough
- runny nose and sneezing
- temporarily reduces fever

Warnings

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

do not exceed recommended dose

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

- new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

| | |
|---------------------------------------|---|
| adults and children 12 years and over | ■ take 2 capsules every 6 hours while symptoms last ■ do not take more than 8 capsules in 24 hours |
| Children 4 to under 12 years | ask a doctor |
| Children under 4 years | do not use |

Other information

- store at 20° to 25°C (68° to 77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- protect from light
- **do not use if blister unit is torn or broken**

Inactive ingredients

D&C yellow no. 10, FD&C blue no. 1, gelatin, glycerin, lecithin, mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-141-03

TYLENOL ®

FOR ADULTS

COLD & FLU

**MULTI SYMPTOM
LIQUID GELS**

Acetaminophen,

Dextromethorphan HBr
Pain Reliever-Fever Reducer,
Cough Suppressant

DAY

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH

Actual Size

3X16 LIQUID GELS

Acetaminophen,

Dextromethorphan HBr, Doxylamine Succinate

Pain Reliever-Fever Reducer,
Cough Suppressant, Antihistamine

NIGHT

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH • RUNNY NOSE

Actual Size

3X8 LIQUID GELS

TOTAL **72** LIQUID GELS*

*LIQUID FILLED CAPSULES



TYLENOL COLD PLUS FLU MULTI-SYMPOTM LIQUID GELS DAY PLUS NIGHT

acetaminophen, dextromethorphan hydrobromide and doxylamine succinate kit

Product Information

| | | | |
|---------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:50580-141 |
|---------------------|----------------|---------------------------|---------------|

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:50580-141-03 | 3 in 1 PACKAGE | 06/30/2025 | |
| 1 | | 1 in 1 KIT; Type 1: Convenience Kit of Co-Package | | |

Quantity of Parts

| Part # | Package Quantity | Total Product Quantity |
|--------|------------------|------------------------|
| Part 1 | 2 BLISTER PACK | 16 |
| Part 2 | 1 BLISTER PACK | 8 |

Part 1 of 2

TYLENOL COLD PLUS FLU MULTI-SYMPATOM LIQUID GELS DAY

acetaminophen, dextromethorphan hydrobromide capsule, liquid filled

Product Information

Item Code (Source) NDC:50580-130

Route of Administration ORAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|----------------------------------|----------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg |
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| GELATIN (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| MINERAL OIL (UNII: T5L8T28FGP) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| POVIDONE (UNII: FZ989GH94E) | |
| WATER (UNII: 059QF0KO0R) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |

Product Characteristics

| | | | |
|----------|--------|--------------|----------|
| Color | orange | Score | no score |
| Shape | OVAL | Size | 21mm |
| Flavor | | Imprint Code | TY;DT |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|--|----------------------|--------------------|
| 1 | | 2 in 1 CARTON | | |
| 1 | | 8 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 09/25/2024 | |

Part 2 of 2

TYLENOL COLD PLUS FLU MULTI-SYMPTOM GELS NIGHT

acetaminophen, dextromethorphan hydrobromide and doxylamine succinate capsule, liquid filled

Product Information

| | |
|-------------------------|---------------|
| Item Code (Source) | NDC:50580-131 |
| Route of Administration | ORAL |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|----------|
| DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL) | DOXYLAMINE SUCCINATE | 6.25 mg |
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 15 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| GELATIN (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| WATER (UNII: 059QF0KO0R) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| MINERAL OIL (UNII: T5L8T28FGP) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| POVIDONE (UNII: FZ989GH94E) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |

Product Characteristics

| | | | |
|----------|-------|--------------|----------|
| Color | green | Score | no score |
| Shape | OVAL | Size | 21mm |
| Flavor | | Imprint Code | TY;NT |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|---------------------|----------------------|--------------------|
|---|-----------|---------------------|----------------------|--------------------|

| | | | |
|----------|--|--|--|
| 1 | 1 in 1 CARTON | | |
| 1 | 8 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|---|-----------------------------|---------------------------|
| OTC Monograph Drug | M012 | 09/25/2024 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|---|-----------------------------|---------------------------|
| OTC Monograph Drug | M012 | 06/30/2025 | |

Labeler - Kenvue Brands LLC (118772437)

Revised: 2/2026

Kenvue Brands LLC