

**EQUALINE NIGHTTIME COLD AND FLU RELIEF- acetaminophen,  
dextromethorphan hbr, doxylamine succinate solution  
United Natural Foods, Inc. dba UNFI**

-----  
**SuperValu Inc. Nighttime Cold & Flu Relief Drug Facts**

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

## Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

## Other information

- **each 30 mL contains:** sodium 38 mg
- store at 20-25°C (68-77°F)

## Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

## Questions?

**1-855-423-2630**

## Package/Label Principal Display Panel

compare to Vicks<sup>®</sup> NyQuil<sup>®</sup> Cold & Flu active ingredients

EQUALINE<sup>®</sup>

nighttime cold & flu relief

acetaminophen (pain reliever/fever reducer)

dextromethorphan HBr (cough suppressant)

doxylamine succinate (antihistamine)

powerful nighttime relief

relieves:

- aches & fever

- runny nose
- sneezing
- cough

original flavor

ALCOHOL 10%

12 FL OZ (355 mL)

compare to Vicks®  
NyQuil® Cold & Flu  
active ingredients\*

NDC 41163-335-40

**EQUALINE®**

# nighttime cold & flu relief

## acetaminophen

(pain reliever/fever reducer)  
dextromethorphan HBr  
(cough suppressant)  
doxylamine succinate  
(antihistamine)

powerful nighttime relief

relieves:

- aches & fever
- runny nose
- sneezing
- cough

original flavor  
ALCOHOL 10%

12 FL OZ (355 mL)

: 33540 EL F7

**DO NOT USE IF  
PRINTED NECKBAND  
IS BROKEN OR MISSING**

Distributed by UNFI  
Providence, RI 02908 USA  
Gluten Free

**100% Quality  
GUARANTEED**

### PARENTS:

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

Like it or let us make it right.  
That's our quality promise.  
855-423-2630

### Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/ fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

**Uses** temporarily relieves common cold/flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose and sneezing

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**PEEL BACK AT  
CORNER FOR MORE  
INFORMATION**

: 33540 EL B1

### Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble

ADHESIVE AREA  
• NO VARNISH • NO TYPE

### Drug Facts (continued)

■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed – see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor

ADHESIVE AREA  
• NO VARNISH • NO TYPE

**AD**  
**NO COATING**

urinating due to an enlarged prostate gland ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema ■ a sodium-restricted diet

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks

**NO COA**

children under 4 yrs do not use

**Other information** ■ each 30 mL contains: sodium 38 mg ■ store at 20-25°C (68-77°F)

**Inactive ingredients**  
alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions?** 1-855-423-2630

\*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Cold & Flu.

## EQUALINE NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41163-335
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	

## Product Characteristics

<b>Color</b>	GREEN (clear, bright green)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	FRUIT (anise / cooling menthol aroma)	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-335-38	296 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/17/2011	01/17/2014
2	NDC:41163-335-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2012	
3	NDC:41163-335-34	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/28/2014	02/04/2017

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/17/2011	

**Labeler** - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 10/2024

United Natural Foods, Inc. dba UNFI