HEALTH MART FAMOTIDINE- famotidine tablet, film coated Strategic Sourcing Services LLC

McKesson Famotidine Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10
 to 60minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- · read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- · protect from moisture

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Maximum Strength Pepcid® AC active ingredient

MAXIMUM STRENGTH

Famotidine Tablets, 20 mg

Acid Reducer

Just One Tablet Prevents & Relieves Heartburn Due to Acid Indigestion

ACTUAL SIZE



DO NOT USE IF PRINTED BLISTER UNIT IS BROKEN OR OPEN OTHER END

Drug Facts

Active ingredient (in each tablet) Famotidine 20 mg.

Purpose

Acid reduce

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn as sociated with acid indigestion and sour stomach brought on by eating or drinking certainfood and beverages

Warnings

Allergy a lert: Do not use if you are allergic to famotidine orotheracid reducers

- if you have trouble or pain swallowing food, vomiting with blood, or bbody or black stools. These may be signs of a serious condition. See your doctor.
- withotheracid reducers

Ask adoctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious con dition.
- heartburn with lightheaded ness, sweating, or dizziness
- chest painor shoulder pain with shortness of breath; sweating; pain spreading to arms, neckors hould ers; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight bss
- nausea orvomiting stomachpain kidney disease

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doc tor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnantor breast-feeding, ask a health professional

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

■ adults and children 12 years and over:

- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn ■ do not use more than 2 tablets in 24 hours
- children u nder 12 years: ask a doctor

Other information

- read the directions and warnings before use
 keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients carnauba wax, colbidal silicon dioxide, croscarmellos e sodium, lactos e monohy drate, magnesium stearate, microcry stalline cellulose, polyethy lene glycol, polyv in yl alcohol, talc,

Tips for Managing Heartburn

• Do not lie flator bend over after eating • Do not wear tight-fitting clothing around the stomach

• Do not lie flator bend over after eating • Do not wear tight-fitting clothing around the stomach

• Do not ast before bedtime - Raises the head of your hed • Avoid heartburn-causing foods
such as rich, spicy, fattly or friedfoods, chocolate, caffeine, alcohol, and certain fruits and
vegetables • Eat slowly and avoid big meals • If overweight, lose weight • Out smoking

CONVENIENT RECLOSING TAB



Drug Facts (continued)

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Maximum Strength Pepcid® AC.

M⊆KESSON

Another Quality Product Distributed by McKesson 6555 State Highway 161, Irving, TX 75039 Money Back Guarantee healthmart.com



HEALTH MART FAMOTIDINE

famotidine tablet, film coated

Product Information

HUMAN OTC DRUG NDC:62011-0143 **Product Type Item Code (Source)**

ORAL **Route of Administration**

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8) **FAMOTIDINE** 20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	WHITE	Score	no score	
Shape	ROUND	Size	8mm	
Flavor		Imprint Code	L194	
Contains				

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:62011- 0143-1	25 in 1 CARTON	03/20/2012				
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product					
2	NDC:62011- 0143-2	1 in 1 POUCH; Type 0: Not a Combination Product	06/15/2016	06/15/2016			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA077351	03/20/2012		

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 7/2021 Strategic Sourcing Services LLC