

FOOT PAIN RELIEF GEL-PATCH- lidocaine and menthol patch
Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.

Foot Pain Relief Gel-Patch 15×22×30

Active Ingredient

Lidocaine 4% w/w Purpose: Topical anesthetic

Menthol 1% w/w Purpose: Topical analgesic

Purposes

Topical anesthetic (Lidocaine)

Topical analgesic (Menthol)

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, neck, shoulders, elbows.

Warnings

For external use only

When using this product:

- use only as directed
- avoid contact with the eyes, mucous membranes or rashes
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Do not use:

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- If you are allergic to any ingredients of this product

Stop use and ask a doctor if:

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering

- condition worsen
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding,

Ask a health professional before use

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center immediately.

Directions

Adult and Children above 12 years old:

- Clean and dry the affected area
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application

Children under 12 years of age: Consult a doctor

Inactive Ingredients

Dihydroxyaluminium Aminoacetate, Edetate Disodium, Glycerin, Hydroxyacetophenone, Kaolin, L(+)-Tartaric Acid, Mineral oil, Polysorbate 80, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylene Glycol, Sodium Polyacrylate, Titanium Dioxide, Water.

Questions or Comments

supports@sumifunofficial.com

Package label. Principal display panel

Maximum Strength

CUT OPEN HERE

Sumifun®

Foot Pain Relief Gel-Patch

Desensitizes aggravated nerves in the foot and relieves foot pain.

☒ Fast Acting

☒ Numbs Pain Away

☒ Targeted Pain Relief

Lidocaine plus +

1 PATCH
2.52"X4.02"
6.4cm X 10.2cm



Knee



Wrist



Neck



Elbow



Shoulder

Drug Facts

Active ingredient	Purpose
Lidocaine 4%	Topical anesthetic
Menthol 1%	Topical analgesic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, neck, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Apply 1 patch at a time to affected area, not more than 3 to 4 times daily.
- Remove patch from the skin after at most 8-hour application

Children under age of 12: Consult a doctor

Inactive ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Hydroxyacetophenone, Kaolin, L(+)-Tartaric Acid, Mineral oil, Polysorbate 80, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylene Glycol, Sodium Polycrylate, Titanium Dioxide, Water.

Questions or Comments:

support@sumifunofficial.com

Business Cooperation:

business@sumifunofficial.com

Child Resistant Packaging



6 9 2 4 8 2 0 1 2 8 0 6

FOOT PAIN

Relief Gel-Patch

www.sumifunofficial.com
YouTube: @sumifun
Facebook: @sumifunOfficial
Instagram: @sumifunofficial
Twitter: @sumifunOfficial
TikTok: @sumifunofficial

15
PATCHES

Maximum Strength

Sumifun® Foot Pain Relief Gel-Patch

Desensitizes aggravated
nerves in the foot
and relieves foot pain.

- ☒ Fast Acting
- ☒ Numbs Pain Away
- ☒ Targeted Pain Relief



Lidocaine plus +

15 PATCHES
2.52"X4.02"
6.4cm X 10.2cm

Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.
417, Building A, No. 8, Dong Heng San Road,
Zhucun, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC

Usage



Remove patch from
outer package



Partially peel back
protective film



Place on
affected area



Remove remaining
film while
pressing patch to skin

unjinws



Knee



Wrist



Neck



Elbow



Shoulder

Drug Facts

Active ingredient Purpose
Lidocaine 4% Topical anesthetic
Menthol 1% Topical analgesic

Uses

Temporarily relieves muscle soreness and minor joint
pains in the wrist, knees, neck, shoulders, elbows.

Warnings

For external use only

Do not use

- more than 1 patch on your body at a time or with
other topical anesthetics at the same time
- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or
mucosae
- do not bandage tightly
- dispose of used patch in manner that always keeps
product away from children and pets. Used patches
still contain the drug product that can produce serious
adverse effects if a child or pet chews or ingests this
patch.

Stop use and ask a doctor if

- localized skin reactions occur, such as rash, itching,
redness, irritation, pain, swelling and blistering
- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few
days

If pregnant or breast-feeding,
ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.
If swallowed accidentally, get medical help or
contact with Poison Control Center immediately.

Directions

- Adults and Children above 12 years old:
 - Clean and dry the affected area.
 - Tear off the protective film and apply the exposed
part of the patch to the affected area.
 - Carefully remove remaining film while pressing
the patch firmly on the skin.
 - Apply 1 patch at a time to affected area, not more
than 3 to 4 times daily.
 - Remove patch from the skin after at most 8-hour
application.
- Children under age of 12: Consult a doctor

Inactive Ingredients

Dihydroxyaluminum Aminoacetate, Edetate
Disodium, Glycerin, Hydroxyacetophenone,
Kaelin, L(+)-Tartaric Acid, Mineral oil, Polyesterate
80, Polyacrylic Acid, Polyvinylpyrrolidone K90,
Propylene Glycol, Sodium Polycrylate,
Titanium Dioxide, Water.

Questions or Comments:

support@sumifunofficial.com

Business Cooperation:

business@sumifunofficial.com

Child Resistant Packaging





FOOT PAIN RELIEF GEL-PATCH

lidocaine and menthol patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83602-174
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	0.01 g in 1 g
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.04 g in 1 g

Inactive Ingredients				
Ingredient Name				Strength
KAOLIN (UNII: 24H4NWX5CO)				
TARTARIC ACID (UNII: W4888I119H)				
WATER (UNII: 059QF0KO0R)				
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)				
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JNI2J)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
MINERAL OIL (UNII: T5L8T28FGP)				
POLYSORBATE 80 (UNII: 6OZP39ZG8H)				
GLYCERIN (UNII: PDC6A3C0OX)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
POVIDONE K90 (UNII: RDH86HJV5Z)				
HYDROXYACETOPHENONE (UNII: G1L3HT4CMH)				
Product Characteristics				
Color	white	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83602-174-15	15 in 1 BOX	01/24/2026	
1		1 in 1 BAG		
1	NDC:83602-174-01	6 g in 1 PATCH; Type 0: Not a Combination Product		
2	NDC:83602-174-22	22 in 1 BOX	01/24/2026	
2		1 in 1 BAG		
2	NDC:83602-174-01	6 g in 1 PATCH; Type 0: Not a Combination Product		
3	NDC:83602-174-30	30 in 1 BOX	01/24/2026	
3		1 in 1 BAG		
3	NDC:83602-174-01	6 g in 1 PATCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M017	01/24/2026		

Labeler - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Registrant - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Revised: 1/2026

Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.