

CHERRY SORE THROAT LOZENGE- benzocaine lozenge

American Sales

Benzocaine 15 mg
Menthol 3.6 mg

Active ingredient (in each lozenge)

Benzocaine 15 mg

Menthol 3.6 mg

Purpose

Oral anesthetic

Oral anesthetic

Uses temporarily relieves these symptoms:

- occasional minor irritation, pain, sore mouth and sore throat

Warnings

Methemoglobinemia warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use

- for teething
- in children under 2 years of age

Stop use and ask doctor if

- sore throat is severe or irritation, pain or redness persists or worsens
- sore mouth symptoms do not improve in 7 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact Poison Control Center right away.

Directions

adults and children 5 years and older - allow 1 lozenge to dissolve slowly in mouth. May be repeated every 2 hours as necessary or as directed by a doctor.

children under 5 years - do not use

Other information

- store at room temperature
- protect from moisture

Inactive ingredients: Corn starch, FD&C red # 40, glucose syrup, glycerin, medium chain triglycerides, natural and artificial flavors, propylene glycol, soybean oil, sucrose and water.

Questions? Call **1-800-245-2898**, Monday through Friday, 9AM to 4PM EST

No Ink or Coating

No Ink or Coating Lot/Exp Info

CAREONE®
Compare to the active ingredients in Cepacol®
Extra Strength Sore Throat Cherry®

Extra Strength
Sore Throat & Cough Lozenges
Fast, soothing, temporary relief of minor sore throat pain and irritation

Cherry
18 LOZENGES
BENZOCAINE 15.0 mg
MENTHOL 3.6 mg

Cherry

Drug Facts
Active ingredients (in each lozenge)
Benzocaine 15 mg.....Oral anesthetic
Menthol 3.6 mg.....Oral anesthetic

Uses temporarily relieves these symptoms:
■ occasional minor irritation, pain, sore mouth and sore throat

Warnings
Hemoglobinemia warnings: If you experience a serious condition that must be treated promptly because it reduces the amount of oxygen in your blood, this can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: ■ pale, gray, or blue colored skin (cyanosis) ■ immediate medical attention ■ rapid heart rate ■ shortness of breath ■ dizziness or lightheadedness ■ fatigue or lack of energy
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.
Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.
Do not use:
■ for teething ■ in children under 2 years of age
Stop use and ask doctor if:
■ sore throat is severe or irritation, pain or redness persists or worsens ■ sore mouth symptoms do not improve in 7 days
If pregnant or breastfeeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away.

Directions
adults and children 5 years and older: allow 1 lozenge to dissolve slowly in mouth. May be repeated every two hours as necessary or as directed by a doctor
children under 5 years: do not use

Other information ■ store at room temperature ■ protect from moisture

Drug Facts (continued)
Inactive ingredients Corn starch, FD&C Red #40, glucose syrup, glycerin, medium chain triglycerides, natural & artificial flavors, propylene glycol, soybean oil, sucrose and water.

Questions? Call 1-800-245-2898, Monday through Friday, 9AM - 4PM EST

*This product is not manufactured or distributed by Beckett Benckiser owner of the registered trademark Cepacol®.
WARNING: TAMPER-EVIDENT PACKAGING. DO NOT USE IF OUTER BOX OR BUBBLE IS OPEN OR DAMAGED.

FOR PHARMACISTS RECOMMEND

GLUTEN FREE

DISTRIBUTED BY: FOODHOLD U.S.A., LLC
LANDOVER, MD 20785
MADE IN USA OF U.S. AND IMPORTED INGREDIENTS
For product questions or concerns, contact us at 1-877-846-9949
© 2018 Anold Delhaize Licensing Sarl
Quality guaranteed or your money back.

PN-292933

3 41520 00028 7

CHERRY SORE THROAT LOZENGE

benzocaine lozenge

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:41520-043 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------|-----------------|
| MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A) | MENTHOL | 3.6 mg |
| BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) | BENZOCAINE | 15 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---------------------------------|-----------------|
| WATER (UNII: 059QF0KO0R) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|----------|
| Color | red | Score | no score |
| Shape | ROUND | Size | 17mm |
| Flavor | CHERRY | Imprint Code | B |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|---|-----------------------------|---------------------------|
| 1 | NDC:41520-043-18 | 18 in 1 BLISTER PACK; Type 0: Not a Combination Product | 06/22/2016 | |



Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 06/22/2016 | |

Labeler - American Sales (809183973)

Registrant - Bestco Inc. (002149136)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------|---------|-----------|------------------------|
| Bestco Inc. | | 002149136 | manufacture(41520-043) |

Revised: 12/2025

American Sales