

**ANTACID FRUIT CHEWS- calcium carbonate tablet, chewable**  
**American Sales**

-----  
**Calcium Carbonate 750 mg**

**Active ingredient (in each chew)**

Calcium carbonate 750 mg

**Purpose**

Antacid

Uses: Relieves

Heartburn

Sour stomach

Acid indigestion

Upset stomach due to these symptoms

**Warnings**

**Do not use if** you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor or pharmacist before use if you are** now taking a prescription drug. Antacids may interact with certain prescription drugs.

Do not take more than 5 chews in a 24-hour period, or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

When using this product constipation may occur.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Directions**

Adults and children 12 years and older chew and swallow 1 to 2 chews every 2-4 hours as needed. Do not exceed 5 chews in 24 hours. Children under 12 years consult a doctor.

**Other information**

- Each chew contains: calcium 300 mg, magnesium 1 mg
- Contains soy
- Contains FD&C Yellow no. 5 (tartrazine) as a color additive
- Store between 20° to 25°C (68° to 77°F) in a dry place

**Inactive ingredients**

Beeswax, carmine, carnauba wax, citric acid, corn starch, corn syrup, DL-alpha tocopherol, ethyl acetate, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 5 (tartrazine), FD&C yellow no. 6 aluminum lake, gum arabic, hydrogenated coconut oil, medium chain triglycerides, methyl paraben, modified corn

starch, natural and artificial flavors, pregelatinized corn starch, propylene glycol, propyl paraben, phosphoric acid, purified water, shellac, sodium benzoate, sorbic acid, sorbitol, soy lecithin, soybean oil, sucrose, and titanium dioxide.

**Questions? Call 1-800-245-2898.** Monday through Friday, 9AM to 4PM EST

**Drug Facts (continued)**

**Other information**

- Each chew contains: calcium 300mg, magnesium 1mg
- Contains soy
- Contains FD&C Yellow No. 5 (tartrazine) as a color additive
- Store between 20° to 25° C (68° to 77°F) in a dry place

**Inactive ingredients:** Beeswax, carmine, carnauba wax, citric acid, corn starch\*, corn syrup, DL-alpha tocopherol, ethyl acetate, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 5 lake (tartrazine), FD&C yellow no. 6, FD&C yellow no. 6 aluminum lake, gum arabic, hydrogenated coconut oil, medium chain triglycerides\*, methyl paraben, modified corn starch, natural and artificial flavors, pregelatinized corn starch, propylene glycol, propyl paraben, phosphoric acid, purified water, shellac, soybean oil\*, sodium benzoate, sorbic acid, sorbitol, soy lecithin, sucrose, titanium dioxide. \*May contain these ingredients.

**Questions?** Call 1-800-245-2898  
Monday - Friday, 9AM - 4PM EST

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**DISTRIBUTED BY**  
FOODHOLD U.S.A., LLC  
LANDOVER, MD 20785  
1-877-848-9949

©2015 Ahold Licensing Sarl  
Quality guaranteed or your money back.

Flavor Chews® & Chewy not Chalky® are registered trademarks of Bestco Inc.

Compare to the active ingredient as AUKA-SELTZER® RELIEFCHEWS®\*\*

**CAREONE™**

**Antacid**

**Extra Strength Flavor Chews®**

750mg of Calcium Carbonate

**ASSORTED FRUIT**

**RELIEVES:**

- Heartburn
- Sour stomach
- Acid indigestion

**Chewy not Chalky®**

**36 FLAVOR CHEWS**

**Drug Facts**

Active ingredient (in each chew)	Purpose
Calcium carbonate 750 mg	Antacid

**Uses:** Relieves

- Heartburn
- Sour stomach
- Acid indigestion
- Upset stomach due to these symptoms

**Warnings**

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor or pharmacist before use if you are now taking a prescription drug. Antacids may interact with certain prescription drugs.

Do not take more than 5 chews in a 24-hour period, or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

May cause constipation

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

**Directions**

Adults and children 12 years and older chew and swallow 1 to 2 chews every 2-4 hours as needed. Do not exceed 5 chews in 24 hours. Children under 12 years consult a doctor.

## ANTACID FRUIT CHEWS

calcium carbonate tablet, chewable

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41520-035
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK) (CALCIUM CATION - UNII:2M83C4R6ZB)	CALCIUM CARBONATE	750 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SUCROSE</b> (UNII: C151H8M554)	

### Product Characteristics

<b>Color</b>	orange (Orange, Pink and Yellow)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	14mm
<b>Flavor</b>	LEMON (Lemon, Orange and Strawberry)	<b>Imprint Code</b>	FC
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-035-36	36 in 1 BOTTLE; Type 0: Not a Combination Product	06/22/2016	



**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	06/22/2016	

**Labeler** - American Sales (809183973)

**Registrant** - Bestco Inc. (002149136)

### Establishment

Name	Address	ID/FEI	Business Operations
Bestco Inc.		002149136	manufacture(41520-035)

Revised: 1/2024

American Sales