# CLOPHED - chlophedianol hydrochloride, guaifenesin, phenylephrine hydrochloride liquid Portal Pharmaceutical

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### Chlophed

Drug Facts
Active ingredients
(in each 5 mL teaspoonful)
Chlophedianol Hydrochloride 12.5 mg
Guaifenesin 100 mg
Phenylephrine Hydrochloride 5 mg

Cough Suppressant Expectorant Nasal Decongestant

#### Uses

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- helps loosen phelgm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- nasal congestion
- reduces swelling of the nasal passages

#### **Warnings**

Do not exceed recommended dosage.

#### Do not use this product

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

### Ask a doctor before use if you have

- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a

fever, rash or persistent headache. These could be signs of a serious condition.

new symptoms occur

### If pregnant or breast-feeding,

ask a health professional before use.

### Keep out of the reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

#### **Directions**

Do not exceed recommended dosage.

Adults and	2 teaspoonfuls every 6 to 8 hours,
children 12 years	not to exceed 8 teaspoonfuls in a
of age and over:	24 hour period
Children 6 to	1 teaspoonful every 6 to 8 hours,
under 12 years of	not to exceed 4 teaspoonfuls in a
age:	24 hour period
Children under 6 years of age	Consult a doctor

#### Other information

Store at 59° - 86° F(15° - 30° C)

### **Inactive ingredients**

Citric Acid, Glycerin, Orange Vanilla Flavor, propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, and Sorbitol.

#### **Questions? Comments?**

Call 1-787-832-6645

Serious side effects associated with this product may be reported to this number.

### **Product Packaging:**

Packaging below represents the labeling currently used:

Principal display panel and side panel for 118 mL label:

NDC 49963-630-04

Clophed

### Each teaspoonful (5 mL) for oral administration contains:

Chlophedianol HCl	12.5 mg
Guaifenesin	_
Phenylephrine HCl	

### Cough Suppressant/Expectorant/Nasal Decongestant

### Alcohol Free/Dye Free/Gluten Free/Sugar Free

### 4 fl oz (118 mL)


Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

Supplied in a tight, light-resistant container with a child-resistant cap.

See package insert for full prescribing information.

Manufactured for: Portal Pharmaceutical Mayaguez, PR 00680

Rev. 10/09





#### Drug Facts (continued)

 cough due to minor throat and bronchial Irritation ■helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive - nasal congestion reduces swelling of nasal passages

#### Warnings

Do not exceed recommended dosage.

#### Do not use this product

■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this

#### Drug Facts (continued)

#### Ask a doctor before use if you have

■a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema a cough that occurs with too much phiegm (mucus) a heart disease . high blood pressure . thyrold disease adiabetes atrouble urinating due to an enlarged prostate gland

#### Stop use and ask a doctor if

- nervousness dizziness or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be sions of a serious condition
- new symptoms occur

#### Drug Facts (continued)

If pregnant or breast-feeding, ask a ealth professional before use Keep out of the reach of children. in case of accidental overdose, seek professional help or contact a Poison Control Center Immediately.

#### Directions

Do not exceed recommended dosage.

12 years of age and over:

Adults and children 2 teaspoonfuls every 12 years of age 6 to 8 hours, not to exceed 8 teaspoonfuls in a 24 hour period

Children 6 to under 12 years of age:

1 teaspoonful every 6 to 8 hours, not to exceed 4 teaspoonfuls In a 24 hour period Consult a doctor

Children under 6 years of age

Inactive ingredients

Other information

Citric Acid, Glycerin, Orange Vanilla Flavor, Propylene Glycol, Purified Water, Sodium Citrate. Sodium Saccharin, and Sorbitol

Store at 59° - 86° F (15° - 30° C)

Drug Facts (continued)

#### Questions? Comments?

Call 1-787-832-6645 Serious side effects associated with this product may be reported to this number.



### **CLOPHED**

chlophedianol hydrochloride, guaifenesin, phenylephrine hydrochloride liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49963-630
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Chlophedianol Hydrochloride (UNII: 69QQ58998Y) (Chlophedianol - UNII:42C50P12AP)	Chlophedianol Hydrochloride	12.5 mg in 5 mL	
Guaifenesin (UNII: 495W7451VQ) (Guaifenesin - UNII:495W7451VQ)	Guaifenesin	100 mg in 5 mL	
<b>Phenylephrine Hydrochloride</b> (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	5 mg in 5 mL	

Inactive Ingredients			
Ingredient Name	Strength		
Citric Acid (UNII: 2968PHW8QP)			
Glycerin (UNII: PDC6A3C0OX)			
Propylene Glycol (UNII: 6DC9Q167V3)			
Water (UNII: 059QF0KO0R)			
Sodium Citrate (UNII: 1Q73Q2JULR)			
Saccharin Sodium (UNII: SB8ZUX40TY)			
Sorbitol (UNII: 506T60A25R)			

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	ORANGE, VANILLA	Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:49963-630-04	118 mL in 1 BOTTLE			
2	NDC:49963-630-15	15 mL in 1 BOTTLE			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	05/28/2010	

## Labeler - Portal Pharmaceutical (831005199)

Revised: 12/2010 Portal Pharmaceutical