

MQUPIN LIDOCAINE GEL-PATCH- lidocaine gel-patch patch
Zhengzhou Miaoqi Medical Technology Co., Ltd.

83781-014

Active Ingredient

Lidocaine 4%
Menthol 3%

Purpose

Topical Analgesic

Use

For temporary relief of pain

Warnings

For external use only

Do not use

- more than 1 patch at a time
- on wounds or damaged skin
- with a heating pad
- if you have ever had an allergic reaction to this product or any of its ingredients

When Using

- use only as directed
- avoid contact with the eyes, mucous membranes or rashes
- do not bandage tightly

Stop Use

localized skin reactions occur, such as rash, itching, redness, irritation, pain swelling and blistering conditions worsen

- symptoms persist for more than 7 days

■ symptoms clear up and occur again within a few days

Ask Doctor

■ If pregnant or breast-feeding, ask a health professional before use.

■ Children under 12 years of age: consult a doctor

Keep Out Of Reach Of Children

Keep Out Of Reach Of Children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children above 12 years old

- 1.Clean and dry the skin.
- 2.Prepare the adhesive secure patch and peel off its backing.
- 3.Peel off the pain patch's protective film.
- 4.Apply to the target area.
- 5.Press to secure in place.Remove patch from the skin after at most 8 hours of application.

Children under 12 years of age: consult a doctor.

Other information

■ avoid storing product in direct sunlight

■ protect product from excessive moisture

■ store at 20-25°C(68-77°F)

Inactive ingredients

Aluminum Glycinate, Edetate Disodium, Ethylhexylglycerin, Glycerin, Kaolin, Phenoxyethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethylcellulose, Sodium Polyacrylate, Tartaric Acid, Water

Questions

Email:mqupin86@gmail.com

PRINCIPAL DISPLAY PANEL

MOUPIN®

MOUPIN®

Lidocaine Gel-Patch

4% Lidocaine+3% Menthol

Maximum Strength Relief

Lasts Up to 10-12 Hours



10 Patches
3.94" x 5.5" (10cm x 14cm)

Lidocaine Gel-Patch

Drug Facts

Active Ingredient

Lidocaine 4% Topical Analgesic
Menthol 3% Topical Analgesic

Use For temporary relief of pain

Warnings For external use only

Do not use

■ more than 1 patch at a time ■ on wounds or damaged skin ■ with a heating pad ■ if you have ever had an allergic reaction to this product or any of its ingredients

When Using

■ use only as directed ■ avoid contact with the eyes, mucous membranes or rashes
■ do not bandage tightly

Stop Use

■ localized skin reactions occur, such as rash, itching, redness, irritation, pain swelling and blistering conditions worsen
■ symptoms persist for more than 7 days
■ symptoms clear up and occur again within a few days

Ask Doctor

■ If pregnant or breast-feeding, ask a health professional before use.
■ Children under 12 years of age: consult a doctor

Keep Out Of Reach Of Children

■ If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children above 12 years old

1. Clean and dry the skin.
2. Prepare the adhesive secure patch and peel off its backing.
3. Peel off the pain patch's protective film.
4. Apply to the target area.
5. Press to secure in place. Remove patch from the skin after at most 8 hours of application.

Children under 12 years of age: consult a doctor.

Other Information

■ avoid storing product in direct sunlight ■ protect product from excessive moisture
■ store at 20-25°C (68-77°F)

Inactive Ingredients

Aluminum Glycinate, Edetate Disodium, Ethylhexylglycerin, Glycerin, Kaolin, Phenoxethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethylcellulose, Sodium Polyacrylate, Tartaric Acid, Water

Legal Disclaimer: This product is not a medical device and is not intended to replace medical treatment. The described effects are based on user feedback, and individual experiences may vary. Please consult a doctor for any health concerns.

Storage: Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

Shelf Life: 3 Years

Made in China

Questions? Email: mqupin86@gmail.com

MFD: Zhengzhou Miaoqi Medical Technology Co., Ltd.

Date Type: MM/DD/YYYY

LOT:

EXP:

MO.UPIN®

MO.UPIN®

Lidocaine Gel-Patch

Lidocaine Gel-Patch

4% Lidocaine+3% Menthol

Maximum Strength Relief

Lasts Up to 10-12 Hours



12 Patches
3.94" x 5.5" (10cm x 14cm)

Lidocaine Gel-Patch

Drug Facts

Active Ingredient	Purpose
Lidocaine 4%	Topical analgesic
Menthol 3%	Topical Analgesic

Use For temporary relief of pain

Warnings For external use only

Do not use
■ more than 1 patch at a time ■ on wounds or damaged skin ■ with a heating pad ■ if you have ever had an allergic reaction to this product or any of its ingredients

When Using
■ use only as directed ■ avoid contact with the eyes, mucous membranes or rashes
■ do not bandage tightly

Stop Use
■ localized skin reactions occur, such as rash, itching, redness, irritation, pain swelling and blistering conditions worsen
■ symptoms persist for more than 7 days
■ symptoms clear up and occur again within a few days

Ask Doctor
■ If pregnant or breast-feeding, ask a health professional before use.
■ Children under 12 years of age: consult a doctor
Keep Out Of Reach Of Children
■ If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children above 12 years old

1. Clean and dry the skin.
2. Prepare the adhesive secure patch and peel off its backing.
3. Peel off the pain patch's protective film.
4. Apply to the target area.
5. Press to secure in place. Remove patch from the skin after at most 8 hours of application.

Children under 12 years of age: consult a doctor.

Other Information

- avoid storing product in direct sunlight
- protect product from excessive moisture
- store at 20-25°C (68-77°F)

Inactive Ingredients

Aluminum Glycinate, Edetate Disodium, Ethylhexylglycerin, Glycerin, Kaolin, Phenoxethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethylcellulose, Sodium Polyacrylate, Tartaric Acid, Water

Legal Disclaimer: This product is not a medical device and is not intended to replace medical treatment. The described effects are based on user feedback, and individual experiences may vary. Please consult a doctor for any health concerns.

Storage: Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

Shelf Life: 3 Years

Made in China

Questions? Email: mqupin86@gmail.com

MFD: Zhengzhou Miaoqi Medical Technology Co., Ltd.

Date Type: MM/DD/YYYY

LOT:

EXP:

MQUPIN®

MQUPIN®

Lidocaine Gel-Patch

Lidocaine Gel-Patch

4% Lidocaine+3% Menthol

Maximum Strength Relief

Lasts Up to 10-12 Hours



15 Patches
3.94" x 5.5" (10cm x 14cm)

Lidocaine Gel-Patch

Drug Facts

Active Ingredient

Lidocaine 4%	Topical analgesic
Menthol 3%	Topical Analgesic

Use For temporary relief of pain

Warnings

For external use only

Do not use

■ more than 1 patch at a time ■ on wounds or damaged skin ■ with a heating pad ■ if you have ever had an allergic reaction to this product or any of its ingredients

When Using

■ use only as directed ■ avoid contact with the eyes, mucous membranes or rashes

■ do not bandage tightly

Stop Use

■ localized skin reactions occur, such as rash, itching, redness, irritation, pain swelling and blistering conditions worsen

■ symptoms persist for more than 7 days

■ symptoms clear up and occur again within a few days

Ask Doctor

■ If pregnant or breast-feeding, ask a health professional before use.

■ Children under 12 years of age: consult a doctor

Keep Out Of Reach Of Children

■ If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children above 12 years old

1. Clean and dry the skin.
2. Prepare the adhesive secure patch and peel off its backing.
3. Peel off the pain patch's protective film.
4. Apply to the target area.
5. Press to secure in place. Remove patch from the skin after at most 8 hours of application.

Children under 12 years of age: consult a doctor.

Other Information

- avoid storing product in direct sunlight
- protect product from excessive moisture
- store at 20-25°C (68-77°F)

Inactive Ingredients

Aluminum Glycinate, Edetate Disodium, Ethylhexylglycerin, Glycerin, Kaolin, Phenoxyethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethylcellulose, Sodium Polyacrylate, Tartaric Acid, Water

Legal Disclaimer: This product is not a medical device and is not intended to replace medical treatment. The described effects are based on user feedback, and individual experiences may vary. Please consult a doctor for any health concerns.

Storage: Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

Shelf Life: 3 Years

Made in China

Questions? Email: mqupin86@gmail.com

MFD: Zhengzhou Miaoqi Medical Technology Co., Ltd.

Date Type: MM/DD/YYYY

LOT:

EXP:

MO.UPIN®

MO.UPIN®

Lidocaine Gel-Patch

4% Lidocaine+3% Menthol

Maximum Strength Relief

Lasts Up to 10-12 Hours

16 Patches

3.94" x 5.5" (10cm x 14cm)

Lidocaine Gel-Patch

Lidocaine Gel-Patch

Drug Facts

Active Ingredient	Purpose
Lidocaine 4%	Topical analgesic
Menthol 3%	Topical Analgesic

Use

For temporary relief of pain

Warnings

For external use only

Do not use

more than 1 patch at a time on wounds or damaged skin with a heating pad if you have ever had an allergic reaction to this product or any of its ingredients

When Using

use only as directed avoid contact with the eyes, mucous membranes or rashes do not bandage tightly

Stop Use

localized skin reactions occur, such as rash, itching, redness, irritation, pain swelling and blistering conditions worsen symptoms persist for more than 7 days symptoms clear up and occur again within a few days

Ask Doctor

if pregnant or breast-feeding, ask a health professional before use. Children under 12 years of age: consult a doctor

Keep Out Of Reach Of Children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children above 12 years old

1. Clean and dry the skin.

2. Prepare the adhesive secure patch and peel off its backing.

3. Peel off the pain patch's protective film.

4. Apply to the target area.

5. Press to secure in place. Remove patch from the skin after at most 8 hours of application.

Children under 12 years of age: consult a doctor.

Other information

avoid storing product in direct sunlight protect product from excessive moisture store at 20-25°C (68-77°F)

Inactive ingredients

Aluminum Glycinate, Edetate Disodium, Ethylhexylglycerin, Glycerin, Kaolin, Phenoxyethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethylcellulose, Sodium Polyacrylate, Tartaric Acid, Water

Legal Disclaimer:

This product is not a medical device and is not intended to replace medical treatment. The described effects are based on user feedback, and individual experiences may vary. Please consult a doctor for any health concerns.

Storage:

Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

Shelf Life:

3 Years

Made in China

Questions? Email: mqupin86@gmail.com

MFD: Zhengzhou Miaoqi Medical Technology Co., Ltd.

6 979297 400777

Date Type: MM/DD/YYYY

LOT:

EXP:

MO.UPIN®

MO.UPIN®

Lidocaine Gel-Patch

4% Lidocaine+3% Menthol

Maximum Strength Relief

Lasts Up to 10-12 Hours



36 Patches
3.94" x 5.5" (10cm x 14cm)

Lidocaine Gel-Patch

Drug Facts

Active Ingredient	Purpose
Lidocaine 4%	Topical analgesic
Menthol 3%	Topical Analgesic

Use For temporary relief of pain

Warnings For external use only

Do not use ■ more than 1 patch at a time ■ on wounds or damaged skin ■ with a heating pad ■ if you have ever had an allergic reaction to this product or any of its ingredients

When Using ■ use only as directed ■ avoid contact with the eyes, mucous membranes or rashes ■ do not bandage tightly

Stop Use ■ localized skin reactions occur, such as rash, itching, redness, irritation, pain swelling and blistering conditions worsen ■ symptoms persist for more than 7 days ■ symptoms clear up and occur again within a few days

Ask Doctor ■ If pregnant or breast-feeding, ask a health professional before use. ■ Children under 12 years of age: consult a doctor

Keep Out Of Reach Of Children ■ If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children above 12 years old

1. Clean and dry the skin.
2. Prepare the adhesive secure patch and peel off its backing.
3. Peel off the pain patch's protective film.
4. Apply to the target area.
5. Press to secure in place. Remove patch from the skin after at most 8 hours of application.

Children under 12 years of age: consult a doctor.

Other Information

- avoid storing product in direct sunlight ■ protect product from excessive moisture
- store at 20-25°C (68-77°F)

Inactive Ingredients

Aluminum Glycinate, Edetate Disodium, Ethylhexylglycerin, Glycerin, Kaolin, Phenoxyethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethylcellulose, Sodium Polyacrylate, Tartaric Acid, Water

Legal Disclaimer: This product is not a medical device and is not intended to replace medical treatment. The described effects are based on user feedback, and individual experiences may vary. Please consult a doctor for any health concerns.

Storage: Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

Shelf Life: 3 Years

Made in China

Questions? Email: mqupin86@gmail.com

MFD: Zhengzhou Miaoqi Medical Technology Co., Ltd.

Date Type: MM/DD/YYYY

LOT:

EXP:

Lidocaine Gel-Patch

MQUPIN®

MQUPIN®

Lidocaine Gel-Patch

4% Lidocaine+3% Menthol

Maximum Strength Relief

Lasts Up to 10-12 Hours

50 Patches
3.94" x 5.5" (10cm x 14cm)

Lidocaine Gel-Patch

Drug Facts

Active Ingredient

Lidocaine 4%

Menthol 3%

Purpose

Topical analgesic

Topical Analgesic

Use

For temporary relief of pain

Warnings

For external use only

Do not use

more than 1 patch at a time

on wounds or damaged skin

with a heating pad

if you have ever had an allergic reaction to this product or any of its ingredients

When Using

use only as directed

avoid contact with the eyes, mucous membranes or rashes

do not bandage tightly

Stop Use

localized skin reactions occur, such as rash, itching, redness, irritation, pain swelling and blistering conditions worsen

symptoms persist for more than 7 days

symptoms clear up and occur again within a few days

Ask Doctor

If pregnant or breast-feeding, ask a health professional before use.

Children under 12 years of age: consult a doctor

Keep Out Of Reach Of Children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children above 12 years old

Clean and dry the skin.

Prepare the adhesive secure patch and peel off its backing.

Peel off the pain patch's protective film.

Apply to the target area.

Press to secure in place. Remove patch from the skin after at most 8 hours of application.

Children under 12 years of age: consult a doctor.

Other Information

avoid storing product in direct sunlight

protect product from excessive moisture

store at 20-25°C (68-77°F)

Inactive Ingredients

Aluminum Glycinate, Edetate Disodium, Ethylhexylglycerin, Glycerin, Kaolin, Phenoxethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethylcellulose, Sodium Polyacrylate, Tartaric Acid, Water

Legal Disclaimer:

This product is not a medical device and is not intended to replace medical treatment. The described effects are based on user feedback, and individual experiences may vary. Please consult a doctor for any health concerns.

Storage:

Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

Shelf Life:

3 Years

Made in China

Questions? Email: mqupin86@gmail.com

MFD: Zhengzhou Miaoqi Medical Technology Co., Ltd.

Date Type: MM/DD/YYYY

LOT:

EXP:

MQUPIN LIDOCAINE GEL-PATCH			
lidocaine gel-patch patch			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83781-014
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)		MENTHOL	3 g in 100
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)		LIDOCAINE	4 g in 100
Inactive Ingredients			
Ingredient Name			Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
EDETATE DISODIUM (UNII: 7FLD91C86K)			
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)			

DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6)	
KAOLIN (UNII: 24H4NWX5CO)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JN12J)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TARTARIC ACID (UNII: W4888I119H)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83781-014-01	10 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
2	NDC:83781-014-02	12 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
3	NDC:83781-014-03	15 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
4	NDC:83781-014-04	16 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
5	NDC:83781-014-05	36 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
6	NDC:83781-014-06	50 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	01/28/2026	

Labeler - Zhengzhou Miaoqi Medical Technology Co., Ltd. (701762807)

Establishment

Name	Address	ID/FEI	Business Operations
Zhengzhou Miaoqi Medical Technology Co., Ltd.		701762807	manufacture(83781-014)

Revised: 1/2026

Zhengzhou Miaoqi Medical Technology Co., Ltd.