

**MQXUNMI LIDOCAINE PAIN RELIEF PATCH- lidocaine pain relief patch patch
Zhengzhou Miaoqi Medical Technology Co., Ltd.**

83781-015

Active Ingredient

Lidocaine 4%
Menthol 3%

Purpose

Topical Analgesic
wrist, knees, back, neck, hips, shoulders, elbows.

Use

For temporary relief of pain

Warnings

For external use only

Do not use

1. Not for use by pregnant or nursing women, infants, or anyone allergic to any ingredient.
2. Do not use on wounds or damaged skin.
3. Do not use with other patch, ointments, creams, sprays, or liniments.
4. Do not tightly wrap the application area or use local heat sources (such as heating pads) on it.
5. Do not use more than 1 patch at a time.
6. Avoid contact with the eyes.
7. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away

When Using

When Using
use only as directed

- avoid contact with the eyes, mucous membranes or rashes
- do not bandage tightly

Stop Use

- 1.If you experience any signs of worsening or skin damage, such as redness, irritation, swelling, or blistering.
- 2.If symptoms persist for more than 7 days, or if symptoms clear up and then reappear within a few days.
- 3.If you are pregnant or breastfeeding, consult a healthcare professional before use.
- 4.Children under 12 years of age: consult a doctor.

Ask Doctor

- 1.If you experience any signs of worsening or skin damage, such as redness, irritation, swelling, or blistering.
- 2.If symptoms persist for more than 7 days, or if symptoms clear up and then reappear within a few days.
- 3.If you are pregnant or breastfeeding, consult a healthcare professional before use.
- 4.Children under 12 years of age: consult a doctor.

Keep Out Of Reach Of Children

Keep Out Of Reach Of Children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children above 12 years old

- 1.Clean and dry the skin.
- 2.Prepare the adhesive secure patch and peel off its backing.
- 3.Peel off the pain patch's protective film.
- 4.Apply to the target area.
- 5.Press to secure in place.

Children under age of 12: Consult a doctor.

Other information

Store in a clean, dry place outside of direct sunlight. Protect product from excessive moisture.

Inactive ingredients

Aluminum Glycinate, Disodium EDTA, Ethylhexylglycerin, Glycerin, Kaolin, Phenoxyethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethyl Cellulose, Sodium Polyacrylate, Tartaric Acid, Water

Questions

Email:MQXUNMI@gmail.com

PRINCIPAL DISPLAY PANEL

230.00 mm

LIDOCaine



LIDOCaine

LIDOCaine Pain Relief Patch

DRUG FACTS

Active Ingredients

Lidocaine 4%.....
Menthol 3%.....

Purpose
Topical Analgesic
Topical Analgesic

Uses

For temporary relief of pain; wrist, knees, back, neck, hips, shoulders, elbows.



Warnings

For external use only.

When Using

Use as directed; avoid contact with the eyes, mucous membranes or rashes, do not bandage tightly. Do not use

1. Not for use by pregnant or nursing women, infants, or anyone allergic to any ingredient.

2. Do not use on wounds or damaged skin.

3. Do not use with other topical medications, creams, sprays, or liniments.

4. Do not tightly wrap the application area or use local heat sources (such as heating pads) on it.

5. Do not use more than 1 patch at a time.

6. Do not swallow.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Stop Use and Ask a Doctor

1. If you experience any signs of worsening or skin damage, such as redness, irritation, swelling, or blisters.

2. If symptoms persist for more than 7 days, or if symptoms clear up and then reappear within a few days.

3. If you are pregnant or breastfeeding, consult a healthcare professional be-fore use.

Directions

Adult and Children above 12 years old

1. Wash and dry the skin.

2. Prepare the patch and secure patch and peel off its backing.

3. Peel off the pain patch's protective film.

4. Apply to the target area.

5. Store in a dry place. Children under age of 12: Consult a doctor.

Inactive Ingredients

Aluminum Glycolate, Disodium EDTA, EthylhexylGlycerin, Glycerin, Kaolin, Phenoxyethanol, Polyvinylpyrrolidone, Polypropylene Glycol, Sodium Carboxymethyl Cellulose, Sodium Polyacrylate, Tartaric Acid, Water

Other information

Store in a clean, dry place outside of direct sunlight. Protect product from excessive moisture.

QUESTION: Who makes MQXUNMI? (Email me)

Legal Disclaimer: This product is not a medical device and is not intended to replace medical treatment. The described effects are based on user feedback, and individual experiences may vary. Please consult a doctor for any health concerns.

Storage: Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

MFID: Zhengzhou Miaoqi Medical Technology Co.,Ltd.

Shelf Life: 3 years

Made in China

Date Type: MM/DD/YYYY

LOT:

EXP:

MQXUNMI®



230.00 mm

LIDOCAINE



LIDOCAINE



双向插口160*45*230mm

230.00 mm



LIDOCAINE

LIDOCaine Pain Relief Patch
DRUG FACTS

Active ingredients
Lidocaine 4%
Menthol 3%
Purpose
Topical Analgesic
Topical Analgesic

Uses
For temporary relief of pain; wrist, knees, back, neck, hips, shoulders, elbows.



Warnings
For external use only.
When Using

1. Do not use for pregnant or nursing women, infants, or anyone allergic to any ingredient.
2. Do not use on wounds or damaged skin.
3. Do not use with other patch, ointments, creams, sprays, or liniments.
4. Do not wrap a patch over a sensitive area or use local heat sources (such as heating pads) on it.
5. Do not use more than 1 patch at a time.
6. Avoid contact with the eyes.

Keep out of reach of children

If you have questions, call a doctor or contact a Poison Control Center right away.

Stop Use and Ask a Doctor

1. If you experience any signs of worsening or skin damage, such as redness, irritation, swelling, or blistering.
2. If symptoms persist for more than 7 days, or if symptoms clear up and then reappear within a few days.
3. If you are pregnant or breastfeeding, consult a healthcare professional before use.

4. If you are under 12 years of age, consult a doctor.

Children

Adults and Children above 12 years old

1. Clean and dry the skin.

2. Prepare the adhesive secure patch and peel off its backing.

3. Place the patch on the skin and apply protective film.

4. Apply to the target area.

5. Press to secure in place. Children under age of 12: Consult a doctor.

Inactive Ingredients

Aluminum Chloride, Sodium EDTA, Ethyleneglycol, Glycerin, Kaolin, Phenoxethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethyl Cellulose, Sodium Polyacrylate, Tartaric Acid, Water

Other Information

Store in a clean, dry place outside of direct sunlight. Protect product from excessive moisture.

QUESTIONS? Email: MQXUNMI@gmail.com

Legal Disclaimer: This product is not a medical device and is not intended to replace medical treatment.

For personal use only. Individual results may vary. Please consult a doctor for any health concerns.

Storage: Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

MD: Zhengzhou Miaoqi Medical Technology Co.,Ltd.

Shelf Life: 3 years

Made in China

Date Type: MM/DD/YYYY

LOT:

EXP:

6 977395 530235



MQXUNMI®

230.00 mm

LIDOCaine



LIDOCaine

LIDOCaine Pain Relief Patch

DRUG FACTS

Active Ingredients

Lidocaine 4%..... Purpose Topical Analgesic
Menthol 3%..... Topical Analgesic

Uses

For temporary relief of pain; wrist, knees, back, neck, hips, shoulders, elbows.



Warnings

For external use only.

Stop Use and Ask a Doctor

Use only as directed. Avoid contact with the eyes, mucous membranes or rashes, do not bandage tightly. 1.Do not use by pregnant or nursing women, infants, or anyone allergic to any ingredient.

2.Do not use on wounds or damaged skin.

3.Do not use with other patch, ointments, creams, sprays, or liniments.

4.Do not tightly wrap the application area or use local heat sources (such as heating pads) on it.

5.Do not use more than 1 patch at a time.

6.Do not reuse.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Stop Use and Ask a Doctor

1.If you notice any signs of worsening or skin damage, such as redness, irritation, swelling, or blistering.

2.If symptoms persist for more than 7 days, or if symptoms clear up and then reappear within a few days.

3.If you are pregnant or breastfeeding, consult a healthcare professional be-fore use.

4.Children under 12 years of age: consult a doctor.

Directions

Adults and Children above 12 years old

1.Clean and dry the skin.

2.Prepare the adhesive secure patch and peel off its backing.

3.Peel off the pain patch's protective film.

4.Apply to the target area.

5.Remove after 8 hours. Children under age of 12: Consult a doctor.

Storage

Store in a clean, dry place outside of direct sunlight. Protect product from excessive moisture.

QUESTIONS? Email: MQXUNMI@gmail.com

Legal Disclaimer: This product is not a medical device and is not intended to replace medical treatment. The described effects are based on user feedback, and individual experiences may vary. Please consult a doctor for any health concerns.

Storage: Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

MFPI: Zhengzhou Miaoqi Medical Technology Co.,Ltd.

Shelf Life: 3 years

Made In China

Date Type: MM/DD/YYYY

LOT:

EXP:

MQXUNMI®

230.00 mm

LIDOCAINE



LIDOCAINE

LIDOCaine Pain Relief Patch

DRUG FACTS

Active Ingredients	Lidocaine 4% Menthol 3%	Purpose	Topical Anesthetic Topical Analgesic
--------------------	----------------------------	---------	---

Uses For temporary relief of pain in wrist, knees, back, neck, hips, shoulders, elbows.



Warnings

For external use only. When Using Use only as directed. Avoid contact with the eyes, mucous membranes or rashes, do not bandage tightly. Overdosage

1. Not for use by pregnant or nursing women, infants, or anyone allergic to any ingredient.
2. Do not use on wounds or damaged skin.
3. Do not use with other patch, ointments, creams, sprays, or liniments.
4. Do not tightly wrap pain patch over area or use local heat sources (such as heating pads) on it.
5. Do not use more than 1 patch at a time.
6. Avoid contact with the eyes.

Keep out of reach of children

If swallowed, call a doctor or contact a Poison Control Center right away.

Stop Use and Ask a Doctor

1. If you experience any signs of worsening or skin damage, such as redness, irritation, swelling, or blistering.
2. If symptoms persist for more than 7 days, or if symptoms clear up and then reappear within a few days.

3. If you are pregnant or breastfeeding, consult a healthcare professional before use.

4. If under 12 years of age, consult a doctor.

Directions

Adults and Children above 12 years old

1. Clean and dry the skin.
2. Remove the adhesive secure patch and peel off its backing.
3. Stick the patch onto the adhesive secure patch.
4. Peel off the pain patch's protective film.

5. Apply to the target area. 6. Press to secure in place. Children under age of 12: Consult a doctor.



Inactive Ingredients

Aluminum Glycolate, Disodium EDTA, Ethylhexylglycerin, Glycerin, Kaolin, Phenoxethanol, Polyvinylpyrrolidone, Propylene Glycol, Sodium Carboxymethyl Cellulose, Sodium Polyacrylate, Tartaric Acid, Water

Other Information

Store in a clean, dry place outside of direct sunlight. Protect product from excessive moisture.

QUESTIONS? Email: MQXUNMI@gmail.com

Legend: This product is a medical device and is not intended to replace medical treatment. The described effects are based on user feedback, and individual experiences may vary. Please consult a doctor for any health concerns.

Storage: Store in a clean, dry place away from direct sunlight. Protect from excessive moisture.

MF: Zhengzhou Miaoqi Medical Technology Co.,Ltd.

Shelf Life: 3 years

Made in China

Date Type: MM/DD/YYYY

LOT:

EXP:

MQXUNMI®

MQXUNMI LIDOCaine PAIN RELIEF PATCH

lidocaine pain relief patch patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83781-015
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	3 g in 100
LIDOCaine (UNII: 98PI200987) (LIDOCaine - UNII:98PI200987)	LIDOCaine	4 g in 100

Inactive Ingredients

Ingredient Name	Strength
DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6)	
GLYCERIN (UNII: PDC6A3C00X)	
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)	
POVIDONE K90 (UNII: RDH86HJV5Z)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JN12J)

METHYLPARABEN (UNII: A2I8C7HI9T)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

POLYSORBATE 80 (UNII: 6OZP39ZG8H)

TARTARIC ACID (UNII: W4888I119H)

WATER (UNII: 059QF0KO0R)

KAOLIN (UNII: 24H4NWX5CO)

EDETALE DISODIUM (UNII: 7FLD91C86K)

PROPYLPARABEN (UNII: Z8IX2SC1OH)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83781-015-01	10 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
2	NDC:83781-015-02	12 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
3	NDC:83781-015-03	15 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
4	NDC:83781-015-04	16 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
5	NDC:83781-015-05	36 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	
6	NDC:83781-015-06	50 in 1 BOX; Type 0: Not a Combination Product	01/28/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	01/28/2026	

Labeler - Zhengzhou Miaoqi Medical Technology Co., Ltd. (701762807)

Establishment

Name	Address	ID/FEI	Business Operations
Zhengzhou Miaoqi Medical Technology Co., Ltd.		701762807	manufacture(83781-015)

Revised: 1/2026

Zhengzhou Miaoqi Medical Technology Co., Ltd.