

ATAMEL FORTE- acetaminophen tablet, coated
OPMX LLC

ATAMEL Forte w/24 Caplets

Drug Facts

Active ingredient (in each caplet)

- Acetaminophen 650 mg

Purpose

Pain reliever / Fever Reducer

Uses

- Temporarily relieves minor aches and pains due to:
- Minor pain of arthritis
- Muscular aches
- Backache
- Premenstrual and menstrual cramps
- The common cold
- Headache
- Toothache
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- More than 6 caplets in 24 hours, which is the maximum daily amount
- With other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- Skin reddening
- Blisters
- Rash

If skin reaction occurs, stop use and seek medical help right away.

Do not use

- With any other containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- Pain gets worse or lasts more than 10 days
- Fever gets worse or lasts more than 3 days
- New symptoms occur
- Redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children

Keep out of reach of children

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **Do not take more than directed (see overdose warning)**

Adults	<ul style="list-style-type: none">• Take 2 caplets every 8 hours with water• swallow whole; do not crush, chew, split or dissolve• do not take more than 6 caplets in 24 hours• do not use for more than 10 days unless directed by doctor
Under 18 years of age	<ul style="list-style-type: none">• Ask a doctor

Other information

- Store between 20-25°C (60-77°F)
- Do not use if carton is opened.
- Do not use if foil inner seal is broken or missing.

Inactive ingredients

caranuba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin.

Questions or comments?

Call **(619)600-5632** (Mon-Fri 9am - 5pm PST)

Made for:

San Diego, CA 92154

Exclusively distributed by:

OPMX

Chula Vista, CA 91910

Phone: 619-600-5632

NDC 69729-800-24

Acetaminophen 650 mg

Atamel®

Via de Administracion: Oral

Forte

Alivia el Dolor

Pain Relief

Reduce la fiebre

Fever Reducer

Dolores Musculares

Muscles Aches

Dolor de cabeza

Headache

Dolor leve por Artritis

Minor Pain of Arthritis

24 Caplets



Drug Facts
Active ingredient (in each caplet)
 Acetaminophen 650 mg.....Pain reliever / Fever Reducer

Purpose
 Acetaminophen 650 mg.....Pain reliever / Fever Reducer

Uses
 ■ Temporarily relieves minor aches and pains due to:
 ■ Minor pain of arthritis
 ■ Muscular aches
 ■ Backache
 ■ Premenstrual and menstrual cramps
 ■ The common cold
 ■ Headache
 ■ Toothache
 ■ Temporarily reduces fever

Warnings - This product contains acetaminophen. Severe liver damage may occur if you take
 ■ More than 6 caplets in 24 hours, which is the maximum daily amount
 ■ With other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product.
Allergy alert: Acetaminophen may cause severe skin reactions.
 Symptoms may include:
 ■ Skin redness ■ Blisters ■ Rash
 If skin reaction occurs, stop use and seek medical help right away.

Do not use
 ■ With any other containing acetaminophen (prescription or

Drug Facts (continued)
 Stop use and ask a doctor if
 ■ Pain gets worse or lasts more than 10 days
 ■ Fever gets worse or lasts more than 3 days
 ■ New symptoms occur
 ■ Redness or swelling is present
 These could be signs of serious condition.

Keep out reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 ■ Do not take more than directed (see overdose warning)

Adults
 Take 2 caplets every 8 hours with water
 swallow whole, do not crush, chew, split or dissolve
 do not use for more than 10 days unless directed by a doctor

Under 18 years of age
 Ask a doctor

Other information
 ■ Store between 20-25°C (68-77°F)

Acetaminophen 650 mg
Atamel®
 Vía de Administración: Oral **Forte**

NDC 69729-800-24
 Acetaminophen 650 mg
Atamel®
 Vía de Administración: Oral **Forte**

Pain Relief
 Alivia el Dolor

- | | |
|--|---|
| Fever Reducer
Reduce la Fiebre | Muscles Aches
Dolores Musculares |
| Headache
Dolor de Cabeza | Minor Pain of Arthritis
Dolor leve por Artritis |

24 Caplets

Acetaminophen 650 mg
Atamel®
 Vía de Administración: Oral **Forte**

Lot No:
 Exp Date:

NO VANISH AREA

TAMPER EVIDENT:
 Do not use if package is opened or if blister unit is torn, broken or shows any sign of tampering

REV 12.2025 ITEM RRX1159

 8 10053 18166 9



Made for:

 San Diego, CA 92154

Exclusively distributed by:

 Chula Vista, CA 91910
 Phone: 619-600-5632

Drug Facts
Active ingredient (in each caplet) **Purpose**
 Acetaminophen 650 mg.....Pain reliever / Fever Reducer

Uses
 ■ Temporarily relieves minor aches and pains due to:
 ■ Minor pain of arthritis
 ■ Muscular aches
 ■ Backache
 ■ Premenstrual and menstrual cramps
 ■ The common cold
 ■ Headache
 ■ Toothache
 ■ Temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ More than 6 caplets in 24 hours, which is the maximum daily amount
 ■ With other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ Skin reddening ■ Blisters ■ Rash
 If skin reaction occurs, stop use and seek medical help right away.

Do not use
 ■ With any other containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ If you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Drug Facts (continued)
Stop use and ask a doctor if
 ■ Pain gets worse or lasts more than 10 days
 ■ Fever gets worse or lasts more than 3 days
 ■ New symptoms occur
 ■ Redness or swelling is present
 These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out reach of children.**
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 ■ **Do not take more than directed (see overdose warning)**

Adults	Take 2 caplets every 8 hours with water swallow whole; do not crush, chew, split or dissolve do not take more than 6 caplets in 24 hours do not use for more than 10 days unless directed by a doctor
Under 18 years of age	Ask a doctor

Other information
 ■ Store between 20-25°C (68-77°F)
 ■ Do not use if carton is opened.
 ■ Do not use if foil inner seal is broken or missing.

Inactive ingredients caranuba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin.

Questions or comments?
 Call (619)600-5632 (Mon- Fri 9am - 5pm PST)

NO VANISH AREA

Datos del medicamento
Ingrediente activo (en cada cápsula) **Propósito**
 Acetaminofén 650 mg Alivio del dolor / Reductor de fiebre

Usos
 ■ Alivia temporalmente dolores y molestias menores debido a:
 ■ Dolor leve por artritis
 ■ Dolores musculares
 ■ Dolor de espalda
 ■ Cólicos premenstruales y menstruales
 ■ Resfriado común
 ■ Dolor de cabeza
 ■ Dolor de muelas
 ■ Reduce temporalmente la fiebre

Advertencias
Advertencia sobre el hígado: Este producto contiene acetaminofén. Puede ocurrir daño hepático severo si usted toma:
 ■ Más de 6 cápsulas en 24 horas, que es la cantidad máxima diaria
 ■ Con otros medicamentos que contengan acetaminofén
 ■ 3 o más bebidas alcohólicas cada día mientras usa este producto
Alerta de alergia: El acetaminofén puede causar reacciones severas en la piel. Los síntomas pueden incluir:
 ■ Enrojecimiento de la piel ■ Ampollas ■ Erupción cutánea
 Si ocurre una reacción en la piel, suspenda su uso y busque ayuda médica de inmediato.

No usar
 ■ Con ningún otro producto que contenga acetaminofén (de prescripción o sin receta). Si no está seguro de si un medicamento contiene acetaminofén, consulte a un médico o farmacéutico.
 ■ Si es alérgico al acetaminofén o a alguno de los ingredientes inactivos en este producto

Consulte a un médico antes de usar si tiene enfermedad hepática
Consulte a un médico o farmacéutico antes de usar si está tomando el medicamento anticoagulante warfarina

Datos del medicamento (continuación)
Suspenda el uso y consulte a un médico si
 ■ El dolor empeora o dura más de 10 días
 ■ La fiebre empeora o dura más de 3 días
 ■ Aparecen nuevos síntomas
 ■ Hay enrojecimiento o hinchazón
 Estos podrían ser signos de una condición seria.

Si está embarazada o amamantando, consulte a un profesional de salud antes de usar. **Mantener fuera del alcance de los niños.**
Advertencia de sobredosis: en caso de sobredosis, obtenga ayuda médica o contacte a un Centro de Control de Envenenamientos de inmediato (1-800-222-1222). La atención médica rápida es crucial tanto para adultos como para niños, incluso si no nota ningún signo o síntoma.

Instrucciones
 ■ **No tome más de lo indicado (ver advertencia de sobredosis)**

Adultos	Tome 2 cápsulas cada 8 horas con agua tráguelas enteras; no las triture, mastique, parta ni disuelva no tome más de 6 cápsulas en 24 horas no use por más de 10 días a menos que lo indique un médico
Menores de 18 años	Consulte a un médico

Otra información
 ■ Almacene entre 20-25°C (68-77°F)
 ■ No use si el empaque está abierto.
 ■ No usar si el sello interior de aluminio está roto o ausente.

Ingredientes inactivos Cera de carnuba, hidroxietil celulosa, hipromelosa, estearato de magnesio, celulosa microcristalina, almidón modificado, povidona, celulosa pulverizada, almidón pregelatinizado, glicolato de almidón sódico, dióxido de titanio, triacetina.

¿Preguntas o comentarios?
 Llame al (619)600-5632 (Mon- Fri 9am - 5pm PST)

ATAMEL FORTE

acetaminophen tablet, coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:69729-800

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POVIDONE K30 (UNII: U725QWY32X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TRIACETIN (UNII: XHX3C3X673)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL (Oblong)	Size	21mm
Flavor		Imprint Code	OPMX
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-800-24	2 in 1 CARTON	01/29/2026	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/29/2026	

Labeler - OPMX LLC (029918743)

Establishment

Name	Address	ID/FEI	Business Operations
HBC HEALTHCARE PRIVATE LIMITED		854255181	manufacture(69729-800)

Revised: 1/2026

OPMX LLC